

## Spatial Hierarchical Bayes Small Area Inference

**Priyanka Anjoy**

*Assistant Director, National Accounts Division (NAD), National Statistical Office (NSO)  
Ministry of Statistics and Program Implementation (MOSPI), Government of India*

Received: 16 August 2022; Revised: 25 August 2022; Accepted: 28 August 2022

---

### Abstract

The Hierarchical Bayes predictor of small area proportions (HBP) under an area level version of generalized linear mixed model with logit link function is widely used in small area estimation for binary variable. However, this predictor does not account for the presence of spatial effect between contiguous or neighbouring regions. Conditional Autoregressive and Simultaneous Autoregressive specifications do incorporate spatial association while considering the spatial correlation effects among areas. But none of these approaches implement the idea of spatially varying covariates through spatially dependent fixed effect parameters. Such approach in statistics is known as spatial nonstationarity. This article introduces a spatially nonstationary extension to the Hierarchical Bayes predictor of small area proportions that accounts for the presence of spatial nonstationarity. The proposed predictor is referred as the spatial nonstationary Hierarchical Bayes predictor (HBNSP). The impact of survey design information is also explored in the proposed predictor. The empirical results from simulation studies using spatially nonstationary data indicate that the HBNSP method performs better, in terms of relative bias and relative mean squared error, than the alternative HBP method that ignore this spatial nonstationarity. The results further show that use of survey-weight to incorporate the sampling design appears to be imperative when sample data is informative.

*Key Words:* Hierarchical Bayes, Small area estimation, Spatial nonstationarity, Survey-weight.

---

### 1. Introduction

In recent years, small area estimation (SAE) technique has emerged as one of the most important topics in survey estimation because of an increasing demand for reliable small area statistics by various government and international agencies, see for example, Rao and Molina (2015). United Nations Sustainable Development Agenda has also marked the developmental strategy through availing and utilizing disaggregate level statistics in the programmes and planning aimed at uprooting social and regional inequalities. Sample surveys are generally designed so that direct estimators (*i.e.* estimators that use only the sample data from the domain of interest) for larger domains provide reliable estimates for parameters of interest. On many occasions, however, the interest is in estimating parameters for domains that contain only a small number of sample observations or sometimes no sample observations. The term ‘small areas’ is used to describe domains whose sample sizes are not large enough to allow sufficiently

precise direct estimation. Hereafter, refer to these smaller domains as 'small areas' or simply 'areas'. When direct estimation is not possible, one must rely on alternative, model-based methods for producing small area estimates. Further, large scale surveys produce reliable estimates at higher geographical level and such estimates often mask variations which is available at local levels. This restricts targeting of heterogeneity at higher levels of spatial disaggregation and limits the scope for monitoring and evaluation of parameters locally within and across administrative units. Model-based SAE techniques are now widely used in practice to meet the indispensable need of reliable disaggregate level statistics from the existing survey data. Such SAE methods depend on the availability of population level auxiliary information related to the variable of interest and are commonly referred to as indirect methods. The industry standard for SAE is to use unit or area level models (Fay and Herriot, 1979; Battese, Harter and Fuller, 1988). In the former case these models are for the individual survey measurements and include area effects, while in the latter case these models are used to smooth out the variability in the unstable area-level direct estimates. Area-level small area modeling is usually employed when unit-level data are unavailable, or, as is often the case, where model covariates (*e.g.* census variables) are only available in aggregate form. In this article solely focus is on area (or aggregated) level small area modeling.

Fay-Herriot (FH) model is one of the popular examples of aggregated level small area model. For continuous survey variable, this model is widely used in practice and has led the phenomenal development of small area literatures based on this model. However, binary or count data is often of interest in many practical applications. In epidemiological, environmental, poverty related studies such data is much common, where interest generally lies in estimation of proportions. A generalized linear mixed model (GLMM) with logit link function (also referred to as logistic linear mixed model) is commonly used for estimation of small area proportions. The basic structure of area level small area models include sampling model for direct survey estimates and associated sampling error; linking model to link the parameter of interest with area-specific auxiliary variables and random effects. The area random effect in small area models explains unstructured heterogeneity between areas. Two basic approaches for drawing inferences about the small area parameters of interest are known to be popular: The empirical best prediction method is based on frequentist idea to estimate unknown model parameters and the hierarchical Bayes (HB) approach assumes particular prior distributions for the hyperparameters to obtain posterior quantities of the parameter of interest. The HB approach has the flexibility to deal with complex SAE model as it overcomes the difficulties of analytical mean squared error (MSE) estimation in frequentist set up and provides quick and easier posterior variance computation based on Markov Chain Monte Carlo (MCMC) simulation. Refer Jiang and Lahiri (2001), You and Zhou (2011), Liu *et al.* (2014), Rao and Molina (2015) and Chandra *et al.* (2018) for frequentist and Bayesian related studies and various real life applications. This article in particular focuses on estimation of small area proportions in hierarchical Bayes framework. Among the previous literatures, Liu *et al.* (2014) and Anjoy *et al.* (2019) have applied hierarchical Bayes version of GLMM (HBGLMM) to estimate survey-weighted small area proportions considering different cases of known and unknown sampling variance structure (denoted by HBP). The linking model of HBP incorporates random effect which is assumed to be independent and identically distributed. As a result, spatial association between geographical areas cannot be described through this structure of the model. However, in many small area problems like disease prevalence and poverty estimation, spatial contiguity between neighbouring areas is very common. Therefore, induction of spatial variability in GLMM can be a way of reducing the variances or Coefficient of Variation (CV) in final estimates. One approach to incorporating such spatial dependency among the areas is to extend the GLMM to allow for spatially correlated area effects using, for

example, a Simultaneous Autoregressive (SAR) model (Cressie, 1993). This model allows for spatial correlation in the area effects, while keeping the fixed effects parameters spatially invariant (Chandra and Salvati, 2018). There are data situations, where this assumption is inappropriate and parameters associated with the model covariates (*i.e.* the fixed effects parameters) vary spatially. This phenomenon is often referred to as spatial nonstationarity (Brunsdon *et al.*, 1996). An alternative approach to incorporating spatial information in SAE is therefore to assume that the parameters associated with the model covariates vary spatially. In frequentist framework, Chandra *et al.* (2017) has devised the concept of spatial nonstationarity in area level version of GLMM (NSGLMM) for estimating small area proportions. A key feature of this approach is that it tries to capture spatial variability through incorporating spatially varying covariates in the linking model. It is worth noting that Chandra *et al.* (2017) approach does not use the sampling weights or clustering information in estimation of small area proportions under NSGLMM. However, use of this sampling information is essential for valid inference from survey data collected by complex survey designs. Baldermann *et al.* (2018) has also forwarded spatial nonstationarity concept for explaining spatial variability between areas, but their model is for unit-level data. Contrary to the previous studies, this article describes a spatial nonstationary version of hierarchical Bayes approach for SAE that incorporates the sampling information when estimating small area proportions under an area level small area models (denoted by HBNSP). Unlike frequentist approach, the HBNSP method offers the flexibility of MSE estimation through posterior variance computation based on MCMC simulation.

Standard model-based approaches to the analysis often ignore the sampling mechanism. The GLMM technique implicitly considers equal probability sampling (simple random sampling with replacement) within each small area and thus ignores the survey-weight (Chandra *et al.*, 2019). But, this may result in potentially large biases in the final estimates. In FH model for estimation of small area population mean, direct design-based estimators are modeled directly and the survey variance of the associated direct estimator is introduced into the model via the design-based errors. The Horvitz-Thompson estimator, weighted Hájek estimator are the structures here to incorporate survey design information (Hidiroglou and You, 2016). However, this method for continuous data requires extension for binary or count data for estimating more representative small area proportions. Consequently, the strategic idea is to modeling survey-weighted proportions (Liu *et al.*, 2014). Hence, the next attempt in this article is to check the impact of complex survey design information in HBNSP. In next section, two versions of HBNSP predictor denoted as HBNSP1 and HBNSP2 are described, which respectively account for HB modeling of unweighted and survey-weighted small area proportions. In section 3, empirical evaluation studies first include a model-based simulation set up to evaluate the performance of proposed HBNSP as compared to HBP. Secondly, a design-based simulation study is carried out for comparing the performance of HBNSP1 and HBNSP2 which respectively, ignores and considers the modeling of survey-weighted proportions. The article ends with relevant concluding remarks.

## 2. Methodology

Let us consider a finite population  $U$  of size  $N$  which is partitioned into  $D$  distinct small areas or simply areas. The set of population units in area  $i$  is denoted as  $U_i$  with known size  $N_i$ , such that  $U = \bigcup_{i=1}^D U_i$  and  $N = \sum_{i=1}^D N_i$ . A sample  $s$  of size  $n$  is drawn from population  $U$  using a probabilistic mechanism. This resulted in sample  $s_i$  in area  $i$  with size  $n_i$ , so that  $s = \bigcup_{i=1}^D s_i$  and

$n = \sum_{i=1}^D n_i$ . Assume that  $y_{ij}$  be the value of target variable  $y$  for unit  $j$  ( $j=1, \dots, n_i$ ) in small area  $i$ . The target variable with values  $y_{ij}$  has binary response, taking value either 1 or 0. Our aim is to estimate the small domain proportions  $P_i = N_i^{-1} \sum_{j=1}^{N_i} y_{ij}$ . When the sample  $s$  is drawn following a complex survey design, with each unit  $y_{ij}$  in small area  $i$  design weight  $w_{ij}$  is attached, which is alternatively known as survey-weights or sampling weights.

## 2.1. Estimation of small area proportions

The area level version GLMM is widely used for estimation of small area proportions to improve the precision of direct survey estimates. Consider,  $p_i$  be the direct survey estimator for the parameter of interest  $P_i$ . In aggregated level model, it is customary to assume that,

$$p_i = P_i + e_i; i = 1, \dots, D,$$

where  $e_i$ 's are independent sampling error associated with direct estimator  $p_i$ . Sampling error  $e_i$  is assumed to have zero mean and known sampling variance  $\sigma_{ei}^2$ . The linking model of  $P_i$  attempt to relate area-specific auxiliary variables and random effect component,

$$g(P_i) = \mathbf{x}_i' \boldsymbol{\beta} + v_i; i = 1, \dots, D,$$

where the linking function  $g(\cdot)$  is logit for binary data and log for count data,  $\mathbf{x}_i$  represent matrix of area-specific auxiliary variables,  $\boldsymbol{\beta}$  is the regression coefficient or fixed effect parameter vector and  $v_i$  being the area-specific random effect, independent and identically distributed as  $E(v_i) = 0$  and  $\text{var}(v_i) = \sigma_v^2$ . Random area-specific effects are included in the linking model to account for between areas dissimilarities. Working under HB set up, certain prior distributions are assumed for the hyperparameters. For estimating small area proportions  $P_i$ , the sampling and linking models of HBP are represented as,

$$p_i | P_i \sim N(P_i, \sigma_{ei}^2), i = 1, \dots, D \text{ and } \text{logit}(P_i) | \boldsymbol{\beta}, \sigma_v^2 \sim N(\mathbf{x}_i' \boldsymbol{\beta}, \sigma_v^2), i = 1, \dots, D.$$

Following standard literature, prior choice for  $\boldsymbol{\beta}$  is usually taken to be  $N(0, \sigma_0^2)$  and for  $\sigma_v^2$  prior choice is  $IG(a_0, b_0)$ , ( $IG$  stands for Inverse Gamma) where  $\sigma_0^2$  is set to be very large (say,  $10^6$ ) and very small value for  $a_0$  and  $b_0$  (usually  $a_0 = b_0 \rightarrow 0$ ) to reflect lack of prior knowledge about variance parameters (Rao, 2015; You and Zhou, 2011). Then, inferences about the small area parameter of interest are drawn from posterior distribution. Posterior mean is taken as the point estimate of the parameter and posterior variance as a measure of the uncertainty associated with the estimate. However, an inbuilt postulation in HBP is that fixed effect parameter or regression coefficient vector  $\boldsymbol{\beta}$  is spatially invariant, this is what customarily known as spatial stationarity. In contrary, spatial nonstationarity approach tends to describe/define spatially varying regression parameters, *i.e.*, values of the regression coefficients are necessarily different at different spatial locations. Small area estimation of proportions in presence of such spatial nonstationarity is described in next subsection.

## 2.2. Hierarchical Bayes version of spatial nonstationary GLMM

For spatial nonstationary version of HBGLMM or HBNSP, regression coefficients in the small area model may be expressed as explicit functions of the spatial points of the sample observations. Unlike HBP, where one restrict to a single global model with fixed parameter, HBNSP technique defines local relationships to exist between study and auxiliary variables. This approach is quite like the geographically weighted regression (GWR) in a multiple regression framework which takes nonstationary auxiliary variables into consideration (Brunsdon *et al.*, 1996). Let,  $l_i$  be the coordinates of an arbitrarily defined spatial location (longitude and latitude) for  $i^{\text{th}}$  small area; generally, this will be its centroid. Consider,  $\mathbf{l} = (l_1, \dots, l_D)'$  denoting the  $D$  component vector of such spatial locations *i.e.*, having available longitude and latitude for all the  $D$  spatial locations or areas of interest. Assume that nonstationarity is characterized by an area specific vector of fixed effects,

$$\mathbf{x}'_i \boldsymbol{\beta}(l_i) = \mathbf{x}'_i \boldsymbol{\beta} + \mathbf{x}'_i \boldsymbol{\gamma}(l_i),$$

where  $\boldsymbol{\beta}(l_i) = \boldsymbol{\beta} + \boldsymbol{\gamma}(l_i)$  and  $\boldsymbol{\gamma}(l_i) = (\gamma_1(l_i), \dots, \gamma_p(l_i))'$ . The linking model of  $P_i$  in HBNSP attempt to relate nonstationary auxiliary variables and random effect component,

$$\text{logit}(P_i) = \mathbf{x}'_i \boldsymbol{\beta}(l_i) + v_i; i = 1, \dots, D, \text{ with } v_i \sim N(0, \sigma_v^2).$$

Aggregating  $D$  area level models lead to the population level version of the HBNSP as

$$\mathbf{p} = \mathbf{X}\boldsymbol{\beta}(\mathbf{l}) + \mathbf{v} + \mathbf{e} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\Psi}\boldsymbol{\Theta} + \mathbf{v} + \mathbf{e},$$

where  $\mathbf{p} = (p_1, \dots, p_D)'$  is the vector of direct survey estimates,  $\mathbf{X} = (\mathbf{x}'_1, \dots, \mathbf{x}'_D)'$  be  $D \times p$  matrix of auxiliary variates,  $\boldsymbol{\beta}$  is the fixed effect parameter vector,  $\mathbf{v} = (v_1, \dots, v_m)'$  is a vector of domain random effects such that  $\mathbf{v} \sim N(\mathbf{0}, \sigma_v^2 \mathbf{I}_D)$ ,  $\mathbf{I}_D$  is the unit matrix of dimension  $D$ ,  $\mathbf{e} = (e_1, \dots, e_D)'$  is the vector of sampling errors with  $\mathbf{e} \sim N(\mathbf{0}, \boldsymbol{\Omega})$ , where  $\boldsymbol{\Omega} = \text{diag}\{\sigma_{ei}^2; 1 \leq i \leq D\}$  is the matrix of design variances.  $\boldsymbol{\Psi} = \{\text{diag}(\mathbf{x}'_1), \dots, \text{diag}(\mathbf{x}'_D)\}'$  is a  $D \times pD$  matrix of known auxiliary data;  $\boldsymbol{\Theta} = (\boldsymbol{\gamma}'(l_1), \dots, \boldsymbol{\gamma}'(l_D))'$  is a spatial Gaussian random vector of dimension  $pD \times 1$  such that  $E(\boldsymbol{\Theta} | \boldsymbol{\Psi}, \mathbf{l}) = \mathbf{0}$  and covariance matrix  $\text{var}(\boldsymbol{\Theta} | \boldsymbol{\Psi}, \mathbf{l}) = \boldsymbol{\Sigma}_\eta = \mathbf{W} \otimes (\mathbf{c}\mathbf{c}')$ , where  $\otimes$  denotes the Kronecker product. The matrix  $\mathbf{W} = 1 / (1 + L(l_i, l_j))$  defines the spatial distances between sample spatial locations  $(l_i, l_j)$ , specifically distances between centroids of two locations  $(i, j)$ . In general, the only constraint on the vector  $\mathbf{c}$  is that  $\boldsymbol{\Sigma}_\eta = \mathbf{W} \otimes (\mathbf{c}\mathbf{c}')$  is symmetric and non-negative definite. Following Chandra *et al.* (2017), consider  $\mathbf{c} = \sqrt{\eta} \mathbf{1}_p$ , where  $\eta \geq 0$  and  $\mathbf{1}_p$  denotes the unit vector of order  $p$ . So,  $\boldsymbol{\Sigma}_\eta = \eta \mathbf{W} \otimes (\mathbf{1}_p \mathbf{1}_p')$  involves non zero covariance  $\text{cov}(\gamma_k(l_i), \gamma_h(l_j)) = \eta / (1 + L(l_i, l_j))$

between  $\gamma_k(l_i)$  and  $\gamma_h(l_j)$  for sample spatial locations  $(i, j)$ , with  $k \neq h = 1, \dots, p$  and diagonal elements as  $\eta$ . The parameter  $\eta$  denotes the strength of spatial heterogeneity being explained by nonstationary auxiliary variables. In particular  $\eta = 0$  indicates the situation where the model is spatially homogeneous. In HB framework, the sampling and linking models for HBNSP are then expressed as

$$\mathbf{p}|\mathbf{P} \sim N(\mathbf{P}, \mathbf{\Omega}) \text{ and } \text{logit}(\mathbf{P})|\boldsymbol{\beta}, \eta, \sigma_v^2 \sim N(\mathbf{X}\boldsymbol{\beta}, \boldsymbol{\Psi}\Sigma_\eta\boldsymbol{\Psi}' + \sigma_v^2\mathbf{I}_D).$$

The prior for hyper-parameter  $\boldsymbol{\beta}$  is  $N(0, \sigma_0^2)$  and for variance parameters  $\eta$  and  $\sigma_v^2$  prior is  $IG(a_0, b_0)$ , where  $\sigma_0^2$  is set to be very large (say,  $10^6$ ) and very small value for  $a_0$  and  $b_0$ . Note that HBNSP reduces to HBP when  $\eta = 0$ . Gibbs sampling method is implemented to estimate posterior mean  $E(P_i|\mathbf{p})$  and posterior variance  $\text{var}(P_i|\mathbf{p})$ . The required full conditional distribution of parameters under HBP and HBNSP models are given in Section 2.3.

### 2.3. Survey-weighted estimation

The HB modeling of respectively unweighted and survey-weighted small area proportions is a way to check the impact of complex survey design information in the resultant estimates. Survey-weighted direct estimates used for HB modeling purpose have the potentiality to reduce the bias or design error of the final estimates. Consider sample  $s$  of size  $n$  is drawn from population  $U$  using a complex design or at least unequal probability scheme. Let  $p_{ij}$  be the selection probability attached to  $j^{\text{th}}$  sampling unit  $y_{ij}$  in the area  $i$ . The basic design weight can be given by  $w_{ij} = (n_i p_{ij})^{-1}$ . These weights can be adjusted to account for non-response and/or auxiliary information (Hidiroglou and You, 2016). Normalized survey-weights  $d_{ij}$  may also be constructed,  $d_{ij} = w_{ij} \left( \sum_j w_{ij} \right)^{-1}$ . Liu *et al.* (2014) and Anjoy *et al.* (2019) have considered HB modeling of survey-weighted small area proportions, where GLMM structure was used for estimation of area proportions. But the effects of taking informative samples were not discussed. Here, two alternative models of HBNSP are defined to study the impact of design informativeness while aim is to estimate small area proportions in presence of spatial nonstationary auxiliary variables using the above furnished HBNSP technique. Let,  $p_{i.uw}$  be the direct survey unweighted estimator for small area proportion  $P_i$ ,

$$p_{i.uw} = (n_i)^{-1} \sum_{j=1}^{n_i} y_{ij} \text{ and the variance of } p_{i.uw} \text{ is given as } \sigma_{ei.uw}^2 = n_i^{-1} P_i (1 - P_i).$$

The survey-weighted estimator denoted as,  $p_{i.sw}$  and its variance is expressed as,

$$p_{i.sw} = \left( \sum_{j=1}^{n_i} w_{ij} \right)^{-1} \sum_{j=1}^{n_i} w_{ij} y_{ij} \text{ and } \sigma_{ei.sw}^2 = \left( \sum_{j=1}^{n_i} w_{ij} \right)^{-2} \left\{ \sum_{j=1}^{n_i} w_{ij} (w_{ij} - 1) (y_{ij} - P_i)^2 \right\}.$$

Two HBNSP methods are explored for the impact of complex survey design, denoted as HBNSP1 and HBNSP2. These models are furnished below:

**HBNSP1:** Does not incorporate survey-weight

Sampling model:  $\mathbf{p}_{uw}/\mathbf{P} \sim N(\mathbf{P}, \boldsymbol{\Omega}_{uw})$

Linking model:  $g(\mathbf{P})/\boldsymbol{\beta}, \eta, \sigma_v^2 \sim N(\mathbf{X}\boldsymbol{\beta}, \boldsymbol{\Psi}\boldsymbol{\Sigma}_\eta\boldsymbol{\Psi}' + \sigma_v^2\mathbf{I}_D)$

**HBNSP2:** Incorporate survey-weight

Sampling model:  $\mathbf{p}_{sw}/\mathbf{P} \sim N(\mathbf{P}, \boldsymbol{\Omega}_{sw})$

Linking model:  $g(\mathbf{P})/\boldsymbol{\beta}, \eta, \sigma_v^2 \sim N(\mathbf{X}\boldsymbol{\beta}, \boldsymbol{\Psi}\boldsymbol{\Sigma}_\eta\boldsymbol{\Psi}' + \sigma_v^2\mathbf{I}_D)$

The required full conditional distributions of HBNSP1 and HBNSP2 under Gibbs sampler are given as below. Within the Gibbs sampler, particularly Metropolis-Hastings (M-H) algorithm is used for drawing random samples from full conditional distributions of posterior quantities. For HBP model, the full conditional distributions for the Gibbs sampler are given as,

$$P_i/\boldsymbol{\beta}, \sigma_v^2, p_i \propto \frac{1}{P_i(1-P_i)\sqrt{\sigma_{ei}^2\sigma_v^2}} \exp\left(-\frac{(p_i - P_i)^2}{2\sigma_{ei}^2} - \frac{(\log \text{it}(P_i) - \mathbf{x}'_i\boldsymbol{\beta})^2}{2\sigma_v^2}\right),$$

$$\boldsymbol{\beta}/P_i, \sigma_v^2 \sim N\left((\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\log \text{it}(\mathbf{P}), \sigma_v^2(\mathbf{X}'\mathbf{X})^{-1}\right), \text{ and}$$

$$\sigma_v^2|\boldsymbol{\beta}, P_i \sim IG\left(a + \frac{D}{2}, b + \frac{\sum_{i=1}^D (\log \text{it}(P_i) - \mathbf{x}'_i\boldsymbol{\beta})^2}{2}\right).$$

For HBNSP model, the full conditional distributions for the Gibbs sampler are given as,

$$\mathbf{P}/\boldsymbol{\beta}, \eta, \sigma_v^2, \mathbf{p} \propto \left|(\boldsymbol{\Psi}\boldsymbol{\Sigma}_\eta\boldsymbol{\Psi}' + \sigma_v^2\mathbf{I}_D)\right|^{-\frac{1}{2}}|\boldsymbol{\Omega}|^{-\frac{1}{2}} \exp\left[-\frac{1}{2}\{(\mathbf{p} - \mathbf{P})'\boldsymbol{\Omega}^{-1}(\mathbf{p} - \mathbf{P}) + (\log \text{it}(\mathbf{P}) - \mathbf{X}\boldsymbol{\beta})'(\boldsymbol{\Psi}\boldsymbol{\Sigma}_\eta\boldsymbol{\Psi}' + \sigma_v^2\mathbf{I}_D)^{-1}(\log \text{it}(\mathbf{P}) - \mathbf{X}\boldsymbol{\beta})\}\right] \left|\frac{\partial \log \text{it}(\mathbf{P})}{\partial \mathbf{P}}\right|,$$

$$\boldsymbol{\beta}/\mathbf{P}, \eta, \sigma_v^2 \sim \text{MVN}\left[(\mathbf{X}'\boldsymbol{\Pi}^{-1}\mathbf{X})^{-1}(\mathbf{X}'\boldsymbol{\Pi}^{-1}\log \text{it}(\mathbf{P})), (\sigma_v^2\mathbf{I}_D + \boldsymbol{\Psi}\boldsymbol{\Sigma}_\eta\boldsymbol{\Psi}')(\mathbf{X}'\boldsymbol{\Pi}^{-1}\mathbf{X})^{-1}\right],$$

$$\eta|\boldsymbol{\beta}, \sigma_v^2, \mathbf{P} \sim IG\left[a_0 + \frac{D}{2}, b_0 + \frac{(\log \text{it}(\mathbf{P}) - \mathbf{X}\boldsymbol{\beta} - \mathbf{v})'(\log \text{it}(\mathbf{P}) - \mathbf{X}\boldsymbol{\beta} - \mathbf{v})}{2}\right], \text{ and}$$

$$\sigma_v^2|\boldsymbol{\beta}, \eta, \mathbf{P} \sim IG\left[a_1 + \frac{D}{2}, b_1 + \frac{(\log \text{it}(\mathbf{P}) - \mathbf{X}\boldsymbol{\beta} - \boldsymbol{\Psi}\boldsymbol{\Theta})'(\log \text{it}(\mathbf{P}) - \mathbf{X}\boldsymbol{\beta} - \boldsymbol{\Psi}\boldsymbol{\Theta})}{2}\right].$$

where,  $\boldsymbol{\Pi} = \boldsymbol{\Psi}\boldsymbol{\Sigma}_\eta\boldsymbol{\Psi}' + \sigma_v^2\mathbf{I}_D + \boldsymbol{\Omega}$ . Recall that  $\boldsymbol{\Sigma}_\eta = \eta\mathbf{W} \otimes (\mathbf{1}_p\mathbf{1}_p')$  with distance matrix  $\mathbf{W} = 1/(1 + L(l_i, l_j))$ .

For HBNSP1 model, the full conditional distributions for the Gibbs sampler are given as,

$$\begin{aligned} \mathbf{P}|\boldsymbol{\beta}, \eta, \sigma_v^2, \mathbf{p}_{uw} &\sim |(\boldsymbol{\Psi}\boldsymbol{\Sigma}_\eta\boldsymbol{\Psi}' + \sigma_v^2\mathbf{I}_D)|^{-\frac{1}{2}}|\boldsymbol{\Omega}_{uw}|^{-\frac{1}{2}} \exp\left[-\frac{1}{2}\{(\mathbf{p}_{uw} - \mathbf{P})' \boldsymbol{\Omega}_{uw}^{-1}(\mathbf{p}_{uw} - \mathbf{P})\right. \\ &\quad \left.+ (\log \text{it}(\mathbf{P}) - \mathbf{X}\boldsymbol{\beta})' (\boldsymbol{\Psi}\boldsymbol{\Sigma}_\eta\boldsymbol{\Psi}' + \sigma_v^2\mathbf{I}_D)^{-1} (\log \text{it}(\mathbf{P}) - \mathbf{X}\boldsymbol{\beta})\right] \left| \frac{\partial \log \text{it}(\mathbf{P})}{\partial \mathbf{P}} \right|, \\ \boldsymbol{\beta}|\mathbf{P}, \eta, \sigma_v^2 &\sim \text{MVN}\left[\left(\mathbf{X}'\boldsymbol{\Pi}_{uw}^{-1}\mathbf{X}\right)^{-1}\left(\mathbf{X}'\boldsymbol{\Pi}_{uw}^{-1}\log \text{it}(\mathbf{P})\right), \left(\sigma_v^2\mathbf{I}_D + \boldsymbol{\Psi}\boldsymbol{\Sigma}_\eta\boldsymbol{\Psi}'\right)\left(\mathbf{X}'\boldsymbol{\Pi}_{uw}^{-1}\mathbf{X}\right)^{-1}\right], \\ \eta|\boldsymbol{\beta}, \sigma_v^2, \mathbf{P} &\sim \text{IG}\left[a_0 + \frac{D}{2}, b_0 + \frac{(\log \text{it}(\mathbf{P}) - \mathbf{X}\boldsymbol{\beta} - \mathbf{v})' (\log \text{it}(\mathbf{P}) - \mathbf{X}\boldsymbol{\beta} - \mathbf{v})}{2}\right], \text{ and} \\ \sigma_v^2|\boldsymbol{\beta}, \eta, \mathbf{P} &\sim \text{IG}\left[a_1 + \frac{D}{2}, b_1 + \frac{(\log \text{it}(\mathbf{P}) - \mathbf{X}\boldsymbol{\beta} - \boldsymbol{\Psi}\boldsymbol{\Theta})' (\log \text{it}(\mathbf{P}) - \mathbf{X}\boldsymbol{\beta} - \boldsymbol{\Psi}\boldsymbol{\Theta})}{2}\right]. \end{aligned}$$

where,  $\mathbf{p}_{uw} = (p_{1.uw}, \dots, p_{D.uw})'$ ;  $\boldsymbol{\Omega}_{uw} = \text{diag}\{\sigma_{ei.uw}^2; 1 \leq i \leq D\}$  and  $\boldsymbol{\Pi}_{uw} = \boldsymbol{\Psi}\boldsymbol{\Sigma}_\eta\boldsymbol{\Psi}' + \sigma_v^2\mathbf{I}_D + \boldsymbol{\Omega}_{uw}$ .

For HBNSP2 model, the full conditional distributions for the Gibbs sampler are given as,

$$\begin{aligned} \mathbf{P}|\boldsymbol{\beta}, \eta, \sigma_v^2, \mathbf{p}_{sw} &\sim |(\boldsymbol{\Psi}\boldsymbol{\Sigma}_\eta\boldsymbol{\Psi}' + \sigma_v^2\mathbf{I}_D)|^{-\frac{1}{2}}|\boldsymbol{\Omega}_{sw}|^{-\frac{1}{2}} \exp\left[-\frac{1}{2}\{(\mathbf{p}_{sw} - \mathbf{P})' \boldsymbol{\Omega}_{sw}^{-1}(\mathbf{p}_{sw} - \mathbf{P})\right. \\ &\quad \left.+ (\log \text{it}(\mathbf{P}) - \mathbf{X}\boldsymbol{\beta})' (\boldsymbol{\Psi}\boldsymbol{\Sigma}_\eta\boldsymbol{\Psi}' + \sigma_v^2\mathbf{I}_D)^{-1} (\log \text{it}(\mathbf{P}) - \mathbf{X}\boldsymbol{\beta})\right] \left| \frac{\partial \log \text{it}(\mathbf{P})}{\partial \mathbf{P}} \right|, \\ \boldsymbol{\beta}|\mathbf{P}, \eta, \sigma_v^2 &\sim \text{MVN}\left[\left(\mathbf{X}'\boldsymbol{\Pi}_{sw}^{-1}\mathbf{X}\right)^{-1}\left(\mathbf{X}'\boldsymbol{\Pi}_{sw}^{-1}\log \text{it}(\mathbf{P})\right), \left(\sigma_v^2\mathbf{I}_D + \boldsymbol{\Psi}\boldsymbol{\Sigma}_\eta\boldsymbol{\Psi}'\right)\left(\mathbf{X}'\boldsymbol{\Pi}_{sw}^{-1}\mathbf{X}\right)^{-1}\right], \\ \eta|\boldsymbol{\beta}, \sigma_v^2, \mathbf{P} &\sim \text{IG}\left[a_0 + \frac{D}{2}, b_0 + \frac{(\log \text{it}(\mathbf{P}) - \mathbf{X}\boldsymbol{\beta} - \mathbf{v})' (\log \text{it}(\mathbf{P}) - \mathbf{X}\boldsymbol{\beta} - \mathbf{v})}{2}\right], \text{ and} \\ \sigma_v^2|\boldsymbol{\beta}, \eta, \mathbf{P} &\sim \text{IG}\left[a_1 + \frac{D}{2}, b_1 + \frac{(\log \text{it}(\mathbf{P}) - \mathbf{X}\boldsymbol{\beta} - \boldsymbol{\Psi}\boldsymbol{\Theta})' (\log \text{it}(\mathbf{P}) - \mathbf{X}\boldsymbol{\beta} - \boldsymbol{\Psi}\boldsymbol{\Theta})}{2}\right]. \end{aligned}$$

where,  $\mathbf{p}_{sw} = (p_{1.sw}, \dots, p_{D.sw})'$ ;  $\boldsymbol{\Omega}_{sw} = \text{diag}\{\sigma_{ei.sw}^2; 1 \leq i \leq D\}$  and  $\boldsymbol{\Pi}_{sw} = \boldsymbol{\Psi}\boldsymbol{\Sigma}_\eta\boldsymbol{\Psi}' + \sigma_v^2\mathbf{I}_D + \boldsymbol{\Omega}_{sw}$ .

### 3. Empirical Evaluations

This section reports the empirical results on the comparative performances of different estimators of the small area proportions which have been described previously. In particular, empirical performance of the proposed small area estimator HBNSP as compared to HBP is evaluated. Further, empirical performance of HBNSP1 and HBNSP2 also has been evaluated. Two types of simulation studies are used here. Section 3.1 describes the model-based simulation set up to evaluate the performance of HBNSP and HBP. In model-based simulation, population data is generated using a specified model. In section 3.2, a design-based simulation study is presented for comparing the performance of nonstationary process HBNSP1 and HBNSP2 which respectively, ignores and considers the modeling of survey-weighted proportions. Here, the aim is to explore impact of the incorporation of complex survey



information. Simulation studies have been implemented in R. Different performance indicators considered for comparison of small area estimators are as below. Let  $t$  is the subscript for  $T$  simulations.

- $RB_i = 100 \times \left( T^{-1} \sum_{t=1}^T P_i^{(t)} \right)^{-1} \left\{ T^{-1} \sum_{t=1}^T \left( \hat{P}_i^{(t)} - P_i^{(t)} \right) \right\}$  is the percentage relative bias (RB) for  $i^{\text{th}}$  small area, where  $\hat{P}_i^{(t)}$  is the estimate of true population mean  $P_i^{(t)}$  for  $i^{\text{th}}$  for small area at  $t^{\text{th}}$  simulation.
- $RRMSE_i = 100 \times \left( T^{-1} \sum_{t=1}^T P_i^{(t)} \right)^{-1} \left\{ \sqrt{T^{-1} \sum_{t=1}^T \left( \hat{P}_i^{(t)} - P_i^{(t)} \right)^2} \right\}$  is the percentage relative root mean squared error (RRMSE) for  $i^{\text{th}}$  for small area.
- $CR_i = 100 \times T^{-1} \sum_{t=1}^T I \left\{ LB \left( \hat{P}_i^{(t)} \right) \leq P_i^{(t)} \leq UB \left( \hat{P}_i^{(t)} \right) \right\}$  is the percentage coverage rate (CR) for  $i^{\text{th}}$  small area, where  $LB \left( \hat{P}_i^{(t)} \right)$  and  $UB \left( \hat{P}_i^{(t)} \right)$  are respectively Lower Bound ( $LB$ ) and Upper Bound ( $UB$ ) of the estimated population mean  $\hat{P}_i^{(t)}$ .  $I(\cdot)$  indicates an indicator function which takes values 1 if true parameter value  $P_i^{(t)}$  is within the computed interval, otherwise it takes value 0. This CR% particularly demonstrate the credible interval property of HB models.

For design-based simulation,  $P_i^{(t)}$  is equal to  $P_i$  or true population mean. A better model should show smaller values for all the performance indicators except CR. Higher the CR better is the model.

### 3.1. Model-based simulations

In model-based simulations the data were generated using both stationary and nonstationary processes. In stationary data generation process (SDGP), the regression coefficients are spatially invariant. The aim of this simulation set is to examine how HBNSP performs when the data follows spatial stationary process. Here, data is generated via the linking model:

$$\text{logit}(P_i) = 1 + x_i + v_i, \quad i = 1, \dots, D = 100.$$

In case of nonstationary data generation process (NSDGP), data is generated from the following model:

$$\text{logit}(P_i) = 1 + x_i + \sqrt{\eta} \left( \gamma_1(l_i) + \gamma_2(l_i) x_i \right) + v_i, \quad i = 1, \dots, D = 100$$

Here the values of  $x_i$  were independently drawn from the uniform distribution  $x_i \sim \text{Uniform}[0,1]$  and area random effects independently drawn as  $v_i \sim N(0, \sigma_v^2 = 0.0625)$ . Again, the sampling model part is considered as  $p_i = P_i + e_i; i = 1, \dots, D$ . The independent sampling errors  $e_i$  are generated from  $N(0, \sigma_{ei}^2)$  with  $\sigma_{ei}^2$  taking values 0.01, 0.02, 0.03 and 0.04 respectively for equal number of areas. To define *longitude* <sub>$i$</sub>  and *latitude* <sub>$i$</sub>  of spatial locations,

it is assumed that observations have been drawn from a two-dimensional grid consist of a  $(\sqrt{D} \times \sqrt{D})$  points uniformly spaced between -1 to 1 with a distance of  $2/(\sqrt{D}-1)$  between any two neighbouring points along the vertical and horizontal axes. The  $D$  points or spatial locations are arranged in such a way that  $k_1$  varies from -1 to 1 for each given  $k_2$ , which also then varies from -1 to 1. For example, when  $D=100$ , the set  $(k_1, k_2)$  is,  $\{k_1, k_2 = -1, -0.77, -0.55, -0.33, -0.11, 0.11, 0.33, 0.55, 0.77, 1\}$ . Further,  $(\gamma_1(l_i), \gamma_2(l_i))'$  has been defined as a random draw from  $N(0, \mathbf{W} \otimes \mathbf{I}_2)$  with  $\mathbf{W} = 1/(1+L(l_i, l_j))$  being the distance matrix between spatial locations  $(l_i, l_j)$ . The values of  $\eta$  have been used as 0.5, 1, 2, 4 in this study. This simulation set up is followed from Chandra *et al.* (2017).

The process of generating data and estimation of small area proportions by implementing HBP and HBNSP methods was independently replicated  $T = 500$  times from both stationary and nonstationary data generation process. The empirical performance and relative efficiency of the proposed HBNSP is compared with the HBP which excludes spatial nonstationarity structure. Performance of the small area HB estimators under each model is compared with respect to different prior cases for variance parameter  $\sigma_v^2$ . Specifically,  $IG(0.01, 0.01)$  and  $IG(0.1, 0.1)$  prior cases were taken up for such sensitivity analysis with respect to prior for variance parameter  $\sigma_v^2$ . However, the result from prior  $IG(0.1, 0.1)$  are only reported. As inferences based on different non-informative priors were found to be similar. The prior for hyperparameter  $\beta$  was taken as  $N(0, 10^6)$ . The prior for  $\eta$  was taken to be same as  $\sigma_v^2$ . To implement the Gibbs sampler, three independent chains are used each of length 10000. The first 5000 iterations are deleted as “burn-in” periods. Further, following Gelman and Rubin (1992), potential scale reduction factor ( $\hat{R}$ ) is used to monitor the convergence of the M-H within Gibbs sampler. The  $\hat{R}$  value close to 1 is expected and equal to 1 implies stationarity.

Table 1 shows the average values of relative biases (RB), relative root mean squared errors (RRMSE) and coverage rates (CR) for HBP and HBNSP methods investigated in model-based simulations. In Table 1 these values are presented as percentage and averages over the small areas of interest ( $D = 100$ ). Summary statistics of RB, RRMSE and CR for HBP and HBNSP methods for different values of  $\eta$  under NSDGP for  $D = 64$  and 100 areas are reported in Appendix (Table A1-A2). The differences between two small area predictors HBP and HBNSP in Table 1 are essentially as one would expect. When the underlying data is stationary (*i.e.*, data generated through SDGP), with identical value of RB, value of RRMSE of HBP is marginally smaller than the HBNSP. In contrast, in presence of nonstationarity in data (*i.e.*, data generated through NSDGP), the HBNSP method performs consistently better than the HBP method both in terms of RB and RRMSE for all values of nonstationarity parameter  $\eta$ . Additionally, HBNSP has shown better coverage properties. Noncoverage rate is marginally higher for HBP method.

**Table 1: The average values of percentage relative biases (%RB), percentage relative root mean squared errors (%RRMSE) and percentage coverage rates (%CR) for HBP and HBNSP methods in model-based simulation. Averaged  $D=100$  areas**

Criterion	SDGP		NSDGP							
			$\eta=0.5$		$\eta=1$		$\eta=2$		$\eta=4$	
	HBP	HBNSP	HBP	HBNSP	HBP	HBNSP	HBP	HBNSP	HBP	HBNSP
RB	0.065	0.065	-0.974	-0.478	-0.891	-0.402	-0.747	-0.359	-0.602	-0.403
RRMSE	4.289	4.447	5.636	4.635	5.130	4.242	4.593	4.047	4.622	4.281
CR	71	86	81	92	89	95	93	95	93	94

### 3.2. Design-based simulations

As in real life small area applications, one cannot be confident that our data ideally follow an assumed model, rather a working model is fitted. The endeavor of design-based simulation is to evaluate the performance of different SAE methods in the context of a realistic population, where a model assumption is essentially an approximation. For this simulation study, debt-investment survey (AIDIS-2013) data of National Statistical Office for rural areas of the state of Karnataka in India is used. The sample size of AIDIS-2013 is 2340 units (rural households including both indebted and non-indebted) spread over 30 districts of Karnataka. The AIDIS sample data is considered as fixed population of size 2340 units (or households) and 30 districts as small areas. Population size of small areas ranges between a minimum of 55 to a maximum of 112 with an average of 78 households. The variable of interest  $y_{ij}$  is binary which takes value 1 if a household is indebted and 0 otherwise. The aim is to estimate the proportions of indebted farm households (*i.e.* or incidence of indebtedness in farm households) at the district level. Here, Probability Proportional to Size with Replacement (PPSWR) samples were drawn independently within each small area instead of Simple Random Sampling to take into account the effect of varying sampling weights. Motivated from the simulation set up in Hidiroglou and You (2016), PPSWR sampling was employed as follows: a size measure  $z_{ij}$  is defined for a given unit  $y_{ij}$ . Using these  $z_{ij}$  values, selection probabilities  $p_{ij} = z_{ij} \left( \sum_j z_{ij} \right)^{-1}$  are computed and used to select PPSWR samples of equal size  $n_i$  from each small area. Then PPSWR samples of sizes  $n_i = 10, 15, 20$  and  $25$  were drawn from each small area based on selection probabilities  $p_{ij}$ . This selection probability, computed from a size measure  $z_{ij}$  is a linear combination of two auxiliary variables, namely Household size and Area operated (in hectare). The basic design weight calculated as,  $w_{ij} = (n_i p_{ij})^{-1}$ . Further, two cases were considered for fitting HBNSP models. Case 1- No auxiliary variable is included in the HB models and linking model contains only intercept and random effect (*i.e.*, random mean form of model). Case 2- Available auxiliary variable (Area operated, in hectare) is used as covariate in the HB models and linking model contains intercept, one auxiliary variable and random effect (*i.e.*, random intercept form of model). The prior for hyperparameter  $\beta$  was  $N(0, 10^6)$ . The prior for  $\eta$  and  $\sigma_v^2$  was taken to be  $IG(0.1, 0.1)$ . Gibbs sampling method is implemented with three independent chains each of length 10000; the first 5000 iterations are deleted as “burn-in” periods. To monitor the convergence success potential scale reduction factor  $\hat{R}$  is observed. The  $\hat{R}$  value close to 1 determines that the MCMC sampler converged very well.

Table 2 presents the average values of RB, RRMSE and CR for the small area predictors defined by HBNSP1 and HBNSP2 methods investigated in design-based simulations under case 1. The average values of RB, RRMSE and CR for HBNSP1 and HBNSP2 under case 2 are reported in Table 3. Figure 1 plots the average values of bias for HBNSP1 and HBNSP2


methods in design-based simulations under case-1 (left side) and case-2 (right side). From these results, it is evident that design bias of survey-weighted predictor HBNSP2 is smaller than HBNSP1. Further, the values of RB for survey-weighted predictor reduces with sample size, which shows the property of design consistency of small area predictor HBNSP2. The RRMSE values are also smaller for HBNSP2 and having the same decreasing trend with increment of small area sample sizes. Investigation on coverage properties of both the models shows that noncoverage rate is higher for HBNSP1 model as compared to the other. As number of areas increases, HBNSP2 shows the better coverage percentage.

**Table 2. The average values of percentage relative biases (%RB), percentage relative root mean squared errors (%RRMSE) and percentage coverage rates (%CR) for HBNSP1 and HBNSP2 methods in design-based simulation under case 1**

Criterion	Method	$n_i = 10$	$n_i = 15$	$n_i = 20$	$n_i = 25$
RB	HBNSP1	2.52	3.28	4.15	4.10
	HBNSP2	1.97	1.74	1.45	1.35
RRMSE	HBNSP1	24.23	23.02	23.80	24.08
	HBNSP2	23.37	15.57	13.35	12.73
CR	HBNSP1	89	87	83	76
	HBNSP2	91	96	97	97

**Table 3: The average values of percentage relative biases (%RB), percentage relative root mean squared errors (%RRMSE) and percentage coverage rates (%CR) for HBNSP1 and HBNSP2 methods in design-based simulation under case 2**

Criterion	Method	$n_i = 10$	$n_i = 15$	$n_i = 20$	$n_i = 25$
RB	HBNSP1	3.45	3.77	4.90	4.88
	HBNSP2	2.41	1.67	1.02	1.07
RRMSE	HBNSP1	28.03	24.64	25.52	25.34
	HBNSP2	24.42	17.00	15.41	13.25
CR	HBNSP1	85	84	80	77
	HBNSP2	91	95	96	97



**Figure 1: Comparison of bias of HBNSP1 and HBNSP2 (HBNSP1: Solid line, HBNSP2: Dash line) under case 1 (left side) and case 2 (right side)**

#### 4. Concluding Remarks

The article describes a spatial nonstationary extension of the area level version of the hierarchical Bayes generalized linear mixed model and considers SAE of proportions under this model. The corresponding predictor is referred to as the spatial nonstationary hierarchical Bayes predictor (HBNSP) for small area proportions. This predictor can account for the

presence of spatial nonstationarity in the data where the parameters associated with the model covariates vary spatially.

Empirical results based on simulation studies provide evidence that the proposed HBNSP predictor is more efficient than the alternative hierarchical Bayes predictor under the area level generalized linear mixed model when there is a spatial nonstationarity in the data. The MSE estimation of the HBNSP predictor derived from associated posterior variance also performed reasonably well, with good coverage performance for nominal confidence intervals based on it. It is worth noting that in this article empirical studies were also carried out using survey weights to incorporate the sampling design in SAE. This seems more realistic to implement survey weighted estimation than assuming that the sampling design is customary non-informative.

The Census in India, like in other countries, usually has limited scope in collection of data. It focuses mainly on basic social and demographic information and that too at decennial interval. On the other hand, NSSO conducts regular surveys on several socio-economic indicators, but outcome is restricted to generate national and state level estimates, not administrative units below state because of small sample sizes for such units. Due to emphasis on disaggregate level Sustainable Development Goal indicators, Government of India as well as different State Governments are now struggling with generation of disaggregated level statistics. The SAE is only indispensable alternative to meet the growing demand for such disaggregated level statistics needed for decentralized policy planning. The SAE methodology discussed in this article can be used for calculating disaggregate level estimates of prevalence and proportions which is common in most of the socio-economic and health surveys.

## References

- Anjoy P., Chandra H. and Basak P. (2019). Estimation of disaggregate-level poverty incidence in Odisha under area-level Hierarchical Bayes small area model. *Social Indicators Research*, **144**, 251-273.
- Baldermann C., Salvati N. and Schmid, T. (2018). Robust small area estimation under spatial non-stationarity. *International Statistical Review*.  
DOI: <https://doi.org/10.1111/insr.12245>.
- Battese G. E., Harter R. M. and Fuller W. A. (1988). An error-component model for prediction of country crop areas using survey and satellite data. *Journal of the American Statistical Association*, **83**, 28-36.
- Brunsdon C., Fotheringham A. S. and Charlton M. E. (1996). Geographically weighted regression: A method for exploring spatial nonstationarity, *Geographical Analysis*, **28**, 281–298.
- Chandra H., Chambers R. and Salvati N. (2019). Small area estimation of survey weighted counts under aggregated level spatial model. *Survey Methodology*, **45**, 31-59.
- Chandra H., Kumar S. and Aditya K. (2018). Small area estimation of proportions with different levels of auxiliary data. *Biometrical Journal*, **60**, 395–415.
- Chandra H. and Salvati N. (2018). Small area estimation for count data under a spatial dependent aggregated level random effects model. *Communications in Statistics - Theory and Methods*, **47(5)**, 1234 -1255.
- Chandra H., Salvati N. and Chambers, R. (2017). Small area prediction of counts under a non-stationary spatial model. *Spatial Statistics*, **20**, 30-56.
- Cressie N. (1993). *Statistics for Spatial Data*. Wiley, New York.

- Fay R. E. and Herriot R. A. (1979). Estimates of income for small places: An application of James-Stein procedures to census data. *Journal of the American Statistical Association*, **74**, 269-277.
- Gelman A. and Rubin D. (1992). Inference from Iterative Simulation using Multiple Sequences. *Statistical Science*, **7**, 457-511.
- Hidiroglou M. A. and You Y. (2016). Comparison of unit level and area level small area estimators. *Survey Methodology*, **42**, 41-61.
- Jiang J. and Lahiri P. (2001). Empirical best prediction for small area inference with binary data. *Annals of the Institute of Statistical Mathematics*, **53**, 217-243.
- Liu B., Lahiri P. and Kalton G. (2014). Hierarchical Bayes modeling of survey-weighted small area proportions. *Survey Methodology*, **40**, 1-13.
- Rao J. N. K. and Molina I. (2015). *Small Area Estimation*. 2nd Edition. New York: John Wiley & Sons.
- You Y. and Zhou M. Q. (2011). Hierarchical Bayes small area estimation under a spatial model with application to health survey data. *Survey Methodology*, **37**, 25-37.

## APPENDIX

**Table A1: Summary statistics of percentage relative biases (%RB), percentage relative root mean squared errors (%RRMSE) and percentage coverage rates (%CR) for HBP and HBNSP methods in model-based simulations for different values of  $\eta$  under spatial nonstationary data generation process for  $D= 100$  small areas.**

Criterion	RB		RRMSE		CR	
	HBP	HBNSP	HBP	HBNSP	HBP	HBNSP
$\eta=0.5$						
Minimum	-9.50	-8.26	2.11	1.88	22	39
Q1	-3.70	-2.46	4.28	3.41	76	91
Mean	-0.97	-0.48	5.64	4.63	81	92
Median	-0.83	-0.50	5.39	4.38	86	95
Q3	0.76	1.09	6.46	5.47	93	98
Maximum	22.30	17.34	23.72	18.52	100	100
$\eta=1$						
Minimum	-9.30	-8.01	1.60	1.477	33	41
Q1	-3.48	-2.17	3.53	2.86	85	95
Mean	-0.89	-0.40	5.13	4.24	89	95
Median	-0.86	-0.32	4.72	4.03	94	98
Q3	0.79	1.00	6.06	5.16	98	99
Maximum	23.08	17.11	25.24	18.30	100	100
$\eta=2$						
Minimum	-8.73	-7.63	1.36	1.22	39	44
Q1	-3.25	-2.21	2.93	2.62	91	95
Mean	-0.75	-0.36	4.59	4.05	93	95
Median	-0.58	-0.06	4.25	3.73	98	99
Q3	0.79	0.89	5.63	5.05	99	100
Maximum	23.17	19.33	25.13	20.66	100	100
$\eta=4$						
Minimum	-8.17	-7.57	1.28	1.15	43	49

Q1	-3.14	-2.56	2.90	2.69	91	94
Mean	-0.60	-0.40	4.62	4.28	93	94
Median	-0.35	-0.24	4.27	3.75	98	99
Q3	0.78	0.83	5.72	5.28	99	99
Maximum	37.61	34.28	39.94	36.31	100	100

**Table A2: Summary statistics of percentage relative biases (%RB), percentage relative root mean squared errors (%RRMSE) and percentage coverage rates (%CR) for HBP and HBNSP methods in model-based simulations for different values of  $\eta$  under spatial nonstationary data generation process for  $D=64$  small areas**

Criterion	RB		RRMSE		CR	
	HBP	HBNSP	HBP	HBNSP	HBP	HBNSP
$\eta=0.5$						
Minimum	-8.21	-6.54	1.61	1.62	64	79
Q1	-2.16	-1.52	3.19	3.12	97	98
Mean	-0.34	0.00	4.18	4.10	97	98
Median	0.23	0.08	4.32	4.05	99	99
Q3	1.34	1.86	4.86	4.71	100	100
Maximum	5.76	5.66	8.63	7.13	100	100
$\eta=1$						
Minimum	-7.79	-6.35	1.39	1.49	68	81
Q1	-1.99	-1.45	2.98	2.37	98	99
Mean	-0.26	0.08	3.91	3.83	98	98
Median	0.35	0.29	4.05	3.83	100	99
Q3	1.33	1.86	4.61	4.61	100	100
Maximum	5.30	5.19	8.19	7.29	100	100
$\eta=2$						
Minimum	-7.35	-6.25	1.42	1.45	67	77
Q1	-1.76	-1.29	2.87	2.09	99	99
Mean	-0.20	0.12	3.86	3.83	98	98
Median	0.37	0.28	4.08	3.69	100	100
Q3	1.26	1.73	4.72	4.54	100	100
Maximum	4.92	4.54	7.77	7.28	100	100
$\eta=4$						
Minimum	-7.02	-5.88	1.58	1.55	68	72
Q1	-1.58	-1.09	3.26	3.42	97	98
Mean	0.30	0.28	4.40	4.36	98	98
Median	0.40	0.36	4.32	4.11	99	99
Q3	1.39	1.44	5.24	4.94	100	100
Maximum	6.25	5.80	9.48	9.45	100	100