

~~Revised~~

DRAFT

THE NATIONAL SAMPLE SURVEY
TWENTYEIGHTH ROUND
OCTOBER 1973 - JUNE 1974

NUMBER 292

TABLES WITH NOTES ON
MORBIDITY

ISSUED BY
NATIONAL SAMPLE SURVEY ORGANISATION
DEPARTMENT OF STATISTICS : MINISTRY OF PLANNING
GOVERNMENT OF INDIA
SEPTEMBER 1980

Note of Caution

Being the scanned copy of old NSS report, this document may suffer from following limitations -

- i. Poor Quality of the Scanned images.
- ii. Page(s) missing in between.
- iii. Improper sequencing/arrangement.

TABLES WITH NOTES ON
MORBIDITY

<u>CONTENTS</u>		<u>Page</u>
CHAPTER ONE	: Introduction	1
CHAPTER TWO	: Concepts and definition	2
CHAPTER THREE	: Summary findings	3
APPENDIX I	: Sampling Design and Estimation Procedure	i
Appendix II	: Statistical Tables :	iii
	List of tables	
	Tables	1

NATIONAL SAMPLE SURVEY
28TH ROUND : OCTOBER 1973 - JUNE 1974
TABLES WITH NOTES ON MORBIDITY
CHAPTER ONE

INTRODUCTION

1.1. In the twenty eight round of the National Sample Survey (NSS) information ~~particulars~~ on morbidity was collected along with the enquiry on births, deaths, migration and disability^{a/}. Information on morbidity was collected with reference to last 2 weeks preceding the date of survey from all members present in the sample households on the date of enquiry as well as visitors in the households. Morbidity was defined as sickness/injury/poisoning.

Information on sickness/injury/poisoning affecting members of the household including visitors during the 2 weeks preceding the date of enquiry together with ancillary information like age and sex of the person, nature of the ailment etc. was collected. Also along with information on morbidity data on chronic ailments prevailing among the members on the date of survey were collected.

1.2. The survey covered the whole of India excepting Ladakh district of Jammu & Kashmir, Mizoram, Arunachal Pradesh, Andaman & Nicobar Islands, Lakshadweep, Minicoy and Amindivi Islands, Dadra and Nagar Haveli, rural areas of Chandigarh and Nagaland, some rural areas in Tripura and in the districts of Sarguja and Bastar of Madhya Pradesh and ~~Chandrapur~~ Chandrapur and Amravati of Maharashtra.

1.3. The survey started in October 1973 and continued upto June 1974 and the survey period of nine months was split up into three sub-rounds of three months duration each. Data were collected from the sample households by the interview method. Broadly, the sample design was a stratified two stage one with village/urban blocks being first stage units and households as the second stage units and self-weighting at the state level separately for the rural and the urban sectors. In all, from about 1.29 lakhs of rural and 0.67 lakh urban households the relevant data on morbidity were collected. The present tables on morbidity are based on such data. The sample design and procedure of estimation are described in Appendix I.

a/ : Information on morbidity was also collected in seventeenth round of NSS from about 8,600 rural households and 4,400 urban households distributed over 4,310 villages & 2,237 urban blocks.

CHAPTER TWO
CONCEPTS AND DEFINITIONS

2.1. Household normally resident members and visitors : Persons who usually lived together and took food from the same kitchen were considered as usual or normally resident members of the household. Other members not usually residing in the household were classified as visitors. If a person lived in a place and took food at some other place, he was considered usual member at the place where he lived. Inmates (including the staff) of a mess, hotel, orphanage, etc. were treated as single member households. However, a family living in any one of the establishments was treated as a separate household by itself.

2.2. Morbidity was defined as sickness which included injury and poisoning also. Sickness meant any deviation from the state of physical and mental well being with a specific cause. Chronic ailments ~~and the associated~~ ~~xxxxxxx~~ were not, however, treated as morbidity.

2.3. The following categories of spells of morbidity could occur during a reference period.

- (a) Spells starting before the first day of the reference period and terminating within the reference period.
- (b) Spells starting and terminating within the reference period.
- (c) Spells starting within the reference period and continuing on the date of survey.
- (d) Spells starting before the first day of the reference period and continuing on the date of survey.

2.4. Incidence rate is defined by

$$\frac{\text{estimated number of spells (b + c)}}{\text{estimated population exposed to risk}} \times 1000$$

$$= 1000 \times \frac{\text{estimated number of spells starting within the reference period}}{\text{estimated population exposed to risk}}$$

The incidence rate measures the frequency of morbidity commencing during the reference period

2.5. Prevalence rate is defined by

$$\frac{\text{estimated number of spells (a+b+c+d)}}{\text{estimated population exposed to risk}} \times 1000$$

$$= 1000 \times \frac{\text{estimated number of spells prevailing in the reference period}}{\text{estimated population exposed to risk}}$$

The prevalence rate measures the frequency of morbidity prevailing in the reference period.

CHAPTER THREE
SUMMARY FINDINGS

3.1. Table 1 below presents incidence and prevalence rates of morbidity for all-India separately by states. From the table it is seen that at all-India level the incidence rate of morbidity per one thousand persons was 12.57 and 13.53 respectively for the rural and the urban sectors^{b/}.

Table 1 : Incidence and prevalence rate of morbidity by states and all-India, rural & urban areas.

states	incidence rate		prevalence rate	
	rural	urban	rural	urban
(1)	(2)	(3)	(4)	(5)
1. Andhra Pradesh	20.89	18.20	31.94	27.84
2. Assam	8.46	5.94	17.40	11.66
3. Bihar	4.69	6.05	10.16	11.42
4. Gujarat	4.48	2.85	10.17	6.26
5. Haryana	9.23	8.46	15.96	16.47
6. Jammu & Kashmir	11.42	9.22	17.79	12.80
7. Karnataka	8.44	7.24	14.92	13.35
8. Kerala	37.29	33.77	71.21	61.84
9. Madhya Pradesh	10.34	11.34	20.89	21.72
10. Maharashtra	17.78	20.06	27.57	32.18
11. Orissa	14.53	17.19	24.20	24.38
12. Punjab	19.25	16.68	27.47	24.62
13. Rajasthan	10.20	7.46	17.33	12.55
14. Tamil Nadu	20.65	21.01	33.99	31.98
15. Uttar Pradesh	7.95	6.27	13.22	10.54
16. West Bengal	11.35	17.30	27.71	33.85
17. All-India ^{b/}	12.57	13.53	22.46	22.77

[1/ All states and union territories under NSS coverage]

It also appears from the table that there were wide variation in the incidence rate between the states. Kerala with an incidence rate of 37.29 and 33.77 respectively in the rural and the urban sectors tops the list. This was followed by Tamil Nadu with a rate of 20.65 and 21.01 in the rural and urban sectors. The states Andhra, Maharashtra, Bihar, Orissa, Punjab and Tamil Nadu had incidence rates both in the rural and the urban sectors higher than the all-India rate. With regard to prevalence rate it is seen from the table that at all-India the prevalence rate of temporary ailments was 22.46 and 22.77 respectively in the rural and the urban sectors. Kerala here again tops the list with a rate of 71.21 in the rural and 61.84 in the urban sectors. This was followed by Tamil Nadu with a rate of 33.99 and 31.98. The states Andhra, Maharashtra, Orissa and Punjab had rates higher than the all-India rates.

^{b/} The incidence and prevalence rates of morbidity of NSS 28th round as estimated are somewhat lower than the rates as observed in the 17th round of the NSS presented in NSS No.129 : Report of Pilot Enquiry on Morbidity. A seminar meeting was held in July 1977 under the auspices of NSSO to examine vital rates and migration and morbidity rates of NSS 28th round and the consensus was that the morbidity rates could be released even though the estimates of NSS 28th round appeared somewhat lower than those of the Pilot Enquiry on Morbidity.

3.2. Table 2 presents incidence and prevalence rates of morbidity by type of ailment for all-India. From the table it is seen that excepting the group 'others' and 'not recorded' the incidence rate of influenza was highest both in the rural & urban areas. It was 2.2 per 1000 persons in both the sectors. In the rural sector influenza was followed by malaria with a rate of 1.1. The incidence rate of malaria in the urban sector was not so high. The prevalence rate of influenza also exceeded the corresponding rates of other ailments in both the sectors. Malaria and dysentery also had a comparatively high rate. The prevalence rates estimated for influenza, malaria and dysentery were 3.1, 1.8 and 1.3 in rural areas and the corresponding rates for the urban sector being 3.2, 1.2 and 1.3.

Table 2 : Incidence and prevalence rate of morbidity by type of temporary ailments for all-India, rural and urban areas

type of ailments (1)	incidence rate		prevalence rate	
	rural (2)	urban (3)	rural (4)	urban (5)
1. cholera	0.03	0.03	0.05	0.04
2. typhoid	0.12	0.19	0.35	0.44
3. dysentery	0.74	0.79	1.34	1.29
4. diarrhoea	0.27	0.22	0.46	0.33
5. diphtheria	0.02	0.01	0.03	0.01
6. whooping cough	0.33	0.25	0.84	0.60
7. tetanus	0.01	0.01	0.02	0.03
8. acute poliomyelities	0.01	0.02	0.03	0.05
9. small pox	0.38	0.44	0.63	0.79
10. measles	0.17	0.14	0.31	0.30
11. mumps	0.03	0.08	0.05	0.12
12. malaria	1.13	0.71	1.77	1.15
13. influenza	2.16	2.15	3.10	3.24
14. pneumonia	0.13	0.05	0.20	0.11
15. food poisoning	0.02	0.04	0.03	0.04
16. accident	0.39	0.54	0.80	1.01
17. others	4.34	5.16	8.27	8.51
18. not recorded	2.29	2.70	4.18	4.71
19. all ailments	12.57	13.53	22.46	22.77

3.3. Incidence and prevalence rates of morbidity broken down by age is shown in table 3. Both incidence and prevalence rates of sickness as revealed in table 3 was highest in the first year of age and gradually diminished as age increased. It was minimum in the age group 15-24 and then again rose higher with the age. The pattern was observed both in the rural and urban areas of all-India. The incidence rates per 1000 person at age '0' were estimated as 21.09 for rural and 30.29 for urban sectors. The corresponding prevalence rates for rural and urban sectors were 35.72 and 49.34 respectively.

Table 3 : Incidence and prevalence rate of temporary ailments by age for all-India, rural and urban

age in years	incidence rate		prevalence rate	
	rural	urban	rural	urban
(1)	(2)	(3)	(4)	(5)
0	21.09	30.29	35.72	49.34
1-4	20.65	24.36	35.50	39.04
5-14	10.04	11.02	17.15	18.14
15-24	9.53	9.30	16.38	14.97
25-44	12.04	13.64	22.10	23.13
45-59	13.64	13.41	25.51	25.24
60 & above	13.67	12.97	28.78	25.57
not recorded	103.98	-	180.54	-
all ages	12.57	13.53	22.46	22.77

3.4. Average duration of selected temporary ailments which terminated in recovery during the reference period is presented in table 3 by type of ailment. It is seen from the table that at all-India level average duration of all temporary ailments taken together was about 8.2 and 6.7 days in the rural and the urban sectors respectively. There was, however, some variation by type of ailments. In the rural sector, the duration of typhoid was 18.8 days while the duration of measles, mumps, small pox & malaria was about 8-10 days. In the urban sector similarly the duration of typhoid was about 16 days while the duration of small pox, mumps, measles & malaria was about 8-10 days.

Table 4 : Average duration (days) of selected temporary ailments terminating in recovery by type of ailments for all-India, rural and urban areas

type of ailments	duration of temporary ailments	
	rural	urban
(1)	(2)	(3)
1. cholera	5.70	5.32
2. typhoid	18.84	16.43
3. diphtheria	5.15	7.00
4. whooping cough	10.52	6.92
5. small pox	10.33	10.13
6. measles	9.61	8.31
7. mumps	9.02	8.27
8. malaria	8.36	6.54
9. influenza	6.66	5.76
10. pneumonia	7.94	7.46
11. food poisoning	12.95	11.14
12. accident	10.73	7.20
13. all ailments	8.16	6.68

3.5. Information on chronic ailments prevailing among the members of the household are presented in table 5. It is seen that the incidence rate of all chronic ailments taken together was 2098 and 1962 per 1,00,000 persons in the rural and urban sectors. It is also seen that asthma was the most prevalent chronic ailment, both in the rural and urban areas. High blood pressure with a rate of 44 in the rural sector compared to 132 in the urban sector appears to be more urban specific. The number of persons suffering from this ailment came to 376 in the rural areas and 355 in the urban areas per 1,00,000 persons. The next most important chronic diseases were rheumatism and tuberculosis in both the rural and urban areas, the number of cases being respectively 251 and 117 in rural whereas 146 and 137

Table 5 : Number of persons suffering from chronic diseases per 1,00,000 persons by type of chronic disease for all-India, rural & urban areas

type of chronic disease (1)	number of persons	
	rural (2)	urban (3)
1. tuberculosis	117	137
2. leprosy	40	25
3. syphilis	8	6
4. cancer	12	14
5. thyroid trouble or goitre	22	16
6. diabetes	39	80
7. mental illness	19	18
8. epilepsy	28	17
9. rheumatic fever	45	26
10. high blood pressure	44	132
11. bronchitis	41	43
12. asthma	376	355
13. peptic ulcer	89	66
14. kidney stone or kidney trouble	37	40
15. arthritis	22	17
16. rheumatism	251	146
17. stroke	13	13
18. piles	65	61
19. others	194	128
20. not recorded	636	629
21. all types	2098	1962

in urban areas. High blood pressure with a rate of 132 in the urban sector compared to 44 in the rural sector appears to be more urban specific.

APPENDIX - I

SAMPLING DESIGN AND ESTIMATION PROCEDURE

A stratified two stage design was adopted for the survey. The first stage units were villages in the rural sector and blocks in the urban sector and the second stage units were households from which the information was collected.

The whole of India was divided into 524 basic strata so that the basic strata did not cut across district boundaries. A district with more than 1.5 million 1971 census rural population formed 2 or more basic strata (by grouping contiguous tehsils) depending on their rural population. In Gujarat, however districts with population less than 1.5 million were divided into two strata in consideration of dry areas and concentration of tribal population at the request of SSB of that state.

Within each State and Union Territory the allotted number of villages and blocks were distributed over the constituent basic strata in proportion to their rural and urban population respectively.

The 1971 census list of villages was used as the sampling frame for selection of villages. Sample villages within each basic stratum were selected with probability proportional to population and with replacement in the form of two independent sub-samples.

All cities and towns within each basic stratum were grouped into a number of urban sub-strata. All towns with 1971 census population less than 20,000 formed sub-stratum 1, towns with 1971 population 20,000 to 49,999 formed sub-stratum 2 and those with 1971 population 50,000 to 99,999 formed sub-stratum 3. Each city with 1971 population 1,00,000 or more constituted a separate urban sub-stratum. Within each urban stratum, the allocated number of blocks was further allocated to the constituent sub-strata circular systematically with probability proportional to urban populations of the respective sub-strata. Within each urban.

sub-stratum, the allocated number of blocks were selected with probability proportional to size and with replacement.

Villages and blocks with large population content were divided into two or more compact areal sub-divisions of nearly equal population content. One of these sub-divisions (called hamlet in a village and sub-block in a block) was selected at random and the survey was confined to the selected sub-division only.

All households in a sample village/hamlet group were arranged into 6 classes on the basis of their means of livelihood viz., (i) landless labour, (ii) cultivator and (iii) others and whether (i) visited health centre or (ii) not. All households in a sample block/sub-block were also arranged into 6 classes by occupation divisions (0-1 and 2,3,4,5,6 and 7-8-9 and X). From the arranged frame in a village/block, a sample of households was selected linear systematically using the intervals and random start specified. Sampling intervals were so fixed as to make the design selfweighting at the state level separately for rural and urban sectors.

Unbiased method of estimation was adopted for estimating aggregate values of the characteristics, say, X, Y, Z, etc. under enquiry. Estimates of ratios were derived by dividing unbiased estimates of the numerators by the unbiased estimates of the denominators.

SNC.

Appendix - II
Statistical Tables

A list of tables included in this
is given below :

	<u>page no.</u>
Table (1R) : Incidence rate of temporary ailments by type of ailment separately by sex for different states and all-India : rural households, NSS 28th round.	1
Table (1v) : Incidence rate of temporary ailments by type of ailment separately by sex for different states and all-India : Urban households, NSS 28th round.	4
Table (2R) : Incidence rate of temporary ailments by age separately by sex for different states and all-India : rural households, NSS 28th round.	7
Table (2U) : Incidence rate of temporary ailments by age separately by sex for different states and all-India : urban households, NSS 28th round.	9
Table (3R) : Prevalence rate of temporary ailments by type of ailments separately by sex for different states and all-India : rural households, NSS 28th round	11
Table (3U) : Prevalence rate of temporary ailments by type of ailments separately by sex for different states and all-India : Urban households, NSS 28th round.	14
Table (4R) : Prevalence rate of temporary ailments by age separately by sex for different states and all-India : rural households, NSS round	17
Table (4U) : Prevalence rate of temporary ailments by age separately by sex for different states and all-India : urban households, NSS 28th round.	19
Table (5R) : Average duration (days) of temporary ailments terminating in recovery by type of ailment separately by sex for different states and all-India : rural households, NSS 28th round.	21

	<u>page no.</u>
Table (5U) : Average duration (days) of temporary ailments terminating in recovery by type of ailment separately by sex for different states and all-India : Urban households, NSS 28th round.	24
Table (6R) : Average duration (days) of temporary ailments terminating in recovery by age separately by sex for different states and all-India : rural households, NSS 28th round.	27
Table (6U) : Average duration (days) of temporary ailments terminating in recovery by age separately by sex for different states and all-India : urban households, NSS 28th round.	29
Table (7R) : Number of persons suffering from chronic diseases per 100,000 persons by type of chronic disease separately by sex for different states and all-India : rural households, NSS 28th round.	31
Table (7U) : Number of persons suffering from chronic disease per 100,000 persons by type of chronic disease separately by sex for different states and all-India : urban households, NSS 28th round.	34
Table (8R) : Number of persons suffering from chronic diseases per 100,000 persons by age separately by sex for different states and all-India : rural households, NSS 28th round.	37
Table (8U) : Number of persons suffering from chronic diseases per 100,000 persons by age separately by sex for different states and all-India : urban households, NSS 28th round.	39

Table (1R) : Incidence rate of temporary ailments by type of ailments separately by sex for different states and all-India, rural households ; NSS 28th round

states	sex	t y p e o f a i l m e n t s																			
		cholera	typhoid	dysentery (all forms)	diarrhoea	diphtheria	whooping cough	tetanus	acute poliomy- elitis	small pox	measles	mumps	malaria	influenza	pneumonia	food poisoning	accident	others	not re corded	all types of ailments	number of sample ailments
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)
Andhra Pradesh	M	-	0.13	0.58	0.14	-	0.31	0.05	-	0.70	0.13	-	1.95	4.68	0.18	-	0.35	7.28	5.44	21.92	492
	F	-	0.09	0.40	0.05	-	0.23	-	-	0.63	0.18	0.05	2.35	3.95	0.09	0.05	0.09	5.38	6.30	19.84	436
	T	-	0.11	0.49	0.09	-	0.27	0.02	-	0.66	0.16	0.02	2.14	4.32	0.14	0.02	0.22	6.36	5.87	20.89	928
Assam	M	-	0.30	2.34	0.14	0.36	0.29	-	-	0.29	-	-	0.96	3.04	0.07	-	0.14	0.43	1.07	9.43	130
	F	-	0.08	0.97	0.16	0.16	0.16	-	-	0.16	-	-	0.42	3.71	0.08	-	0.40	0.48	0.56	7.34	91
	T	-	0.19	1.69	0.15	0.27	0.23	-	-	0.23	-	-	0.70	3.35	0.08	-	0.27	0.46	0.84	8.46	221
Bihar	M	0.04	0.04	0.43	0.21	-	0.18	-	-	0.39	0.04	-	0.14	0.96	0.14	-	0.18	2.62	0.11	5.48	154
	F	0.04	0.04	0.15	0.11	-	0.11	0.04	-	0.18	-	-	0.07	0.77	0.04	-	0.04	2.20	0.11	3.90	106
	T	0.04	0.04	0.29	0.16	-	0.14	0.02	-	0.29	0.02	-	0.11	0.86	0.09	-	0.11	2.41	0.11	4.69	260
Gujarat	M	0.08	-	0.16	0.16	-	0.65	-	-	0.16	-	0.08	0.89	0.72	-	-	0.40	0.32	0.56	4.18	52
	F	0.08	0.17	-	0.09	-	0.08	-	-	0.26	-	-	0.94	1.37	0.08	-	0.42	0.61	0.68	4.79	56
	T	0.08	0.08	0.08	0.13	-	0.37	-	-	0.21	-	0.04	0.92	1.04	0.04	-	0.42	0.45	0.62	4.48	108
Haryana	M	0.12	0.12	0.18	0.18	-	0.43	-	-	0.18	0.37	-	1.29	0.61	0.37	0.06	0.31	1.41	4.42	10.05	164
	F	0.14	0.34	0.34	0.07	0.07	0.28	-	-	0.21	0.28	-	0.76	0.28	0.07	-	-	0.76	4.75	8.35	121
	T	0.13	0.23	0.26	0.13	0.03	0.36	-	-	0.19	0.32	-	1.04	0.45	0.23	0.03	0.16	1.10	4.57	9.23	285
Jammu & Kashmir	M	-	0.35	1.29	0.70	0.03	0.38	-	-	0.03	0.07	-	0.53	4.29	0.21	0.03	0.14	2.11	0.73	10.89	318
	F	0.04	0.35	1.48	0.89	-	0.54	-	-	0.04	0.16	0.04	0.40	5.22	0.20	0.04	0.04	1.76	0.78	11.98	308
	T	0.02	0.35	1.38	0.79	0.02	0.46	-	-	0.04	0.11	0.02	0.47	4.73	0.20	0.04	0.09	1.94	0.76	11.42	621

+ M = male, F = female, T = total

Table (1R)Contd. : Incident rate of temporary ailments by type of ailments separately by sex for different states and all-India, rural households : NSS 28th round

states	sex +	type of ailments																			
		cholera	typhoid	dysentery (all forms)	diarrhoea	diphtheria	whooping cough	tetanus	acute poliomyelitis	small pox	measles	mumps	malaria	influenza	pneumonia	food poisoning	accident	others	not recorded	all type of ailments	number of sample ailments
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)
Karnataka	M	-	-	1.06	-	-	0.64	-	0.07	0.14	-	-	0.14	0.71	-	0.07	0.07	2.06	3.69	8.65	122
	F	-	-	0.51	-	-	0.51	-	-	0.29	0.07	0.07	-	0.65	-	-	0.22	2.03	3.85	8.20	113
	T	-	-	0.79	-	-	0.57	-	0.04	0.22	0.04	0.04	0.07	0.68	-	0.04	0.14	2.04	3.77	8.44	235
Kerala	M	-	-	2.58	0.84	0.07	0.35	-	-	0.28	0.63	-	-	5.24	-	-	1.82	15.92	11.52	39.25	562
	F	-	0.07	2.26	1.40	-	0.53	-	-	0.13	4.00	-	-	5.58	0.07	-	0.40	13.95	10.10	35.49	534
	T	-	0.03	2.42	1.12	0.03	0.44	-	-	0.20	0.82	-	-	5.41	0.03	-	1.09	14.91	10.79	37.29	1096
Madhya Pradesh	M	-	0.07	0.11	0.22	0.07	0.36	-	0.04	0.43	0.25	0.11	3.09	0.25	0.22	0.07	0.22	5.53	-	11.04	307
	F	-	0.04	0.26	0.15	0.11	0.26	-	-	0.42	0.04	0.08	2.99	0.30	0.15	0.19	0.04	4.62	-	9.65	255
	T	-	0.06	0.18	0.18	0.09	0.31	-	0.02	0.42	0.15	0.09	3.04	0.28	0.18	0.13	0.13	5.08	-	10.34	562
Maharashtra	M	-	0.17	1.08	0.12	-	0.50	0.04	-	0.25	0.33	0.08	1.38	5.43	0.12	-	0.63	4.63	5.20	19.96	479
	F	0.04	0.15	0.49	0.15	-	0.49	-	-	0.19	0.41	-	1.24	4.26	0.07	-	0.41	2.67	4.13	14.70	391
	T	0.02	0.16	0.80	0.14	-	0.51	0.02	-	0.22	0.39	0.04	1.35	4.98	0.10	-	0.53	3.72	4.80	17.78	870
Orissa	M	-	0.08	1.23	-	0.08	0.46	-	-	1.08	0.54	-	1.00	2.69	0.15	-	0.38	5.38	4.62	17.69	230
	F	-	0.08	0.83	0.45	-	0.15	-	-	0.15	0.15	0.08	1.35	1.80	-	-	0.22	3.62	2.56	11.44	152
	T	-	0.08	1.03	0.23	0.04	0.30	-	-	0.61	0.34	0.04	1.18	2.24	0.08	-	0.30	4.49	3.57	14.53	382
Punjab	M	0.66	0.46	1.84	0.86	-	1.12	-	0.13	0.13	0.33	0.07	6.92	1.71	1.12	-	0.92	-	4.41	22.46	341
	F	0.15	0.51	0.51	0.07	-	0.81	-	-	0.22	0.81	0.07	5.36	1.61	0.59	-	-	-	3.89	15.63	213
	T	0.42	0.49	1.21	0.49	-	0.97	-	0.07	0.17	0.56	0.07	6.18	1.67	0.87	-	0.49	-	4.17	19.25	554

+ M = male, F = female, T = total

Table (1A)Contd. : Incidence rate of temporary ailments by type of ailments separately by sex for different states and all-India, rural households : NSS 28th round

states	sex		type of ailments																	number of sample ailments	
	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)		(21)
	M	-	0.37	0.15	0.15	-	0.37	0.08	-	0.60	-	-	3.21	0.68	0.37	-	0.76	2.69	1.65	11.08	148
	F	-	0.16	0.48	0.08	-	0.08	-	-	0.32	-	-	3.26	0.89	0.32	-	0.24	1.35	2.08	9.26	116
	T	-	0.27	0.31	0.12	-	0.23	0.04	-	0.46	-	-	3.23	0.78	0.35	-	0.51	2.04	1.86	10.20	204
	M	-	0.20	1.55	0.20	-	0.54	-	-	0.40	0.14	-	0.14	1.55	-	-	0.74	10.88	3.84	20.18	299
	F	-	0.06	1.70	0.26	-	0.39	-	-	0.79	0.13	-	-	0.72	-	-	0.26	11.66	5.13	21.10	321
	T	-	0.13	1.63	0.23	-	0.46	-	-	0.60	0.14	-	0.07	1.13	-	-	0.50	11.27	4.49	20.65	620
	M	-	0.03	0.35	0.35	0.03	0.45	-	-	0.61	0.19	-	0.77	1.17	0.29	-	0.69	4.34	-	9.27	348
	F	0.06	0.18	0.24	0.15	0.03	0.18	-	0.03	0.39	-	0.03	0.75	0.99	0.09	-	0.27	2.94	-	6.33	211
	T	0.03	0.10	0.30	0.25	0.03	0.32	-	0.01	0.51	0.10	0.01	0.76	1.16	0.20	-	0.49	3.68	-	7.95	559
	M	-	0.26	1.36	0.51	-	0.17	0.08	0.04	0.26	0.09	0.04	0.09	4.44	0.04	-	1.07	3.07	0.81	12.37	290
	F	0.09	0.27	1.26	0.63	-	0.04	-	-	0.27	0.13	0.05	0.09	4.05	-	-	0.27	2.48	0.85	10.26	228
	T	0.04	0.26	1.32	0.57	-	0.11	0.04	0.02	0.26	0.11	0.04	0.09	4.25	0.02	-	0.68	2.79	0.83	11.35	518
	M	0.03	0.12	0.84	0.28	0.03	0.40	0.02	0.01	0.43	0.18	0.02	1.16	2.25	0.18	0.01	0.56	4.76	2.25	13.53	4675
	F	0.03	0.11	0.64	0.26	0.02	0.26	0.00	0.00	0.33	0.15	0.03	1.09	2.06	0.09	0.02	0.22	3.90	2.34	11.55	3937
	T	0.03	0.12	0.74	0.27	0.02	0.33	0.01	0.01	0.38	0.17	0.03	1.13	2.16	0.13	0.02	0.39	4.34	2.29	12.57	8612

* including smaller states and union territories. M = male, F = female, T = total

Table (1U) : Incidence rate of temporary ailments by type of ailments separately by sex for different states and all-India, urban households : NSS 28th round

states	sex	type of ailments																	number of sample ailments							
		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)		(18)	(19)	(20)	(21)	(22)		
Andhra Pradesh	M																									
	F																									
	T																									
Assam	M																									
	F																									
	T																									
Bihar	M																									
	F																									
	T																									
Gujarat	M																									
	F																									
	T																									
Haryana	M																									
	F																									
	T																									
Jammu & Kashmir	M																									
	F																									
	T																									

+ M = male, F = female, T = total

Table (1 U) Contd : Incidence rate of temporary ailments by type of ailments separately by sex for different states and all-India, urban households, NSS 28th round

states	sex	type of ailments																	number of sample ailments		
		(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)		(20)	(21)
	M		0.15	0.62	0.31		0.15						0.75	0.61	0.31		0.31	4.73	0.30	8.39	55
	F		0.33			0.16	0.16			0.17		0.51	0.32					3.13	0.50	6.45	39
	T		0.23	0.33	0.16	0.16	0.16		0.08	0.08		1.11	0.47	0.16				3.96	0.40	7.46	94
	M			1.13	0.49		0.50		1.19	0.21	0.21			1.92				10.11	4.97	21.08	298
	F		0.07	1.25	0.22	0.30	0.30		0.73	0.14			1.76					12.26	3.98	20.93	286
	T		0.04	1.18	0.36	0.40	0.40		0.96	0.18	0.11		1.84					11.17	4.48	21.01	584
	M	0.06	0.11	0.17	0.06		0.39		0.50	0.11			0.33	0.77	0.06			3.26		6.54	118
	F		0.13	0.27		0.20	0.20		0.66	0.07			0.20	1.40	0.07			2.79		5.99	90
	T	0.03	0.12	0.21	0.03	0.30	0.30		0.57	0.09			0.27	1.06	0.06			3.05		6.27	208
	M	0.08	0.30	1.80	0.30		0.07		0.23	0.30	0.30			4.37				6.26	2.71	18.60	247
	F	0.10	0.10	1.43	0.28	0.10	0.10		0.38	0.38	0.10			4.48				4.88	2.57	15.65	164
	T	0.09	0.21	1.64	0.30	0.08	0.08		0.30	0.33	0.21			4.42				5.63	2.65	17.30	411
	M	0.03	0.17	0.78	0.23	0.01	0.29		0.39	0.15	0.09		0.73	2.10	0.06			5.33	2.74	13.87	2306
	F	0.03	0.21	0.81	0.20	0.21	0.21		0.49	0.13	0.06		0.69	2.22	0.05			4.97	2.67	13.14	1855
	T	0.03	0.19	0.79	0.22	0.01	0.25		0.44	0.14	0.08		0.71	2.15	0.05			5.16	2.70	13.53	4161
	M																				
	F																				
	T																				

* including smaller states and union territories

+ M = male, F = female, T = total

Table (1U) Contd : Incidence rate of temporary ailments by type of ailments separately by sex for different states and-India, urban households : NSS 28th round

states	SEX		type of ailments																	number of sample	
	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)		(21)
Karnataka	M		0.12	0.72		0.12	0.72			0.12	0.49		0.36	1.20	0.12		0.24	1.68	2.87	8.76	73
	F			0.48	0.24		0.12			0.36			0.12	1.21				1.09	2.07	5.69	47
	T		0.06	0.60	0.12	0.06	0.42			0.24	0.24		0.24	1.21	0.06		0.12	1.39	2.48	7.24	120
Kerala	M			0.84	0.66		0.17			0.36	0.17	0.66		6.10	0.18		2.89	11.29	8.96	32.28	190
	F			3.43	0.67		0.34			0.18	1.01	0.50		6.70			0.18	11.41	10.80	35.22	205
	T			2.13	0.67		0.26			0.27	0.59	0.59		6.40	0.09		1.54	11.35	9.88	33.77	395
Madhya Pradesh	M	0.11	0.21	0.53	0.21		0.21			0.53	0.11		2.54	0.64	0.21		0.53	6.89		12.72	120
	F	0.23	0.70	0.35	0.35		0.47	0.12	0.23	0.58			1.17	0.58	0.12		0.12	4.78		9.80	84
	T	0.17	0.44	0.44	0.28		0.33	0.06	0.11	0.56	0.06		1.89	0.61	0.17		0.33	5.89		11.34	204
Maharashtra	M		0.26	1.16	0.47		0.31		0.05	0.21	0.05	0.05	0.95	3.74			0.47	7.99	5.99	21.70	413
	F		0.25	0.92	0.37		0.12	0.06		0.19	0.06	0.06	0.74	3.72			0.49	5.37	5.75	18.10	293
	T		0.26	1.05	0.43		0.23	0.03	0.03	0.20	0.06	0.06	0.85	3.73			0.48	6.78	5.87	20.06	706
Orissa	M		0.22	2.93	0.45		0.45			1.13			0.45	2.48			0.90	6.08	4.51	19.60	87
	F			1.59						1.59			0.53	1.86			0.53	2.92	5.32	14.34	54
	T		0.12	2.32	0.24		0.24			1.34			0.49	2.19			0.73	4.63	4.89	17.19	141
Punjab	M	0.19		0.58			0.77			0.19	0.38	0.19	3.64	1.73	0.19	0.19	0.19	4.99	1.53	14.95	78
	F	0.22	0.87	0.43	0.43		0.65			0.22			6.72	2.60	1.08		0.43	2.60	2.38	18.63	86
	T	0.20	0.41	0.51	0.20	0.20	0.71			0.20	0.20	0.10	5.09	2.14	0.61	0.10	0.31	3.87	1.93	16.68	164

+ M = male, F = female, T = total

Table (2R): Incidence rate of temporary ailments by age separately by sex for different states and all-India, rural households; NSS 28th round.

s t a t e s--		age (years)							not recor- ded	all ages	number of sample ailments
		0	1-4	5-14	15-24	25-44	45-59	60 & above			
(1)		(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
1. Andhra Pradesh	male	26.10	39.31	14.67	19.82	21.27	24.59	23.88		21.92	492
	female	22.68	33.72	15.90	17.56	19.64	20.47	15.62		19.84	436
	total	24.40	36.55	15.26	18.67	20.47	22.46	19.90		20.89	928
2. Assam	male	4.88	12.01	8.65	5.67	7.85	23.66	14.51		9.43	130
	female	14.64	9.57	7.64	5.03	7.18	6.95	6.83		7.34	91
	total	8.76	10.94	8.16	5.34	7.57	12.52	11.69		8.46	221
3. Bihar	male	10.65	10.50	3.87	3.41	5.63	5.35	6.86		5.48	154
	female	4.83	8.88	2.10	2.10	4.26	4.85	3.62		3.90	106
	total	7.83	9.69	3.06	2.78	4.92	5.09	5.21		4.69	260
4. Gujarat	male	14.76	3.87	3.11	0.87	5.62	9.39	2.83		4.18	52
	female	3.95	8.37	4.47	2.88	3.71	7.56	4.41		4.79	56
	total	9.48	6.24	3.74	1.83	4.69	8.46	3.61		4.48	108
5. Haryana	male	17.18	13.14	6.03	6.52	12.35	14.02	16.82		10.05	164
	female	14.92	12.25	4.53	6.36	11.53	7.78	11.31		8.35	121
	total	16.12	12.70	5.33	6.43	11.94	10.97	14.69		9.23	285
6. Jammu & Kashmir	male	21.19	21.54	8.66	6.29	9.92	12.86	10.35		10.89	313
	female	15.34	17.96	10.18	7.06	11.58	14.62	18.58		11.98	308
	total	18.32	19.83	9.39	6.55	10.68	13.71	13.55		11.42	621
7. Karnataka	male	14.36	13.17	6.40	4.59	11.13	11.93	6.14		8.65	122
	female	8.52	12.40	6.84	5.53	8.79	8.16	11.57		8.20	113
	total	11.43	12.77	6.62	5.06	9.93	10.04	8.69		8.44	235
8. Kerala	male	64.15	82.25	41.36	30.98	28.75	31.78	19.60		39.25	562
	female	51.29	70.34	36.44	25.79	32.34	32.99	20.19		35.49	534
	total	57.63	76.46	38.90	28.26	30.68	32.41	19.91		37.29	1096
9. Madhya Pradesh	male	24.77	12.94	8.18	9.69	11.39	12.89	13.55		11.04	307
	female	14.11	18.97	7.62	6.04	7.56	8.99	15.32		9.65	255
	total	19.53	15.95	7.91	7.91	9.57	10.86	14.36		10.34	562
10. Maharashtra	male	32.32	37.82	15.20	12.02	18.93	25.05	20.29		19.96	479
	female	29.96	26.74	13.85	9.03	12.93	17.30	11.80		14.70	391
	total	32.22	33.23	13.01	10.66	16.36	21.52	16.54		17.78	870
11. Orissa	male	34.58	31.03	15.92	14.01	17.65	12.88	12.16		17.69	230
	female	9.63	14.97	11.54	6.89	14.28	7.60	13.97		11.44	152
	total	22.80	23.04	13.70	10.36	15.97	10.01	13.01		14.53	382

Table (2R) Contd. : Incidence rate of temporary ailments by age separately by sex for different states and all-India, rural households : No 28th round.

RURAL

s t a t e s		age (years)							all ages	number of sample ailments	
		0	1-4	5-14	15-24	25-44	45-59	60 & not above recorded			
(1)		(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
12. Punjab	male	48.05	36.43	19.87	18.41	18.26	18.55	26.51		22.46	341
	female	24.12	22.11	12.72	12.46	16.16	15.55	20.08		15.63	213
	total	37.02	29.78	16.58	15.59	17.23	17.08	23.81		19.25	554
13. Rajasthan	male	13.40	13.75	8.00	13.09	10.62	10.51	16.47		11.08	148
	female	20.17	7.44	7.69	10.65	8.18	11.73	10.81		9.26	116
	total	16.68	10.80	7.84	11.92	9.40	11.15	13.76		10.20	264
14. Tamil Nadu	male	28.24	32.58	16.22	12.87	22.27	19.02	24.63		20.18	299
	female	42.67	28.05	12.95	16.08	23.01	25.74	30.27		21.10	321
	total	35.53	30.36	14.59	14.52	22.65	22.47	27.31		20.65	620
15. Uttar Pradesh	male	18.46	14.39	6.33	6.30	8.68	11.89	12.63		9.27	348
	female	4.00	11.90	4.16	4.63	6.17	7.54	8.12		6.33	211
	total	11.69	13.22	5.36	5.53	7.45	9.67	10.60		7.95	559
16. West Bengal	male	33.43	18.94	10.45	11.01	9.45	10.72	17.88		12.37	290
	female	31.25	15.36	9.04	5.82	7.89	13.63	14.19		10.26	228
	total	32.39	17.21	9.76	8.52	8.70	12.22	16.21		11.35	518
17. all-India*	male	24.08	22.36	10.57	10.44	12.88	14.82	14.91	157.82	13.53	4675
	female	17.88	18.81	9.46	8.63	11.20	12.55	12.31	32.04	11.55	3937
	total	21.09	20.65	10.04	9.53	12.04	13.64	13.67	103.98	12.57	8612

*including smaller states and Union territories.

SNC..

Table (2U) : Incidence rate of temporary ailments by age separately by sex for different states and all-India, urban households ; NSS 28th round.

URBAN

s t a t e s		age (years)							all number ages of sample ailments	number of sample ailments	
		0	1-4	5-14	15-24	25-44	45-59	60 & not above recor- ded			
(1)		(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
1. Andhra Pradesh	male	19.35	27.80	17.38	14.38	18.60	13.96	14.42		17.80	232
	female	54.88	44.25	11.89	11.52	18.71	13.53	16.60		18.62	235
	total	37.62	36.02	14.67	13.01	18.65	13.74	15.59		18.20	467
2. Assam	male		7.30	1.64	5.84	5.23	13.79			5.15	24
	female		23.83	6.53	9.07	1.17	3.40	5.74		7.23	26
	total		15.22	3.94	7.25	3.65	9.65	2.79		5.94	50
3. Bihar	male	19.24	12.44	5.70	4.67	6.45	7.10	6.11		6.87	70
	female	21.16	5.21	3.15	4.76	6.62	6.95	1.96		5.23	45
	total	20.15	8.99	4.48	4.71	6.52	7.02	3.99		6.05	115
4. Gujarat	male	28.68	0.45	0.96	2.83	4.03	3.79	10.47		2.47	31
	female	6.54	4.19	3.20	3.40	3.45	4.15	2.49		3.53	25
	total	18.21	1.35	1.55	3.09	3.76	3.96	6.38		2.85	56
5. Haryana	male	8.20	17.72	6.99	10.47	16.02	13.02	9.39		11.73	57
	female	20.83	4.29	4.35	1.17	6.30	6.65			4.73	20
	total	15.04	11.30	5.74	6.28	11.56	9.88	5.58		8.46	77
6. Jammu & Kashmir	male	23.48	10.03	7.31	7.27	9.18	11.18	20.78		9.32	106
	female	25.11	14.83	6.35	5.71	10.75	10.14	8.56		9.04	90
	total	24.26	12.30	6.86	6.53	9.90	10.69	15.62		9.22	196
7. Karnataka	male	25.14	23.84	5.48	3.84	8.22	10.78	6.76		8.76	73
	female		12.46	2.78	2.94	7.92	6.37	7.87		5.69	47
	total	13.03	18.14	4.14	3.41	8.08	8.52	7.34		7.24	120
8. Kerala	male	97.20	72.93	33.10	19.75	27.71	21.66	17.94		32.28	190
	female	114.18	76.46	34.62	21.46	28.99	29.37	34.94		35.22	205
	total	104.95	74.59	33.82	20.60	28.37	25.54	27.53		33.77	395
9. Madhya Pradesh	male	20.50	18.60	12.16	7.90	12.08	16.42	14.56		12.72	120
	female	35.00	18.72	6.15	4.42	13.17	10.06	3.90		9.80	84
	total	27.03	18.69	9.27	6.33	12.54	13.36	9.06		11.34	204
10. Maharashtra	male	45.41	38.00	17.86	16.94	21.42	24.65	16.21		21.70	413
	female	22.81	32.98	17.85	13.94	17.57	16.32	10.57		18.10	293
	total	35.37	35.52	17.85	15.65	19.77	20.86	13.25		20.06	706
11. Orissa	male	47.20	28.40	21.38	11.33	17.54	19.07	22.42		19.60	87
	female	43.48	21.69	11.93	10.37	8.44	23.26	20.58		14.34	54
	total	45.45	25.33	17.01	10.92	13.52	21.22	21.46		17.19	141

Table(2U) Contd. : Incidence rate of temporary ailments by age separately by sex for different states and all-India, urban households : NSS 28th round.

URBAN

s t a t e s		age (years)							60 & not above recor- ded	all ages of sample ailments	number of ailments
		0	1-4	5-14	15-24	25-44	45-59	(8)			
(1)		(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
12. Punjab	male	34.19	33.47	13.77	11.04	14.56	5.91	12.04		14.95	78
	female	28.57	37.22	16.80	11.21	20.21	12.36	21.98		18.63	86
	total	31.12	35.22	15.25	11.11	17.15	9.05	16.53		16.68	164
13. Rajasthan	male	25.16	16.76	3.27	7.85	10.97	4.92	8.30		8.39	55
	female	5.79	8.22	4.69	5.41	3.84	9.35	17.84		6.45	39
	total	15.08	12.47	3.95	6.77	8.59	7.17	13.30		7.46	94
14. Tamil Nadu	male	41.96	41.96	19.12	9.50	24.12	15.40	18.82		21.08	298
	female	20.06	34.88	14.10	14.42	24.18	24.60	26.93		20.93	286
	total	31.19	38.48	16.63	11.97	24.14	20.01	23.07		21.01	584
15. Uttar Pradesh	male	2.57	7.84	4.53	6.48	6.03	8.09	10.31		6.54	118
	female	16.54	7.89	6.49	4.05	5.38	4.94	5.25		5.99	90
	total	13.76	7.85	5.43	5.46	5.75	6.63	8.04		6.27	208
16. West Bengal	male	71.95	52.43	17.92	11.59	11.28	23.18	14.41		18.60	247
	female	42.59	39.20	16.00	9.73	13.35	8.90	12.93		15.65	164
	total	58.09	46.07	16.99	10.76	12.09	17.38	13.67		17.30	411
17. all-India*	male	33.69	23.92	11.40	9.61	13.87	14.36	13.09		13.87	2306
	female	26.61	24.86	10.61	8.96	13.30	12.36	12.86		13.14	1855
	total	30.29	24.36	11.02	9.30	13.64	13.41	12.97		13.53	4161

SNC.

Table (3R) : Prevalence rate of temporary ailments by type of ailments separately by sex for different states and all-India, rural households : NSS 28th round

states	sex +	type of ailments																			
		cholera	typhoid	dysentery (all forms)	diarrhoea	diphtheria	whooping cough	tetanus	acute poliomyelitis	small pox	measles	mumps	malaria	influenza	pneumonia	food poisoning	accident	others	not recorded	all types of ailments	number of sample ailments
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)
Andhra Pradesh	M	-	0.36	0.88	0.18	-	0.53	0.05	-	0.96	0.18	-	3.05	6.58	0.22	0.04	0.75	10.28	8.79	32.85	737
	F	-	0.22	0.62	0.05	-	0.63	-	-	0.80	0.18	0.09	3.21	5.61	0.14	0.05	0.40	8.71	10.24	30.95	681
	T	-	0.29	0.75	0.12	-	0.58	0.02	-	0.88	0.18	0.05	3.13	6.10	0.18	0.05	0.58	9.51	9.52	31.94	1418
Assam	M	0.14	0.44	3.56	0.37	0.50	0.72	0.07	-	0.94	-	0.07	1.91	5.94	0.14	-	0.29	1.16	3.30	19.55	270
	F	-	0.16	1.86	0.40	0.16	0.40	-	-	0.73	-	0.08	0.98	6.53	0.08	-	0.40	0.97	2.25	15.00	186
	T	0.08	0.31	2.75	0.38	0.34	0.57	0.04	-	0.83	-	0.08	1.47	6.21	0.11	-	0.34	1.07	2.82	17.40	456
Bihar	M	0.07	0.07	1.13	0.46	-	0.50	-	-	0.71	0.14	-	0.32	1.49	0.18	-	0.43	5.85	0.35	11.70	330
	F	0.04	0.11	0.81	0.22	-	0.40	0.04	0.04	0.51	0.04	-	0.18	0.99	0.04	-	0.04	4.76	0.40	8.62	235
	T	0.05	0.09	0.97	0.34	-	0.45	0.02	0.02	0.61	0.09	-	0.25	1.24	0.11	-	0.23	5.31	0.38	10.16	565
Gujarat	M	0.08	0.16	0.16	0.32	-	1.13	-	-	0.24	-	0.08	2.91	0.80	0.08	-	0.97	1.04	1.37	9.34	116
	F	0.08	0.26	0.26	0.09	-	0.52	-	-	0.34	-	0.08	3.51	1.62	0.08	-	0.60	1.29	2.32	11.05	129
	T	0.08	0.21	0.21	0.21	-	0.83	-	-	0.29	-	0.08	3.20	1.20	0.08	-	0.78	1.17	1.83	10.17	245
Haryana	M	0.12	0.31	0.37	0.31	-	1.23	-	-	0.31	0.43	-	1.78	0.74	0.55	0.06	0.37	2.70	7.67	16.95	276
	F	0.14	0.34	0.48	0.28	0.07	0.69	-	-	0.83	0.28	-	1.10	0.28	0.14	-	0.07	1.93	8.25	14.88	216
	T	0.13	0.32	0.42	0.29	0.03	0.97	-	-	0.55	0.36	-	1.46	0.52	0.36	0.03	0.23	2.34	7.95	15.96	492
Jammu & Kashmir	M	-	0.70	2.16	1.01	0.07	0.87	-	-	0.03	0.07	-	0.99	6.06	0.38	0.03	0.35	3.44	1.53	17.69	508
	F	0.08	0.63	2.14	1.09	-	1.08	-	-	0.08	0.16	0.04	0.67	7.00	0.23	0.04	0.12	3.09	1.44	17.89	460
	T	0.04	0.66	2.15	1.05	0.04	0.97	-	-	0.06	0.11	0.02	0.84	6.50	0.31	0.04	0.24	3.27	1.49	17.79	968

+ M = male, F = female, T = total

Table (3R) Contd. : Prevalence rate of temporary ailments by type of ailments separately by sex for different states and all-India, rural households : NSS 28th round

states	sex +	type of ailments																			
		cholera	typhoid	dysentery (all forms)	diarrhoea	diphtheria	whooping cough	tetanus	acute poliomyelitis	small pox	measles	mumps	malaria	influenza	pneumonia	food poisoning	accident	others	not recorded	all types of ailments	number of sample ailments
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)
Karnataka	M	-	0.14	1.35	-	-	1.13	-	0.07	0.14	-	0.14	0.14	1.56	-	0.24	0.35	3.76	3.09	16.08	227
	F	-	0.07	0.65	-	-	1.16	-	0.07	0.29	0.07	0.15	0.15	1.16	-	-	0.36	3.12	6.46	13.71	189
	T	-	0.11	1.00	-	-	1.15	-	0.07	0.22	0.04	0.14	0.14	1.36	-	0.11	3.36	3.44	6.78	14.92	416
Kerala	M	0.14	0.07	3.77	1.05	0.07	1.96	-	-	0.84	0.98	0.07	0.14	7.26	0.14	-	3.56	30.73	23.81	74.59	1068
	F	0.13	0.20	3.46	1.73	-	1.73	-	-	0.20	1.46	0.07	-	6.78	0.27	-	1.79	27.44	22.72	67.98	1023
	T	0.14	0.14	3.61	1.40	0.03	1.84	-	-	0.51	1.23	0.07	0.07	7.01	0.20	-	2.66	29.04	23.26	71.21	2091
Madhya Pradesh	M	-	0.36	0.25	0.32	0.14	0.79	-	0.04	0.72	0.97	0.14	4.95	0.47	0.39	0.07	0.36	11.84	0.04	21.85	609
	F	-	0.11	0.26	0.23	0.11	0.79	0.04	-	0.91	0.42	0.08	4.96	0.49	0.26	0.19	0.11	10.86	-	19.82	524
	T	-	0.24	0.26	0.28	0.13	0.79	0.02	0.02	0.81	0.70	0.11	4.96	0.48	0.33	0.13	0.24	11.37	0.02	20.89	1123
Maharashtra	M	-	0.75	1.58	0.21	-	1.34	0.04	0.08	0.46	0.46	0.17	2.34	7.14	0.21	-	1.13	7.83	7.91	31.65	758
	F	0.04	0.38	0.83	0.15	-	1.62	0.04	-	0.27	0.49	0.04	2.07	5.88	0.08	-	0.75	4.92	6.34	23.90	635
	T	0.02	0.57	1.19	0.18	-	1.48	0.04	0.04	0.36	0.47	0.10	2.19	6.48	0.14	-	0.93	6.31	7.09	27.57	1393
Orissa	M	-	0.38	1.77	0.15	0.08	1.15	-	0.08	1.15	0.62	-	2.46	3.39	0.15	-	0.92	8.85	7.93	29.08	378
	F	-	0.08	1.58	0.45	-	0.22	-	-	0.38	0.37	0.16	1.73	2.33	-	0.08	0.52	6.18	5.34	19.42	258
	T	-	0.23	1.67	0.30	0.04	0.68	-	0.04	0.77	0.49	0.08	2.09	2.85	0.08	0.04	0.72	7.50	6.62	24.20	636
Punjab	M	0.66	0.72	2.24	0.92	-	1.45	-	0.20	0.46	0.53	0.07	8.96	2.50	1.32	-	1.12	2.77	6.52	30.44	462
	F	0.15	0.66	0.95	0.15	-	2.35	0.07	-	0.51	1.10	0.07	6.53	2.57	0.88	0.07	0.22	1.91	6.02	24.21	330
	T	0.42	0.69	1.63	0.56	-	1.87	0.03	0.10	0.49	0.80	0.07	7.81	2.53	1.11	0.03	0.69	2.36	6.28	27.47	792

+ M = male, F = female, T = total

Table (3R) Contd. : Prevalence rate of temporary ailments by type of ailment separately by sex for different states and all-India, rural households : NSS 26th round

state	sex +	type of ailments																			
		cholera	typhoid	dysentery (all forms)	diarrhoea	diphtheria	whooping cough	tetanus	acute poliomyelitis	small pox	measles	mumps	malaria	influenza	pneumonia	food poisoning	accident	others	not recorded	all types of ailments	number of sample ailments
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)
Rajasthan	M	0.08	1.05	0.30	0.15	-	0.89	0.08	-	0.75	-	-	4.26	0.82	0.74	-	1.13	5.36	3.67	19.28	258
	F	0.08	0.87	0.48	0.08	-	0.47	-	0.08	0.56	-	-	4.38	1.28	0.72	-	0.48	2.77	3.13	15.38	193
	T	0.08	0.96	0.39	0.12	-	0.69	0.04	0.04	0.66	-	-	4.29	1.04	0.73	-	0.81	4.10	3.38	17.33	451
Tamil Nadu	M	0.13	0.40	2.78	0.27	-	0.88	-	0.07	1.21	0.34	-	0.14	2.16	-	-	1.01	7.86	7.37	34.62	512
	F	-	0.19	2.77	0.59	-	0.79	-	-	1.18	0.13	-	-	1.84	-	-	0.39	7.44	8.02	33.34	507
	T	0.07	0.30	2.78	0.43	-	0.83	-	0.03	1.19	0.24	-	0.07	2.00	-	-	0.70	7.65	7.70	33.99	1019
Uttar Pradesh	M	-	0.29	0.53	0.51	0.03	0.93	-	0.03	0.77	0.56	-	1.04	1.62	0.29	-	1.36	7.27	-	15.23	572
	F	0.06	0.30	0.51	0.42	0.03	0.60	-	0.06	0.48	0.03	0.03	0.96	1.50	0.12	0.03	0.42	5.40	-	10.95	365
	T	0.03	0.30	0.52	0.47	0.03	0.78	-	0.04	0.63	0.31	0.01	1.00	1.57	0.21	0.01	0.92	6.39	-	13.22	937
West Bengal	M	-	1.02	3.62	1.75	-	0.47	0.13	0.04	0.55	0.17	0.09	0.13	7.03	0.17	-	2.44	0.54	2.43	30.58	717
	F	0.18	0.50	3.32	1.12	-	0.50	-	-	0.54	0.13	0.05	0.14	5.62	-	-	1.26	8.77	2.61	24.74	550
	T	0.09	0.77	3.47	1.45	-	0.48	0.06	0.02	0.55	0.15	0.07	0.13	6.35	0.08	-	1.86	9.66	2.52	27.71	1267
* All-India	M	0.05	0.42	1.46	0.50	0.04	0.90	0.02	0.03	0.70	0.37	0.05	1.85	3.26	0.26	0.02	1.09	8.96	4.13	24.11	8219
	F	0.05	0.28	1.21	0.41	0.02	0.78	0.01	0.02	0.55	0.24	0.05	1.69	2.93	0.14	0.03	0.51	7.55	4.23	20.70	6918
	T	0.05	0.35	1.34	0.46	0.03	0.84	0.02	0.03	0.63	0.31	0.05	1.77	3.10	0.20	0.03	0.80	8.27	4.18	22.46	15137

* including smaller states and union territories.

+ M = male, F = female, T = total

Table (3U) : Prevalence rate of temporary ailments by type of ailment separately by sex for different states and all-India, urban households : NSS 28th round

states	type of ailments																				
	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)
sex +	M	F	T	M	F	T	M	F	T	M	F	T	M	F	T	M	F	T	M	F	T
Andhra Pradesh	-	-	0.54	0.92	-	-	0.15	-	0.08	0.77	0.15	0.08	2.45	3.76	-	-	1.61	8.44	9.74	28.69	374
Assem	-	-	0.63	0.87	0.16	-	0.71	-	0.08	1.43	0.16	0.16	2.22	3.72	-	-	1.66	7.68	7.45	26.93	340
Bihar	-	-	0.58	0.90	0.08	-	0.43	-	0.08	1.09	0.16	0.12	2.34	3.74	-	-	1.64	8.07	8.61	27.84	714
Gujarat	-	-	0.42	1.72	0.42	-	-	-	-	0.42	-	-	-	3.65	-	-	-	0.85	2.53	10.01	47
Haryana	-	-	-	2.49	-	-	-	-	-	0.82	-	-	1.13	1.93	-	-	0.27	5.24	1.91	13.79	50
Jammu & Kashmir	-	-	0.24	2.06	0.24	-	-	-	-	0.59	-	-	0.49	2.89	-	-	0.12	2.76	2.27	11.66	97
All-India	-	-	0.19	0.97	0.19	-	0.29	-	-	0.68	-	0.29	0.29	1.94	0.10	-	0.48	6.01	0.78	12.21	126
	-	-	0.12	0.58	0.35	-	0.12	-	0.12	0.46	0.12	-	0.23	2.08	-	-	0.12	4.86	1.39	10.55	91
	-	-	0.16	0.79	0.26	-	0.21	-	0.05	0.58	0.05	0.16	0.26	2.00	0.05	-	0.32	5.48	1.05	11.42	217
	-	-	0.08	0.08	-	-	0.24	-	0.08	0.16	0.08	-	1.27	0.80	0.08	-	0.40	0.96	0.80	5.03	63
	0.14	0.05	0.70	-	-	2.28	-	-	-	0.29	-	0.29	2.68	0.71	-	-	0.14	1.41	1.83	8.47	60
	0.05	0.30	0.30	0.05	-	0.26	-	-	0.05	0.21	0.05	0.10	1.78	0.76	0.05	-	0.31	1.12	1.17	6.26	123
	0.21	0.62	0.62	0.82	0.21	0.62	0.71	0.66	-	0.21	-	-	1.23	0.82	0.41	-	2.26	7.18	4.72	19.31	94
	-	-	-	0.71	0.24	0.71	0.66	0.66	0.24	-	-	0.24	0.24	0.71	0.47	-	0.47	6.37	2.83	13.23	56
	0.11	0.33	0.33	0.77	0.22	0.66	0.66	0.66	0.11	0.11	-	0.11	0.77	0.77	0.44	-	1.43	6.80	3.84	16.47	150
	-	-	0.26	0.53	0.62	-	0.27	-	0.09	-	-	-	0.09	6.72	0.18	-	0.35	2.22	1.33	12.66	144
	0.10	0.61	0.10	1.11	0.61	0.30	0.30	0.30	-	0.10	-	-	0.20	7.82	0.10	-	0.40	1.02	1.02	12.98	129
	0.05	0.61	0.19	0.80	0.61	0.28	0.28	0.28	0.05	0.05	-	-	0.14	7.28	0.14	-	0.37	1.66	1.18	12.80	273

+ M = male, F = female, T = total

Table (30) Contd. : Prevalence rate of temporary ailments by type of ailment separately by sex for different states and all-India, urban household : NSS 28th round

states	sex +	type of ailments																			
		cholera	typhoid	dysentery (all forms)	diarrhoea	diphtheria	whooping cough	tetanus	acute poliomyelitis	small pox	measles	mumps	malaria	influenza	pneumonia	food poisoning	accident	others	not recorded	all types of ailments	number of sample ailments
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)
Karnataka	M	-	0.12	1.32	-	0.12	1.08	0.12	-	0.61	0.49	-	0.36	2.53	0.12	-	0.36	3.13	4.56	14.92	124
	F	-	-	0.60	0.24	-	0.85	-	0.12	0.61	-	-	0.12	2.92	-	-	-	2.05	4.25	11.76	97
	T	-	0.06	0.97	0.12	0.06	0.97	0.06	0.06	0.61	0.24	-	0.24	2.73	0.06	-	0.18	2.59	4.40	13.35	221
Kerala	M	-	0.36	2.02	1.00	0.17	1.53	-	-	0.36	0.52	0.83	-	7.79	0.18	-	5.30	21.18	21.99	63.23	371
	F	-	0.18	4.31	1.18	-	1.02	-	-	0.35	1.59	0.50	-	9.30	-	-	0.89	19.01	22.00	60.44	351
	T	-	0.27	3.61	1.09	0.08	1.28	-	-	0.35	1.11	0.67	-	8.54	0.09	-	3.11	20.10	21.99	61.84	722
Madhya Pradesh	M	0.11	0.64	1.17	0.21	-	0.85	0.11	0.11	0.95	0.11	0.71	3.71	1.17	0.32	-	1.06	12.93	-	23.56	222
	F	0.23	1.40	0.47	0.70	-	1.28	0.12	0.35	0.82	-	-	3.62	0.70	0.47	-	0.23	9.33	-	19.72	169
	T	0.17	1.00	0.83	0.44	-	1.06	0.11	0.22	0.89	0.06	0.06	3.66	0.94	0.39	-	0.67	11.22	-	21.72	391
Maharashtra	M	-	0.68	1.63	0.53	0.05	1.11	-	0.10	0.42	0.32	0.21	1.53	6.22	0.05	-	1.63	12.35	8.66	35.49	675
	F	-	0.50	1.29	0.37	-	0.31	0.06	0.06	0.37	0.12	0.06	1.05	5.89	-	-	0.99	8.95	8.28	28.30	458
	T	-	0.60	1.48	0.46	0.03	0.74	0.03	0.08	0.40	0.23	0.14	1.31	6.06	0.03	-	1.33	10.78	8.48	32.18	1133
Orissa	M	-	0.22	3.38	0.68	-	0.45	-	-	2.03	0.45	-	1.80	2.93	-	-	1.13	9.69	6.76	29.52	131
	F	-	0.53	1.59	-	-	0.27	-	-	2.12	-	-	1.06	1.86	-	-	0.53	3.98	6.38	18.32	69
	T	-	0.37	2.56	0.37	-	0.37	-	-	2.07	0.24	-	1.46	2.44	-	-	0.85	7.07	6.58	24.38	200
Punjab	M	0.38	0.38	0.96	-	0.19	1.34	-	-	0.19	0.77	0.19	4.80	2.11	0.58	0.38	0.58	6.71	3.64	23.20	121
	F	0.22	1.08	0.65	0.43	-	1.95	-	-	0.87	-	-	7.37	3.90	1.52	-	0.65	4.33	3.25	26.22	121
	T	0.31	0.71	0.81	0.20	0.10	1.63	-	-	0.51	0.41	0.10	6.00	2.95	1.02	0.20	0.61	5.60	3.46	24.62	242

+ M = male, F = female, T = total

Table (3U) contd. : Prevalence rate of temporary ailments by type of ailment separately by sex for different states and all-India, urban households : NSS 28th round

states	sex		type of ailments																	number of ailments	all types of ailments
	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)		
Rajasthan	M	-	0.15	0.62	0.31	-	0.46	-	-	0.45	-	-	1.20	0.61	0.46	-	0.63	6.41	1.66	12.96	85
	F	-	0.82	0.32	0.33	-	0.33	-	-	0.19	0.17	-	1.67	0.32	0.34	-	0.50	5.28	1.85	12.12	73
	T	-	0.47	0.48	0.32	-	0.39	-	-	0.33	0.08	-	1.43	0.47	0.40	-	0.57	5.86	1.75	12.55	153
Tamil Nadu	M	-	0.07	1.83	0.70	-	0.85	0.07	-	1.89	0.49	0.21	-	2.55	-	-	0.71	15.51	6.7	11.62	448
	F	-	0.22	1.61	0.43	-	0.51	0.07	-	1.83	0.58	0.14	-	2.05	-	-	0.44	17.72	7.20	32.35	443
	T	-	0.14	1.73	0.57	-	0.68	0.07	-	1.64	0.53	0.18	-	2.30	-	-	0.58	16.50	6.96	31.98	891
Uttar Pradesh	M	0.05	0.33	0.39	0.17	-	0.50	-	0.05	0.66	0.39	-	0.44	1.49	0.06	-	1.10	5.52	-	11.23	203
	F	-	0.40	0.33	-	0.20	-	0.20	-	0.86	0.20	-	0.33	1.66	0.13	-	0.40	5.25	-	9.76	147
	T	1.03	0.36	0.36	0.09	-	0.36	-	0.03	0.75	0.30	-	0.39	1.57	0.09	-	0.78	5.40	-	10.54	350
West Bengal	M	0.08	0.68	3.99	0.30	-	0.45	0.06	-	1.28	0.75	0.30	0.15	7.45	0.07	0.53	2.35	11.84	5.35	35.78	475
	F	0.10	0.57	2.48	0.86	-	0.29	-	0.08	1.34	0.67	0.19	0.10	7.07	0.10	0.18	2.01	10.40	4.96	31.40	329
	T	0.08	0.63	3.32	0.55	-	0.38	0.03	0.04	1.31	0.71	0.25	0.13	7.26	0.08	0.38	2.32	11.21	5.18	33.85	804
* All-India	M	0.03	0.38	1.41	0.30	0.03	0.65	0.03	0.04	0.77	0.33	0.14	1.13	3.25	0.31	0.06	1.29	8.78	4.66	23.39	3849
	F	0.04	0.50	1.15	0.37	-	0.53	0.02	0.06	0.82	0.27	0.10	1.17	3.14	0.12	0.02	0.68	8.21	4.76	22.06	3104
	T	0.04	0.44	1.29	0.33	0.01	0.60	0.03	0.05	0.79	0.30	0.12	1.15	3.24	0.11	0.04	1.01	8.51	4.71	22.77	6953

* including smaller states and union territories. + M = male, F = female, T = total

Table (4R): Prevalence rate of temporary ailments by age separately by sex for different states and all-India, rural households : 28th round

RURAL,

s t a t e s	a g e (y e a r s)									
	0	1-4	5-14	15-24	25-44	45-59	60 & above	not recorded	all ages	number of sample ailments
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
Andhra	male	36.66	52.49	21.70	27.22	32.60	40.05	47.23	32.85	737
pradesh	female	36.46	46.29	22.45	26.18	31.59	37.14	35.45	30.95	681
	total	36.56	49.40	22.09	26.70	32.17	38.55	41.56	31.94	1418
Assam	male	7.32	22.52	15.77	14.86	16.89	55.67	31.68	19.55	270
	female	25.62	19.08	12.99	10.89	15.67	18.08	20.48	15.00	186
	total	14.61	21.00	14.44	12.83	16.39	30.60	27.56	17.40	456
Bihar	male	16.75	18.75	8.83	7.73	12.32	13.54	15.82	11.70	330
	female	11.27	15.46	5.94	3.98	8.68	12.00	11.87	8.62	235
	total	14.08	17.13	7.48	5.90	10.43	12.74	13.82	10.16	565
Gujarat	male	18.44	11.65	7.25	5.29	10.55	17.09	8.54	9.34	116
	female	11.65	21.63	7.77	6.75	10.40	15.08	13.18	11.05	129
	total	15.12	16.89	7.50	5.99	10.48	16.08	10.83	10.17	245
Haryana	male	28.64	20.90	10.12	12.36	20.92	20.69	30.45	16.95	276
	female	29.83	19.17	8.15	9.52	19.10	22.64	20.11	14.88	216
	total	29.23	20.07	9.21	11.02	19.99	21.61	26.43	15.96	492
Jammu & kashmir	male	29.08	31.45	12.85	8.81	15.55	23.61	29.60	17.69	508
	female	26.40	24.80	13.91	12.00	17.53	23.24	28.99	17.89	460
	total	27.76	28.24	13.36	10.29	16.49	23.42	29.36	17.79	968
Karnataka	male	25.86	18.18	9.96	11.29	18.56	26.84	24.58	16.08	227
	female	17.04	17.98	8.36	8.09	15.73	18.55	28.90	13.71	189
	total	21.44	18.07	9.19	9.70	17.12	22.69	26.62	14.92	416
Kerala	male	134.12	166.35	72.81	55.33	59.50	57.58	40.23	74.59	1068
	female	108.27	147.37	67.83	47.48	59.74	59.64	43.26	67.98	1023
	total	121.02	157.16	70.31	51.23	59.66	58.63	41.81	71.21	2091
Madhya Pradesh	male	43.32	23.66	16.71	17.48	22.31	28.56	30.42	21.85	609
	female	28.24	33.80	16.16	13.10	18.74	17.66	28.10	19.82	524
	total	35.91	28.73	16.45	15.36	20.63	22.90	29.24	20.89	1133
Maharashtra	male	43.64	58.40	24.13	20.49	28.42	41.21	38.22	31.65	759
	female	48.88	41.93	19.69	13.45	23.11	30.46	25.94	23.90	635
	total	46.29	50.07	21.90	16.47	25.62	35.28	32.02	27.57	1393

Contd.

Table (4R) / Prevalence rate of temporary ailments by age separately by sex for different states and all-India, rural households : NSS 28th Round

states		age (years)									number of sample ailments
		0	1-4	5-14	15-24	25-44	45-59	60 & above	not recorded	all ages	
(1)		(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
Orissa	male	43.23	43.18	24.02	25.13	27.42	25.02	42.04		29.08	378
	female	16.08	27.91	19.94	11.48	20.32	15.84	27.95		19.42	258
	total	30.40	35.50	21.95	18.13	23.88	20.02	35.48		24.20	636
Punjab	male	66.37	50.77	24.51	23.02	24.24	29.98	41.10		30.44	462
	female	37.52	33.15	18.42	18.28	24.73	31.10	31.04		24.21	330
	total	53.08	42.60	21.68	20.76	24.50	30.55	36.85		27.47	792
Rajasthan	male	18.66	22.68	14.46	17.75	19.52	25.49	28.64		19.28	258
	female	20.17	16.15	11.70	24.18	16.31	17.23	19.24		15.35	193
	total	19.39	19.62	13.17	17.00	17.91	21.18	24.15		17.33	451
Tamil Nadu	male	41.21	53.17	24.33	19.29	40.00	37.07	55.37		34.62	512
	female	55.25	42.90	19.86	24.27	37.98	39.93	56.02		33.34	507
	total	48.32	48.11	22.12	21.88	38.96	38.54	55.14		33.99	1019
Uttar Pradesh	male	26.38	21.89	9.58	10.57	16.04	19.48	22.43		15.23	572
	female	14.99	20.17	6.15	7.54	10.94	12.18	16.62		10.95	365
	total	21.06	21.08	8.04	9.14	13.53	15.77	19.83		13.22	937
West Bengal	male	74.86	47.34	25.20	23.18	26.39	30.73	44.72		30.58	717
	female	66.05	38.76	19.82	15.42	23.84	27.26	35.46		24.74	550
	total	70.65	43.19	22.57	19.44	24.87	28.94	40.29		27.71	1267
All-India*	male	38.71	37.83	18.24	18.00	23.44	28.30	31.32	157.82	24.11	8219
	female	32.55	33.07	15.98	14.78	20.76	22.97	25.96	210.95	20.70	6918
	total	35.72	35.50	17.15	16.38	22.10	25.51	28.78	180.54	22.46	15137

including Smaller states and union Territories.

Table (4U) : Prevalence rate of temporary ailments by age separately by sex for different states and all-India, urban households : NSS 28th round

states		URBAN									
		a g e (years)							all ages	number of sample ailments	
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)			(10)
Andhra	male	48.40	43.47	25.79	19.69	31.32	22.99	36.86		28.69	374
Pradesh	female	70.14	61.39	17.54	16.45	25.62	26.31	26.28		26.93	340
	total	59.56	52.42	21.72	18.13	28.62	24.72	31.19		27.84	714
Assam	male	12.77	16.82	4.11	9.69	7.46	18.28	32.60		10.01	47
	female		29.00	13.13	15.30	7.04	6.80	28.71		13.79	50
	total	6.10	22.65	8.34	12.14	7.29	13.71	30.70		11.66	97
Bihar	male	33.66	22.95	9.26	8.41	11.01	16.23	12.23		12.21	126
	female	21.16	13.56	7.88	6.80	13.70	15.06	3.92		10.55	91
	total	27.71	18.48	8.60	7.76	12.20	15.67	7.99		11.42	217
Gujarat	male	34.55	2.71	2.10	5.09	8.55	6.34	23.50		5.03	63
	female	6.54	5.59	6.41	6.14	11.40	13.77	10.02		8.47	60
	total	21.30	3.40	3.24	5.56	9.88	9.91	16.59		6.26	123
Jammu & Kashmir	male	23.48	14.14	10.06	9.98	12.58	15.88	26.99		12.66	144
	female	25.10	17.64	8.67	7.80	16.12	19.36	17.20		12.98	129
	total	24.26	15.82	9.40	8.98	14.19	17.53	22.85		12.80	273
Karnataka	male	50.41	30.65	10.53	6.07	14.06	24.33	13.54		14.92	124
	female	10.77	16.95	7.89	6.46	17.36	10.13	17.63		11.76	97
	total	31.32	23.78	9.22	6.26	15.69	17.01	15.72		13.35	221
Kerala	male	171.36	142.92	62.00	38.69	49.08	54.96	53.63		63.23	371
	female	133.80	138.39	59.62	40.38	45.77	54.29	63.01		60.44	351
	total	154.40	140.76	60.84	39.53	47.38	54.61	58.91		61.84	722
Madhya Pradesh	male	32.79	35.14	22.04	15.78	24.59	26.69	22.87		23.56	222
	female	65.00	33.27	13.11	6.96	23.27	27.91	19.52		19.72	169
	total	47.29	34.24	17.74	11.78	23.95	27.26	21.15		21.72	391
Maharashtra	male	72.85	62.02	29.33	25.99	33.12	45.67	34.72		35.49	675
	female	34.19	49.14	25.27	21.23	29.64	27.63	22.16		28.30	458
	total	55.72	55.67	23.37	23.94	31.64	37.45	28.13		32.18	1133

Table (4U) Contd. : Prevalence rate of temporary ailments by age separately by sex for different states and all-India, urban households : NSS 28th round

URBAN

s t a t e s	a g e (y e a r s)									n u m b e r o f s a m p l e a i l m e n t s
	0	1-4	5-14	15-24	25-44	45-59	60 & a b o v e	n o t r e c o r - d e d	a l l a g e s	
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
Orissa	male	66.04	38.54	33.36	14.74	26.73	43.60	22.42	29.52	131
	female	43.48	28.92	14.91	11.85	11.60	31.01	28.81	18.32	69
	total	55.55	34.14	24.83	13.49	20.04	37.13	25.75	24.38	200
Punjab	male	51.29	40.91	17.82	15.28	27.61	23.63	15.05	23.20	121
	female	35.71	43.78	21.00	17.30	30.32	26.79	29.31	26.22	124
	total	42.89	42.25	19.37	16.20	29.68	25.16	21.49	24.62	242
Rajasthan	male	31.63	18.17	6.54	11.33	16.46	12.88	19.60	12.96	85
	female	23.16	18.19	8.80	7.31	9.21	15.63	27.83	12.12	73
	total	27.22	18.18	7.62	9.55	14.76	14.28	23.92	12.55	158
Tamil Nadu	male	67.14	58.13	28.13	16.13	34.26	26.00	34.87	31.62	448
	female	43.18	56.35	20.32	21.84	36.87	37.24	43.89	32.35	443
	total	55.35	57.28	24.28	18.97	35.19	31.63	39.61	31.98	891
Uttar Pradesh	male	33.40	11.76	8.87	8.10	10.44	15.64	17.19	11.23	203
	female	18.91	11.52	9.14	6.62	10.78	9.88	10.50	9.76	147
	total	26.38	11.63	8.97	7.49	10.60	12.97	14.18	10.54	350
West Bengal	male	127.14	97.84	33.16	20.92	25.73	41.48	30.28	35.78	475
	female	80.50	66.40	30.91	20.37	27.11	24.01	34.51	31.40	329
	total	107.37	82.73	32.07	20.67	26.26	34.38	32.40	33.85	804
All-India	male	56.96	38.87	18.88	15.26	23.39	27.02	26.05	23.39	3849
	female	41.10	39.25	17.28	14.56	22.80	23.28	25.06	22.06	3104
	total	49.34	39.04	18.14	14.97	23.13	25.24	25.57	22.77	6953

* including smaller states and union Territories.

Table (5.R): Average duration (days) of temporary ailments terminating in recovery by type of ailment separately by sex for different states and all-India, rural households : NSS 28th round

states	sex +	type of ailments																			
		cholera	typhoid	dysentery (all forms)	diarrhoea	diphtheria	whooping cough	tetanus	acute poliomyelitis	small pox	measles	mumps	malaria	influenza	pneumonia	food poisoning	accident	others	not recorded	all types of ailments	number of sample ailments
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)
Andhra Pradesh	M		19.23	4.12			8.38			6.64	13.94		6.89	7.16	4.00		6.33	7.50	6.25	7.25	266
	F			5.40						7.17	7.00		9.83	6.84				6.43	4.97	6.60	250
	T		19.23	4.65			8.38			6.89	12.23		8.27	7.00	4.00		6.33	7.05	5.46	6.94	516
Assam	M	9.50	8.00	6.44	5.00	7.00	16.00			13.33		10.00	5.89	6.78	9.50		8.67	5.83	9.00	7.28	106
	F		8.00	13.75	6.00	10.00	8.00			14.60			6.33	7.20	8.00			5.00	4.65	8.33	78
	T	9.50	8.00	9.08	5.40	7.75	4.00			14.13		10.00	6.07	6.98	9.00		8.67	5.83	6.08	7.72	184
Bihar	M	3.00	4.00	9.11	3.88		14.57			10.00	7.00		7.20	6.50	10.00		6.75	1.26	10.00	9.36	137
	F	5.00	20.00	7.00	4.67		32.40			8.25	7.00		16.00	7.07				8.08	9.00	9.86	77
	T	4.00	14.67	8.27	4.09		22.00			9.22	7.00		10.50	6.69	10.00		6.75	10.09	9.33	9.54	214
Gujarat	M	5.00			3.00		8.33						6.79	4.00			8.64	2.00	7.16		42
	F	4.00					8.00			10.50			7.71	5.10			2.50	6.24	2.50		54
	T	4.50			3.00		8.25			10.50			7.34	4.92			6.20	5.38	5.60		96
Haryana	M	4.00	20.00	10.00	6.33		6.67			10.00	4.00		8.60	4.33	4.33	3.00	2.00	6.42	12.98		86
	F	1.00		6.50	22.33					10.63	6.50		6.83	8.00				5.00	8.96		55
	T	2.50	20.00	8.25	14.33		6.67			16.90	5.43		7.94	5.25	4.33	3.00	2.00	6.06	11.39		141
Jammu & Kashmir	M		10.85	9.58	5.95		11.66			8.00	6.00		9.60	5.87	18.64	4.00	10.00	8.22	11.00		218
	F	4.00	11.67	5.29	4.94		11.96				6.32		9.01	6.35	10.00			7.99	11.53		189
	T	4.00	11.08	6.35	5.47		11.81			8.00	6.24		9.44	6.11	16.93	4.00	10.00	8.11	11.30		407

+ M = male, F = female, T = total

Table (5R) Contd : Average duration (days) of temporary ailments terminating in recovery by type of ailment separately by sex for different states and all-India, rural households : NSS 28th round

states	sex +	type of ailments																			
		cholera	typhoid	dysentery (all forms)	diarrhoea	diphtheria	whooping cough	tetanus	acute poliomyelitis	small pox	measles	mumps	malaria	influenza	pneumonia	food poisoning	accident	others	not recorded	all types of ailments	number of sample ailments
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)
Karnataka	M		7.00	11.12			4.50						6.00	5.44		15.00	20.00	6.50	4.47	6.74	65
	F			3.50			4.33		30.00			8.00	10.00	5.09			60.05	4.65	5.82	5.66	59
	T		7.00	9.60			4.40		30.00			8.00	8.00	5.25		15.00	9.50	5.61	5.11	6.13	124
Kerala	M	6.50		4.80	5.60		7.50			15.00	10.43		12.00	5.48			9.43	6.74	5.57	8.24	350
	F	6.50		5.93	3.59		10.50			20.00	11.07			6.85	12.00		15.00	6.24	6.33	6.66	321
	T	6.50		5.37	4.33		9.00			15.71	10.86		12.00	6.17	12.00		11.83	6.52	6.96	6.75	671
Madhya Pradesh	M		13.50	6.50	14.17		14.75			10.00	18.80		8.02	11.57	7.40		5.00	8.72		9.28	181
	F		14.00		7.00		10.00	8.00		8.38	7.00		8.41	9.50			5.00	8.59		8.56	109
	T		13.67	6.50	11.78		13.17	8.00		9.00	15.43		8.18	10.82	7.40		5.00	8.67		9.01	290
Maharashtra	M		15.60	4.83	6.50		9.80		45.00	14.24	9.25		7.17	7.86	10.77		6.00	5.85	7.05	7.58	265
	F		21.56	7.38	3.00		8.40			21.00	8.00		8.88	5.97			7.33	8.05	7.74	7.73	220
	T		17.30	5.85	5.80		9.10		45.00	16.49	8.63		8.00	6.97	10.77		6.75	6.80	7.39	7.65	485
Orissa	M		15.50	2.11	12.00	4.00	6.33			10.50	7.40		7.52	6.47	9.00		8.00	7.33	9.81	7.94	173
	F			11.00	4.75		7.00			6.00	8.00	8.00	8.80	6.25			12.00	6.50	11.73	9.08	117
	T		15.50	8.48	6.20	4.00	6.50			8.70	7.50	8.00	7.94	6.39	9.00		9.00	7.65	10.65	8.40	290
Punjab	M	3.88	11.33	4.07	3.78		13.00		14.00	11.20	8.00		6.26	5.65	8.17		3.00	6.00	6.96	6.42	191
	F	5.00	63.00	7.60	4.00		5.00	20.00		8.00	8.00	1.00	6.78	3.92	8.67		8.00	10.58	5.57	7.61	135
	T	4.00	32.00	5.00	3.80		10.71	20.00	14.00	9.78	8.00	1.00	6.46	4.90	8.42		5.50	8.50	6.20	6.91	326

+ M = male, F = female, T = total

Table(5R) Contd. : Average duration (days) of temporary ailments terminating in recovery by type of ailment separately by sex for different states and all-India, rural households : NSS 28th round

States	Sex +	type of ailments																			
		cholera	typhoid	dysentery (all forms)	diarrhoea	diphtheria	whooping cough	tetanus	acute poliomyelitis	small pox	measles	mumps	malaria	influenza	pneumonia	food poisoning	accident	others	not recorded	all types of ailments	number of sample ailments
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)
Rajasthan	M	2.00	28.58				3.00			13.32			13.81	8.00	4.76		6.00	7.36	9.49	11.13	78
	F												10.34	31.74	7.00		7.00	10.13	4.26	10.60	52
	T	2.00	28.47				3.00			13.32			12.20	22.31	5.21		6.50	8.43	7.66	10.92	130
Tamil Nadu	M	11.00	8.66	4.16	3.00		3.50			11.29	7.00		4.00	5.90			6.50	6.18	6.11	6.18	191
	F		19.00	5.48	7.41					10.59	9.00			7.35			2.00	5.11	7.24	6.36	198
	T	11.00	13.82	4.89	6.44		3.50			10.94	8.00		4.00	6.58			3.71	5.62	6.67	6.30	389
Uttar Pradesh	M		23.33	5.00	6.40		14.33			8.00	5.00		8.21	5.65	5.83		23.56	6.59		8.21	146
	F		33.50	6.00	5.00		6.50			4.67			8.26	7.70	3.50		9.86	9.53		8.90	88
	T		27.40	5.20	6.00		11.20			7.09	5.00		8.24	6.68	5.25		17.56	7.49		8.47	234
West Bengal	M		16.53	10.63	5.93		8.33	10.00		13.24	7.67	14.00	2.00	6.24	11.99		14.00	9.90	25.02	9.42	275
	F	8.50	27.40	6.31	4.64		9.20			14.25	5.00		4.50	6.12				8.15	6.63	7.54	178
	T	8.50	20.76	8.95	5.52		8.72	10.00		13.75	7.00	14.00	3.67	6.19	11.99		14.00	9.12	15.33	8.68	453
* All-India	M	5.49	15.92	6.65	5.96	4.38	9.55	10.00	19.98	10.50	10.06	12.48	8.11	6.61	7.98	12.95	12.07	8.15	6.89	7.88	2817
	F	6.06	24.81	7.19	5.25	10.00	12.26	12.26	30.00	10.13	9.05	6.73	8.68	6.72	7.76		8.20	11.23	5.66	8.52	2240
	T	5.70	18.84	6.89	5.66	5.15	10.52	10.87	22.97	10.33	9.61	9.02	8.36	6.66	7.94	12.95	10.73	9.45	6.27	8.16	5057
	M																				
	F																				
	T																				

* Includes smaller states and union territories.

+ M = male, F = female, T = total

Table (5U) : Average duration (days) of temporary ailments terminating in recovery by type of ailment separately by sex for different states and all-India, urban households :
NSS 28th round

states	sex	type of ailments																			
		cholera	typhoid	dysentery (all forms)	diarrhoea	diphtheria	whooping cough	tetanus	acute poliomyelitis	small pox	measles	mumps	malaria	influenza	pneumonia	food poisoning	accident	others	not recorded	all types of ailments	number of sample ailments
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)
Andhra Pradesh	M		24.00	2.50					7.00	9.83	7.00		5.91	5.70			1.50	6.96	5.20	6.35	165
	F		14.40	3.75	1.50				7.00	6.88	5.00		5.24	5.26			11.71	15.26	5.43	8.59	133
	T		18.00	3.33	1.50				7.00	8.14	6.00		5.63	5.46			9.44	10.46	5.27	7.35	298
Assam	M			3.67										6.73				6.00	15.07	8.01	21
	F			7.35										7.00				10.17		7.94	8
	T			5.53										6.99				8.12	15.07	7.99	29
Bihar	M			3.00	10.00		9.00			12.00			5.00	4.11			7.50	4.82	18.00	6.44	54
	F			15.00	3.00						7.00		13.00	4.60			4.00	6.91	10.00	6.65	40
	T			9.00	5.33		9.00			12.00	7.00		8.20	4.37			6.33	5.71	16.67	6.53	94
Gujarat	M		10.00	2.33						10.00	15.00		5.66	4.99			4.32	9.00	3.53	5.85	28
	F		7.00									4.00	8.27	9.00			3.00	7.00	12.94	8.51	24
	T		8.49							10.00	15.00	4.00	7.10	6.32			4.00	7.80	7.27	7.07	52
Haryana	M		33.00	2.33									6.25	15.00	17.00		10.00	4.86	5.88	8.43	28
	F											2.00		2.00	3.00			5.88	6.64	5.33	15
	T		33.00	2.33								2.00	6.25	10.67	10.00		10.00	5.40	6.64	7.35	43
Jammu & Kashmir	M			7.46	6.03								5.00	4.90	12.00			5.80	5.00	5.32	63
	F			7.34	4.00					7.00				4.44				9.08	7.77	5.60	54
	T			7.37	5.35					7.00			5.00	4.68	12.00			6.89	6.16	5.45	117

Table (5U) Contd. : Average duration (days) of temporary ailments terminating in recovery by type of ailment separately by sex for different states and all-India, urban households :
NSS 28th round

states	sex +	type of ailments																						
		cholera	typhoid	dysentery (all forms)	diarrhoea	diphtheria	whooping cough	tetanus	acute poliomyelitis	small pox	measles	mumps	malaria	influenza	pneumonia	food poisoning	accident	others	not recorded	all types of ailments	number of sample ailments			
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)			
Karnataka	M			3.76		7.00	4.47			8.00	8.00		3.00	6.83			4.97	8.08	5.47	6.26	47			
	F			3.49	3.00		4.50			7.50			5.00	4.74				5.36	3.43	4.57	33			
	T			3.67	3.00	7.00	4.48			7.80	8.00		4.01	5.63			4.97	6.10	4.57	5.56	80			
Kerala	M			6.23	7.00						7.00	9.60		4.69			9.80	6.93	7.83	7.07	129			
	F			4.13	7.80						9.40	11.00		5.91			4.00	5.29	8.81	6.45	141			
	T			4.00	7.50						9.09	10.00		5.42			9.33	6.25	8.35	6.75	270			
Madhya Pradesh	M		13.00	4.50	7.00		10.00	10.00	10.00	4.00			7.00	5.40	6.00					7.41	70			
	F	2.00	21.00		9.33		13.50		4.00	7.50			4.92	4.50	11.00		10.00	7.10		7.45	49			
	T	2.00	17.00	4.50	8.40		12.33	10.00	7.00	5.75			6.04	5.14	8.50		10.00	7.30		7.31	119			
Maharashtra	M		14.23	6.43	3.34		8.10		30.00	7.00	11.53		7.64	6.98	8.00					6.57	5.40	4.09	5.92	317
	F		16.33	6.38	1.50			1.00	25.00	7.00			6.19	6.15			12.00	4.86	5.14	5.68	201			
	T		14.13	6.41	2.61		8.10	1.00	27.50	7.00	11.53		7.29	6.66	8.00		8.20	5.19	4.52	5.82	518			
Orissa	M			3.33	7.50		3.00			10.00	10.00		8.00	5.28						5.56	5.80	6.29	62	
	F		15.00	3.20						9.00			4.00	4.33						7.75	5.50	5.92	26	
	T		15.00	3.29	7.50		3.00			9.75	10.00		7.00	5.00						6.00	5.67	6.18	88	
Punjab	M	7.50	30.00	3.33			2.00				8.00		7.54	4.67	3.00					4.67	6.50	6.30	40	
	F	8.00	8.50		6.00		4.00			11.50			5.05	5.18	8.25					3.25	5.25	5.56	50	
	T	7.67	15.67	2.50	6.00		2.50			11.50	8.00		6.86	5.07	7.20					4.23	5.88	5.89	90	

Table (5U) Contd. : Average duration (days) of temporary ailments terminating in recovery by type of ailments separately by sex for different states and all-India, urban households:

states	sex +	type of ailments																			number of sample ailments
		cholera	typhoid	dy-sentery (all forms)	diarrhoea	diphtheria	whooping cough	tetanus	acute poliomy-elities	small pox	measles	mumps	malaria	influenza	pneumonia	food poisoning	accident	others	not recorded	all types of ailments	
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)
Rajasthan	M			1.00	3.00		6.00			8.00			11.86	3.00	5.00			6.54		6.71	27
	F						4.00				4.00		3.27	6.00	3.00			5.35	6.43	4.93	22
	T			1.00	3.00		5.00			8.00	4.00		7.50	4.01	4.00			6.06	5.43	5.90	49
Tamil Nadu	M			6.97	2.25		5.00	10.00		11.39	7.16	7.00		5.01			1.67	5.59	4.69	5.64	192
	F		15.66	5.20	9.34		8.23	2.00		12.85	9.60	8.50		4.84			2.66	5.33	5.69	6.15	189
	T		15.66	6.04	5.27		7.14	6.00		12.15	8.37	1.19		4.93			2.17	5.46	5.15	5.89	381
Uttar Pradesh	M		30.00	3.50				15.00		8.00			11.50	7.18			12.67	6.95		8.33	52
	F		16.50	7.75						7.63			6.00	6.53	7.00		7.00	6.50		7.11	57
	T		21.00	6.33				15.00		7.82			9.14	6.79	7.00		10.40	6.79		7.69	109
West Bengal	M		14.33	5.63	5.33		6.01			15.00	8.55		5.50	5.30		12.43	5.82	11.06	5.64	7.70	213
	F	3.00	18.20	5.60	5.00		8.50			11.83	6.60	7.00	11.00	6.50		6.00	7.78	11.64	5.92	8.29	135
	T	3.00	16.09	5.62	5.14		7.01			13.54	7.85	7.00	7.33	5.84		11.14	6.43	11.25	6.18	7.93	348
* all-India	M	7.50	17.59	5.32	4.79	7.00	6.39	11.44	15.92	10.21	8.57	9.02	6.91	5.78	6.89	12.43	6.76	6.90	5.05	6.58	1548
	F	4.09	15.33	5.57	5.64		7.84	1.56	14.59	10.03	7.98	7.67	6.09	5.75	7.85	6.00	8.06	7.04	6.23	6.80	1212
	T	5.32	16.43	5.44	5.23	7.00	6.92	7.34	15.34	10.13	8.31	8.27	6.54	5.76	7.46	11.14	7.20	6.96	5.55	6.68	2760

+ M = male, F = female, T = total * including smaller states and union territories.

Table (22): Average duration (days) of temporary ailments terminating in recovery by age separately by sex for different states and all-India, rural households : IJSS 28th round.

States		age (years)							not recorded	all ages	number of sample ailments
		0	1-4	5-14	15-24	25-44	45-59	60 & above			
(1)		(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
1. Andhra-Pradesh	male	7.32	5.57	7.53	7.85	7.60	7.81	7.97		7.25	266
	female	5.47	5.38	6.94	5.77	7.45	7.08	7.98		6.60	250
	total	6.40	5.49	7.26	6.49	7.53	7.43	7.98		6.94	516
2. Assam	male	5.00	8.21	7.10	6.71	7.17	5.91	8.00		7.28	106
	female	5.53	11.67	8.72	8.31	7.36	5.89	6.00		8.33	78
	total	5.25	9.50	7.82	7.40	7.61	5.90	7.33		7.72	184
3. Bihar	male	8.67	7.54	7.57	7.50	13.80	7.33	10.18		9.56	137
	female	16.20	11.21	11.71	5.50	7.31	9.20	8.50		9.36	77
	total	13.58	8.76	8.87	6.64	11.76	8.27	9.73		9.54	214
4. Gujarat	male	5.00	4.99	5.46	7.53	8.86	6.38			6.53	42
	female		6.91	5.86	7.75	6.86	7.53	8.00		6.53	54
	total	5.00	5.61	5.67	7.62	7.49	7.24	8.00		6.53	96
5. Haryana	male	5.53	4.71	7.38	5.80	7.67	6.42	34.55		9.91	86
	female	20.50	9.79	11.00	4.33	5.83	9.33	4.50		10.15	55
	total	17.71	7.00	9.00	5.46	6.75	7.00	29.92		10.00	141
6. Jammu & Kashmir	male	10.79	7.47	6.41	8.25	7.24	8.47	9.35		7.62	218
	female	6.66	5.32	7.27	3.28	6.89	7.63	10.16		7.25	189
	total	8.88	6.69	6.77	8.27	7.08	8.08	9.70		7.45	407
7. Karnataka	male	5.00	5.50	8.56	4.45	7.30	5.00	14.00		6.74	65
	female	2.00	3.82	3.64	6.50	5.88	7.00	11.00		5.66	59
	total	4.40	4.62	6.56	5.18	6.60	5.79	11.86		6.13	124
8. Kerala	male	3.77	6.53	6.13	6.77	8.24	5.33	6.86		8.24	350
	female	3.57	7.36	6.44	6.20	6.13	6.97	6.92		6.66	321
	total	8.70	6.92	6.27	6.51	10.00	6.88	6.42		6.75	671
9. Madhya-Pradesh	male	4.85	9.36	7.14	6.53	9.97	13.48	10.89		9.28	181
	female	11.35	7.94	6.85	7.46	7.71	9.92	13.25		8.56	109
	total	8.67	8.10	7.01	6.91	9.20	12.26	11.81		9.01	290

sd/

Table (6R) (Contd.) : Average duration (days) of temporary ailments terminating in recovery by age separately by sex for different states and all-India, rural households : NSS 20th round.

RURAL

States		age years							60 & above	not recorded	all ages	number of sample ailments
		0	1-4	5-14	15-24	25-44	45-59					
(1)		(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	
10. Maharashtra	male	4.50	6.89	7.91	7.35	9.08	6.48	9.02		7.58	265	
	female	7.68	6.65	8.47	6.93	8.74	6.18	9.21		7.73	220	
	total	5.74	6.78	8.17	7.16	8.93	6.35	9.11		7.65	485	
11. Orissa	male	.25	3.48	5.98	6.71	11.05	7.87	11.95		7.94	173	
	female	4.50	12.93	9.41	11.64	7.87	7.18	5.70		9.08	117	
	total	4.35	8.28	7.54	8.26	9.62	7.58	9.87		8.40	290	
12. Punjab	male	6.11	6.33	5.92	7.00	7.41	6.23	4.42		6.42	191	
	female	5.00	10.65	5.42	5.91	6.62	13.50	9.43		7.61	135	
	total	5.71	7.86	5.73	6.52	6.98	10.76	5.77		6.91	326	
13. Rajasthan	male	6.32	9.67	10.42	7.79	9.25	11.91	29.01		11.13	78	
	female		8.36	13.69	8.14	11.60	7.99	8.51		10.60	52	
	total	6.32	9.32	11.79	7.93	10.46	10.16	23.96		10.92	130	
14. Tamil Nadu	male	4.21	5.01	6.37	4.92	6.90	7.79	5.18		6.18	191	
	female	4.45	6.81	6.53	4.49	6.42	9.82	5.37		6.36	198	
	total	4.32	5.83	6.45	4.61	6.67	8.66	5.28		6.30	389	
15. Uttar Pradesh	male	5.67	6.33	8.73	9.05	8.58	6.81	11.60		8.21	146	
	female	5.50	6.96	12.42	6.85	10.11	7.00	14.43		8.90	88	
	total	5.60	6.65	9.79	8.23	9.09	6.88	12.76		8.47	234	
16. West Bengal	male	7.55	10.45	9.88	9.14	8.10	9.55	9.62		9.42	275	
	female	6.78	9.12	8.00	7.03	7.36	8.54	5.50		7.54	178	
	total	6.71	10.00	9.19	8.33	7.77	9.20	7.97		8.68	453	
17. all-India *	male	6.00	7.01	7.52	7.41	8.69	8.02	10.39	2.58	7.88	2817	
	female	7.88	7.61	7.89	13.53	7.41	7.86	8.00	11.34	8.52	2240	
	total	6.80	7.27	7.68	10.22	8.11	7.95	9.37	6.96	8.16	5057	

* including smaller states and Union Territories.

contd.

Table (6U) : Average duration (days) of temporary ailments terminating in recovery by age separately by sex for different states and all-India, urban households : NSS 28th round.

URBAN

States		age (years)						60 & above	not recorded	all ages of sample ailments	number of ailments
		0	1-4	5-14	15-24	25-44	45-59				
(1)		(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
10. Maharashtra	male	4.01	5.56	5.52	6.43	6.14	6.25	7.15		5.92	317
	female	10.40	5.20	5.7	6.74	5.21	5.07	7.00		5.68	201
	total	5.78	5.41	5.55	6.54	5.80	5.88	7.12		5.82	518
11. Orissa	male	8.67	5.13	6.83	4.50	6.26	7.29	3.00		6.29	62
	female	9.00	5.75	8.57	1.33	6.00	5.17	3.00		5.92	26
	total	8.80	5.33	7.32	3.44	6.24	6.31	3.00		6.18	88
12. Punjab	male	1.50	6.60	5.43	6.83	7.00	7.25	5.00		6.30	40
	female	5.67	5.71	7.00	4.10	5.08	4.75	8.00		5.56	50
	total	4.00	6.24	6.42	5.13	5.95	6.00	7.00		5.89	90
13. Rajasthan	male	10.00	4.84	11.74	3.61	9.19		3.99		6.71	27
	female	6.00	4.95	4.19	3.03		6.38	6.45		4.93	22
	total	8.67	4.89	7.04	3.41	9.19	6.38	5.63		5.90	49
14. Tamil Nadu	male	4.90	4.57	5.86	7.70	5.65	5.92	4.62		5.64	192
	female	8.92	5.37	7.08	5.79	5.08	7.30	7.28		6.15	189
	total	6.43	4.93	6.47	6.60	5.40	6.84	5.65		5.89	381
15. Uttar Pradesh	male	5.40	6.78	8.67	8.30	11.00	6.20	10.50		8.33	52
	female	7.50	8.83	6.85	5.18	8.64	6.50	7.00		7.11	57
	total	6.33	7.60	7.41	6.67	9.87	6.53	9.33		7.69	109
16. West Bengal	male	5.80	8.25	6.77	6.11	6.73	9.57	15.87		7.70	213
	female	7.12	6.60	10.37	6.95	6.55	10.25	9.76		8.29	135
	total	6.38	7.65	8.39	6.47	6.67	9.74	12.09		7.93	348
17. all-India*	male	5.19	6.28	5.88	6.52	6.80	8.20	7.85		6.58	1548
	female	6.73	6.07	7.02	6.27	6.09	6.99	12.32		6.80	1212
	total	5.85	6.18	6.40	6.41	6.51	7.67	9.99		6.68	2760

* including smaller states and Union Territories.

Table (7R) : Number of persons suffering from chronic diseases per 100,000 persons by type of chronic disease separately by sex for different states and all-India, rural households : NSS 28th round

states	type of chronic diseases																						
	tuberculosis	leprosy	syphilis	cancer	thyroid trouble or jointre	diabetes	mental illness	epilepsy	rheumatic fever	high blood pressure	bronchitis	asthma	peptic ulcer	kidney stone or kidney trouble	arthritis	rheumatism	stroke	piles	others	not recorded	all types	number of sample ailments	
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)	
1. Andhra Pradesh	m	126	91	13	13	9	17	9	22	34	26	27	277	73	22		189	13	17		774	1752	404
	f	59	23	9	26	9	9	13	13	35	83	13	231	40		177	31	13		867	1651	371	
	t	93	57	11	19	9	13	11	17	35	54	20	255	57	11		183	22	15		820	1702	775
2. Assam	m	51	36	14	14	43	65		14	36	79	79	281	194	116		158	14	51	554	144	1943	270
	f	24	9			56	16	16	8	40	16	16	88	129	40	8	153		32	459	72	1182	147
	t	38	23	8	8	49	42	8	11	38	49	49	190	163	80	4	155	8	42	509	110	1584	417
3. Bihar	m	172	63	18	4	14	25	21	39	18	25	25	549	151	25	11	141	11	120	1076	28	2536	720
	f	98	22		4	11	4	18	15	26	15	18	489	91	26	4	219	7	40	1028	11	2146	588
	t	136	43	9	4	13	14	20	27	21	20	21	519	122	25	7	179	9	81	1053	20	2343	1308
4. Gujarat	m	105	8		8	24	24		16	16	24	24	249	8	16	32	8	16	18	81	113	780	97
	f	26	17	9			8	9		8	17		162	34		17			9	42	60	418	49
	t	66	12	4	4	12	17	4	8	12	21	13	207	21	8	25	4	8	8	62	89	605	146
5. Haryana	m	54	24		6	12	18	18	12	90	18	60	409	30	30	26	138		120	968		2043	340
	f	107	7		20	7		13	33	60	20	20	354	27	7	107	87	7	87	1015		1978	296
	t	79	16		13	9	9	16	22	76	19	41	383	28	19	70	114	3	104	991		2012	636
6. Jammu & Kashmir	m	45	12	2	5	17	5	17	33	38	93	14	159	69	66		84	3	43		259	934	542
	f	63	4	2	6	23	17	4	17	64	99	23	91	86	31		90	6	16		321	963	500
	t	54	8	2	5	20	11	11	10	50	96	18	127	77	49		87	5	30		288	948	1042

m = male, f = female, t = total

Table (9A) Contd. : Number of persons suffering from chronic diseases per 100,000 by type of chronic disease separately by sex for different states and all-India, rural households : NSS 28th round

states	type of chronic diseases																				number of sample ailments		
	tuberculosis	leprosy	syphilis	cancer	thyroid trouble or joitre	diabetes	mental illness	epilepsy	rheumatic fever	high blood pressure	bronchitis	asthma	peptic ulcer	kidney stone or kidney trouble	arthritis	rheumatism	stroke	piles	others	not recorded		all types	
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)	
Karnataka	m	50	7	14	14	83	21	14	7	55	21	346	48	7		28	7	26		582	1330	192	
	f	43	14		14	21	14	14	14	43		257	29			22	14	14		392	919	129	
	t	46	11	7	7	14	53	18	14	11	49	18	302	39	4		25	11	21		487	1137	321
Kerala	m	584	21	7	55	41	309	21	110	103	151	288	1057	206	117		2052	21	584		2504	8231	1200
	f	505	26	7	66	85	269		131	236	144	72	1057	66	59	20	2710	20	164		2861	8498	1294
	t	544	23	7	60	64	289	10	121	171	148	178	1057	134	87	10	2389	20	369		2687	8368	2494
Madhya Pradesh	m	59	53	7	11	25	12	6	39	46	12	78	328	35	46	18	198	14	32	406		1425	413
	f	75	22		4	15	34	19	30	37	26	52	75	26	19	22	157	7	11	220		851	228
	t	80	38	4	7	20	24	15	34	42	20	65	205	31	33	20	178	11	22	316		1165	641
Maharashtra	m	127	119	8	12	16	37	20	16	21	12	12	352	114	33	94	41		65		736	1835	449
	f	67	59		8	16	4	12	12	8	28	12	279	51	16	130	28	4	24		639	1397	354
	t	96	88	4	10	16	20	16	14	14	20	12	315	82	24	112	34	2	44		686	1609	803
Orissa	m	53	61	15		23	46	69		76	38	53	274	160	107	23	343	38	61	1463	46	2949	387
	f	60		15	23	15	30	150	15	128	38	45	128	128	53	38	451	45	22	1172	60	2616	348
	t	57	30	15	11	19	38	110	8	102	38	49	200	144	79	30	397	42	42	1317	53	2781	735
Rajasthan	m	102		45	13	13	13	51	70	26	45	102	774	26	96	6	154	26	204	582		2348	367
	f	130		22		7	50	50	58	65	123	86	468	72	72	14	195	14	86	685		2197	305
	t	115		34	7	10	30	51	65	44	81	95	630	48	85	10	173	20	149	630		2277	672

m = male, f = female, t = total

Table (9R) Contd. : Number of persons suffering from chronic disease per 100,000 persons by type of chronic disease separately by sex for different states and all-India, rural households : NSS 28th round

states	type of chronic diseases																					number of sample ailments		
	tuberculosis	leprosy	syphilis	cancer	thyroid trouble or jointre	diabetes	mental illness	epilepsy	rheumatic fever	high blood pressure	bronchitis	asthma	peptic ulcer	kidney stone or kidney trouble	arthritis	rheumatism	stroke	piles	others	not recorded	all types			
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)		
13 Rajasthan	m	116	7	7	7	29	14	29	36		51	436	37	14	22	88	7	50		468	1425	196		
	f	47			8		23	16	16	47	24	8	272	8		16	116	8	15		493	1117	143	
	t	83	4	4	7	4	26	15	22	41	11	30	357	23	7	19	102	8	33		480	1276	339	
14 Tamil Nadu	m	235	128	13	13		47	14	40		53		242	174		81	13	167	33	53		1147	2453	367
	f	92	59	7	20	7	7	27	33	46	33	13	204	111		7	7	85	13	20		1217	2008	306
	t	163	93	10	17	3	26	20	36	23	43	7	223	142		43	10	125	23	37		1182	2226	673
15 Uttar Pradesh	m	129	31	10	10	16	26	10	29	31	8	34	485	50		21	3	137	10	81		464	1585	177
	f	74	23	3	21	27	35	9	27	65	12	26	345	27		24	6	159	9	15		486	1393	165
	t	103	28	7	15	21	30	10	28	47	8	31	419	39		22	4	147	10	50		474	1493	342
16 West Bengal	m	181	71	21	8	67	86	4	21	29	143	134	634	361		101	34	260	8	160	1137	458	3920	934
	f	58	36	4	4	45	18	15	36	36	138	27	299	98		90	27	368		62	1205	465	2969	665
	t	121	54	13	6	56	54	9	28	32	141	82	472	234		95	30	284	4	113	1170	461	3459	1599
17 * All-India	m	144	54	13	11	22	48	17	30	36	41	56	440	115		48	19	228	14	95	204	625	2260	7783
	f	89	25	4	14	21	30	21	27	54	47	25	309	60		26	26	275	12	33	184	647	1947	6480
	t	117	40	8	12	22	39	19	28	45	44	41	376	89		37	22	251	13	65	194	636	2098	14263

* including smaller states and union territories

m = male, f = female, t = total

Table (9U) : Number of persons suffering from chronic diseases per 100,000 persons by type of chronic disease separately by sex for different states and all-India, urban households : NSS 28th round

states	type of chronic diseases																					
	tuberculosis	leprosy	splifosis	cancer	thyroid trouble or goitre	diabetes	mental illness	epilepsy	rheumatic fever	high blood pressure	bronchitis	asthma	peptic ulcer	kidney stone or kidney trouble	arthritis	rheumatism	stroke	piles	others	not recorded	all types	number of sample ailments
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)
1. Andhra Pradesh	m	98	45	8	8	75	8	8	8	60	38	150	30	30	38	23	15		466	1100	146	
	f	31		8	8	47	8	8		109	16	186	8	8	47	47	23		637	1191	153	
	t	65	23	8	4	4	61	4	8	4	84	27	168	19	19	42	34	19		550	1143	299
2. Assam	m	63		21	63	104			21	229	42	229	231	42	125		63	812	168	2213	106	
	f	27		27	27	81		27	81	109	27	163	27	81	190			1060	136	2063	76	
	t	47		24	47	94		12	47	177	35	201	142	59	153		35	921	154	2148	182	
3. Bihar	m	230	57		10	67	29	48		29	29	527	67	48	105	29	67	910	10	2262	236	
	f	183		23	34	11	34	11		149	34	561	23	57	57	11	46	778	134	2046	179	
	t	209	31	10	21	42	31	31		83	31	542	47	52	83	21	57	850	21	2162	415	
4. Gujarat	m	162	12		12	50			12	37		211	12	13	12	12		164	88	797	64	
	f	56				28			14	70	14	98		14	14	14		56	238	616	44	
	t	112	6		7	39			13	53	7	158	7	13	13	13		113	158	712	108	
5. Haryana	m	121	20			60		40	20	80		623	141	101	20	80		161	1246		2713	135
	f	138		46	46		23	46		92	46	367	69		23	46	23	92	1171		2228	97
	t	129	11	21	21	32	11	43	11	86	21	504	107	54	21	64	11	129	1211		2487	232
6. Jammu & Kashmir	m	48	13		4	4	4	4	26	126	9	152	126	39	70	9	44		114	792	182	
	f	30	20		5	15	10	5	20	160		90	105	30	10	70		10	262	842	168	
	t	39	17		5	9	7	5	23	141	5	124	116	35	5	70	5	28	182	816	350	

MD. m = male, f = female, t = total

Contd.

Table (90) Contd. : Number of persons suffering from chronic diseases per 100,000 persons by type of chronic disease separately by sex for different states and all-India, urban households : NSS 28th round

states	type of chronic diseases																						
	tuberculosis	leprosy	syphilis	cancer	thyroid trouble or jointre	diabetes	mental illness	epilepsy	rheumatic fever	high blood pressure	bronchitis	asthma	peptic ulcer	kidney stone or kidney trouble	arthritis	rheumatism	stroke	piles	others	not recorded	all types	number of sample ailments	
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)	
Karnataka	m	128	23			129	23	12		58		453	47		12	35	23	59		288	1290	111	
	f	82	12		12	83	24	12		141		329				48	12			316	1071	91	
	t	105	17		6	106	23	12		100		392	24		6	41	18	29		302	1181	202	
Kerala	m	768	16		33	480	33	33	182	434	278	1824	248	181		1584		511		2260	8865	534	
	f	308	34		119	66	321		33	268	372	133	1725	135	116	18	2291	51	269	2565	8884	527	
	t	570	25		75	33	401	16	33	225	403	200	1775	192	149	9	1935	25	391	2411	8874	1061	
Madhya Pradesh	m	135	31		21	41	21	10	10	83		425	62	52	10	31		41	456		1429	138	
	f	113			23	45	11	11	23	23	113	11	204	34	23	23	57	11	23	522	1270	112	
	t	125	16		11	32	27	16	16	16	97	5	319	49	38	16	43	5	32	487	1350	250	
Maharashtra	m	190	31		26	103	10	15	5	180	31	330	72	21	16	21	10	46		463	1570	305	
	f	140	18		12	91	24	18	6	218	12	339	24	24	24	42		24		588	1610	265	
	t	167	25		20	97	17	17	6	197	23	334	50	22	20	31	6	36		520	1591	570	
Orissa	m	10	113		22	158	68	23		90		248	23	45	23	135	22	68	946	113	2180	97	
	f	53	53	53			264		26	212	26	132	53	106		344	53	26	952	106	2459	93	
	t	73	85	24	12	85	158	12	12	146	12	195	36	73	12	232	36	49	950	110	2312	190	
Punjab	m	75	56	19		19	37	19	37		130	130	391	19	56		56		37	540		1621	87
	f	107	65	21	43	62	22	43	21		215	86	407	43		22	257		107	472		1993	93
	t	190	60	20	20	40	30	30	30		169	110	398	30	30	10	149		70	508		1794	180

Table (9U) Contd. Number of persons suffering from chronic diseases per 100,000 persons by type of chronic disease separately by sex for different states and all-India, urban households ; NSS 28th round

states	type of chronic diseases																					number of sample ailments		
	tuberculosis	leprosy	syphilis	cancer	thyroid trouble or goitre	diabetes	mental illness	epilepsy	rheumatic fever	high blood pressure	bronchitis	asthma	peptic ulcer	kidney stone or kidney trouble	arthritis	rheumatism	stroke	piles	others	not recorded	all types			
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)		
Rajasthan	m	89	30	15	14	29	15			30	93			45				15		459	834	55		
	f	64				36	48	48	16	16	48	33	114	33			65		32		384	937	57	
	t	77	15	8	7	17	39	31	8	8	39	16	103	16	23		31		23		423	884	112	
Tamil Nadu	m	195	55	21	7	35	138	14	7	6	76	14	270	131	21		21	14	27		997	2049	296	
	f	86	22	21	14		29	14	14	7	138	22	223	57	7	15	79		14		1165	1927	268	
	t	142	39	21	11	18	85	14	10	7	106	18	247	95	14	7	49	7	21		1079	1990	564	
Uttar Pradesh	m	184	32	5	5	16	5	11	11	16	38	276	32	54		54	5	22		416		1182	77	
	f	143	13	20	26	7	26	19	20	32	78	26	176	33	72	7	84	13	7		449		1251	69
	t	165	23	12	15	3	21	12	15	21	44	32	230	32	62	3	68	9	15		431		1213	146
West Bengal	m	111	22		7	37	193	7	22	44	222	163	651	229	74	82	193	37	282		1052	496	3924	529
	f	47	19		19	28	38	9	47	104	348	132	301	122	75	132	319	9	103		1449	601	3902	414
	t	83	21		12	33	125	8	33	70	278	149	497	182	74	104	248	25	203		1227	542	3914	943
All-India	m	169	34	5	8	15	105	13	16	21	108	48	397	86	45	13	113	14	81		582	2003	3373	
	f	102	14	7	20	18	52	22	18	33	159	38	308	44	35	22	182	12	38		680	1931	2912	
	t	137	25	6	14	16	80	18	17	26	132	43	355	66	40	17	146	13	61		629	1962	6285	

* including smaller states and union territories.

m = male, f = female, t = total

Table (18R): Number of persons suffering from chronic diseases per 100,000 persons by age separately by sex for different states and all-India, rural households : NSS 28th round.

states		age (years)							number of sample cases
		0-14	15-25	25-44	45-59	60 & above	not recorded	all ages	
(1)		(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
1. Andhra Pradesh	male	241	388	1731	5346	8114	-	1752	404
	female	284	520	2618	4045	4633	-	1651	371
	total	261	611	2162	4666	6770	-	1702	775
2. Assam	male	265	650	2520	6770	9184	-	1943	270
	female	172	753	2245	3771	3868	-	1182	147
	total	216	702	2396	5379	7241	-	1584	417
3. Bihar	male	455	1084	3490	7204	9223	-	2536	720
	female	367	1088	3067	5784	5727	-	2146	588
	total	411	1084	3273	6466	7460	-	2343	1308
4. Gujarat	male	202	264	953	2035	4117	-	780	97
	female	-	97	593	1343	2182	-	418	49
	total	105	184	777	1687	3163	-	605	146
5. Haryana	male	549	941	2359	6602	7488	-	2043	340
	female	392	884	4040	5424	5119	-	1978	296
	total	476	914	3214	6034	6577	-	2012	636
6. Jammu & Kashmir	male	134	321	1073	2572	4093	-	934	542
	female	96	371	1867	2404	3043	-	963	500
	total	116	344	1443	2491	3684	-	948	1042
7. Karnataka	male	401	285	1168	4069	6375	-	1330	192
	female	417	202	1481	1593	2831	-	919	129
	total	409	288	1325	2840	4716	-	1137	321
8. Kerala	male	2054	2093	10325	22301	35649	-	8231	1200
	female	1805	3515	11457	24547	26859	-	8498	1294
	total	1931	2839	10927	23439	31099	-	8368	2494
9. Madhya Pradesh	male	313	553	1879	3891	5757	-	1425	413
	female	78	461	1329	2448	2088	-	851	228
	total	200	512	1616	3141	3891	-	1165	641
10. Maharashtra	male	438	698	2271	4612	7833	-	1835	449
	female	276	355	2552	3869	3343	-	1397	354
	total	361	506	2413	4221	5652	-	1609	803

Table (3R): contd. Number of persons suffering from chronic diseases per 100,000 persons by age separately by sex for different states and all-India, rural households NSS 28th round.

RURAL

states	age (years)								
	0-14	15-25	25-44	45-59	60 & above	not recorded	re-ages	all ages	number of sample cases
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	
11. Orissa	male	651	1236	3719	7498	11341	-	2949	387
	female	626	1280	3456	7561	7143	-	2616	348
	total	638	1259	3589	7530	9403	-	2781	735
12. Punjab	male	247	759	2459	6689	9717	-	2348	367
	female	269	900	3503	5772	6708	-	2197	305
	total	257	827	2970	6238	8458	-	2277	672
13. Rajasthan	male	246	540	1534	4912	5803	-	1425	196
	female	187	570	1940	2714	2728	-	1117	143
	total	218	554	1731	3770	4345	-	1276	339
14. Tamil Nadu	male	509	1088	2821	6486	8538	-	2453	367
	female	431	1090	3387	4323	3804	-	2008	306
	total	470	1089	3110	5382	6306	-	2226	673
15. Uttar Pradesh	male	263	485	1835	4498	6082	-	1585	177
	female	248	856	1963	3533	3338	-	1393	165
	total	256	661	1897	4009	4848	-	1493	342
16. West Bengal	male	1028	1260	4692	11763	18549	-	3920	934
	female	532	1173	5526	8224	848	-	2969	665
	total	786	1221	5088	9959	13743	-	3459	1599
17. All-India*	male	474	793	2703	6373	9248	13015	2260	7783
	female	361	917	2966	5109	5340	16625	1929	6480
	total	421	855	2833	5726	7404	14582	2098	14263

* including smaller states and Union Territories.

Table (8U) : Number of persons suffering from chronic diseases per 100,000 persons by age separately by sex for different states and all-India, urban households NSS 28th round.

URBAN		states	age (years)					number of sample cases
			0-14	15-24	25-44	45-59	60 & above	
(1)		(2)	(3)	(4)	(5)	(6)	(7)	(8)
1. Andhra Pradesh	male	268	299	1521	3177	4747	1100	146
	female	193	565	2259	2574	3225	1191	153
	total	231	426	1868	2864	3837	1143	299
2. Assam	male	232	670	2845	8379	9231	2213	106
	female	194	992	4222	419	5128	2063	76
	total	214	810	3372	7604	7464	2148	182
3. Bihar	male	267	1249	2679	4797	15726	2262	236
	female	270	605	3451	5385	7589	2046	179
	total	270	985	3022	5069	11583	2162	415
4. Gujarat	male	-	281	1050	1636	6552	797	64
	female	145	137	1245	1232	1728	616	44
	total	69	216	1141	1442	4080	712	108
5. Haryana	male	461	746	3649	7806	12037	2713	135
	female	277	905	3768	5870	8929	2228	97
	total	373	818	3704	6852	10796	2487	232
6. Jammu & Kashmir	male	116	226	856	2605	5465	792	182
	female	94	390	1363	2576	3942	842	168
	total	106	302	1084	2591	4822	816	350
7. Karnataka	male	444	375	1490	3726	5975	1290	111
	female	272	280	1393	3012	4388	1071	91
	total	359	328	1442	3356	5137	1181	202
8. Kerala	male	3684	2256	10721	22137	36613	8865	534
	female	2643	2253	11976	24232	25537	8884	527
	total	3186	2256	11360	23175	30393	8874	1061
9. Madhya Pradesh	male	179	311	1301	5185	8686	1429	138
	female	135	366	2514	3514	3426	1270	112
	total	157	336	1868	4387	5980	1350	250
10. Maharashtra	male	289	655	1744	5465	6241	1570	305
	female	360	741	1949	4435	6544	1610	265
	total	321	692	1831	5000	6399	1591	570

Table (8U): contd. Number of persons suffering from chronic diseases per 100,000 persons by age separately by sex for different states and all-India, urban households : NCS 28th round.

URBAN

states		age (years)						number of sample cases
		0-14	15-24	25-44	45-59	60 & above	all ages	
(1)		(2)	(3)	(4)	(5)	(6)	(7)	(8)
11. Orissa	male	619	1012	2732	6720	8297	2180	97
	female	260	732	4493	6170	6719	2459	93
	total	452	891	3510	6439	7469	2312	190
12. Punjab	male	473	416	1724	4097	8036	1621	87
	female	334	607	2342	7042	7326	1993	93
	total	405	502	1998	5513	7718	1794	180
13. Rajasthan	male	191	280	1343	2256	3243	834	55
	female	313	524	2263	1424	994	937	57
	total	251	388	1778	1839	2061	884	112
14. Tamil Nadu	male	941	489	2080	5412	8637	2049	296
	female	349	1040	3086	4059	6004	1927	268
	total	647	764	2551	4737	7252	1990	564
15. Uttar Pradesh	male	222	637	1363	3361	4592	1182	77
	female	158	683	2156	2619	4357	1251	69
	total	192	657	1718	3017	4486	1213	146
16. West Bengal	male	1596	1217	3565	8910	18640	3924	529
	female	787	1367	5651	10446	12016	3902	414
	total	1207	1284	4379	9531	15316	3914	943
17. All India*	male	591	656	2232	5662	9450	2003	3373
	female	379	763	2982	4954	6388	1931	2912
	total	488	704	2571	5321	7893	1969	6285

* including smaller states and Union Territories.