

*R. 20/3*

DRAFT

THE NATIONAL SAMPLE SURVEY  
TWENTYEIGHTH ROUND  
OCTOBER 1973 - JUNE 1974

NUMBER 292

TABLES WITH NOTES ON  
MORBIDITY

ISSUED BY

NATIONAL SAMPLE SURVEY ORGANISATION  
DEPARTMENT OF STATISTICS : MINISTRY OF PLANNING  
GOVERNMENT OF INDIA

SEPTEMBER 1980

TABLES WITH NOTES ON  
MORBIDITY

	<u>CONTENTS</u>	<u>Page</u>
CHAPTER ONE	: Introduction	1
CHAPTER TWO	: Concepts and definition	2
CHAPTER THREE	: Summary findings	3
APPENDIX I	: Sampling Design and Estimation Procedure	i
Appendix II	: Statistical Tables : List of tables Tables -	iii  1

NATIONAL SAMPLE SURVEY  
28TH ROUND : OCTOBER 1973 - JUNE 1974  
TABLES WITH NOTES ON MORBIDITY  
CHAPTER ONE

INTRODUCTION

1.1. In the twenty eight round of the National Sample Survey (NSS) information ~~experience~~ on morbidity was collected along with the enquiry on births, deaths, migration and disability<sup>a/</sup>. Information on morbidity was collected with reference to last 2 weeks preceding the date of survey from all members present in the sample households on the date of enquiry as well as visitors in the households. Morbidity was defined as sickness/injury/poisoning.

Information on sickness/injury/poisoning affecting members of the household including visitors during the 2 weeks preceding the date of enquiry together with ancillary information like age and sex of the person, nature of the ailment etc. was collected. Also along with information on morbidity data on chronic ailments prevailing among the members on the date of survey were collected.

1.2. The survey covered the whole of India excepting Ladakh district of Jammu & Kashmir, Mizoram, Arunachal Pradesh, Andaman & Nicobar Islands, Lakshadweep, Minicoy and Amindivi Islands, Dadra and Nagar Haveli, rural areas of Chandigarh and Nagaland, some rural areas in Tripura and in the districts of Surguja and Bastar of Madhya Pradesh and <sup>Chandrapur</sup> ~~Gondia~~ and Amravati of Maharashtra.

1.3. The survey started in October 1973 and continued upto June 1974 and the survey period of nine months was split up into three sub-rounds of three months duration each. Data were collected from the sample households by the interview method. Broadly, the sample design was a stratified two stage one with village/urban blocks being first stage units and households as the second stage units and self-weighting at the state level separately for the rural and the urban sectors. In all, from about 1.29 lakhs of rural and 0.67 lakh urban households the relevant data on morbidity were collected. The present tables on morbidity are based on such data. The sample design and procedure of estimation are described in Appendix I.

---

a/ : Information on morbidity was also collected in seventeenth round of NSS from about 8,600 rural households and 4,400 urban households distributed over 4,310 villages & 2,237 urban blocks.

CHAPTER TWO  
CONCEPTS AND DEFINITIONS

2.1. Household normally resident members and visitors : Persons who usually lived together and took food from the same kitchen were considered as usual or normally resident members of the household. Other members not usually residing in the household were classified as visitors. If a person lived in a place and took for at some other place, he was considered usual member at the place where he lived. Inmates (including the staff) of a mess, hotel, orphanage, etc. were treated as single member households. However, a family living in anyone of the establishments was treated as a separate household by itself.

2.2. Morbidity was defined as sickness which included injury and poisoning also. Sickness meant any deviation from the state of physical and mental well being with a specific cause. Chronic ailments ~~and the associated~~ xxxxxxxx were not, however, treated as morbidity.

2.3. The following categories of spells of morbidity could occur during a reference period.

- (a) Spells starting before the first day of the reference period and terminating within the reference period.
- (b) Spells starting and terminating within the reference period.
- (c) Spells starting within the reference period and continuing on the date of survey.
- (d) Spells starting before the first day of the reference period and continuing on the date of survey.

2.4. Incidence rate is defined by

$$\frac{\text{estimated number of spells } (b + c)}{\text{estimated population exposed to risk}} \times 1000$$

$$= 1000 \times \frac{\text{estimated number of spells starting within the reference period}}{\text{estimated population exposed to risk}}$$

The incidence rate measures the frequency of morbidity commencing during the reference period

2.5. Prevalence rate is defined by

$$\frac{\text{estimated number of spells } (a+b+c+d)}{\text{estimated population exposed to risk}} \times 1000$$

$$= 1000 \times \frac{\text{estimated number of spells prevailing in the reference period}}{\text{estimated population exposed to risk}}$$

The prevalence rate measures the frequency of morbidity prevailing in the reference period.

CHAPTER THREE  
SUMMARY FINDINGS

3.1. Table 1 below presents incidence and prevalence rates of morbidity for all-India separately by states. From the table it is seen that at all-India level the incidence rate of morbidity per one thousand persons was 12.57 and 13.53 respectively for the rural and the urban sectors.<sup>b/</sup>

Table 1 : Incidence and prevalence rate of morbidity by states and all-India, rural & urban areas.

states	incidence rate		prevalence rate	
	rural	urban	rural	urban
(1)	(2)	(3)	(4)	(5)
1. Andhra Pradesh	20.89	18.20	31.94	27.84
2. Assam	8.46	5.94	17.40	11.66
3. Bihar	4.69	6.05	10.16	11.42
4. Gujarat	4.48	2.85	10.17	6.26
5. Haryana	9.23	8.46	15.96	16.47
6. Jammu & Kashmir	11.42	9.22	17.79	12.80
7. Karnataka	8.44	7.24	14.92	13.35
8. Kerala	37.29	33.77	71.21	61.84
9. Madhya Pradesh	10.34	11.34	20.89	21.72
10. Maharashtra	17.78	20.06	27.57	32.18
11. Orissa	14.53	17.19	24.20	24.38
12. Punjab	19.25	16.68	27.47	24.62
13. Rajasthan	10.20	7.46	17.33	12.55
14. Tamil Nadu	20.65	21.01	33.99	31.98
15. Uttar Pradesh	7.95	6.27	13.22	10.54
16. West Bengal	11.35	17.30	27.71	33.85
17. All-India <sup>a/</sup>	12.57	13.53	22.46	22.77

<sup>a/</sup> All states and union territories under NSS coverage<sup>b/</sup>

It also appears from the table that there were wide variation in the incidence rate between the states. Kerala with an incidence rate of 37.29 and 33.77 respectively in the rural and the urban sectors tops the list. This was followed by Tamil Nadu with a rate of 20.65 and 21.01 in the rural and urban sectors. The states Andhra, Maharashtra, Bihar, Orissa, Punjab and Tamil Nadu had incidence rates both in the rural and the urban sectors higher than the all-India rate. With regard to prevalence rate it is seen from the table that at all-India the prevalence rate of temporary ailments was 22.46 and 22.77 respectively in the rural and the urban sectors. Kerala here again tops the list with a rate of 71.21 in the rural and 61.84 in the urban sectors. This was followed by Tamil Nadu with a rate of 33.99 and 31.98. The states Andhra, Maharashtra, Orissa and Punjab had rates higher than the all-India rates.

<sup>b/</sup> The incidence and prevalence rates of morbidity of NSS 28th round as estimated are somewhat lower than the rates as observed in the 17th round of the NSS presented in NSS No.129 : Report of Pilot Enquiry on Morbidity. A seminar meeting was held in July 1977 under the auspices of NSSO to examine vital rates and migration and morbidity rates of NSS 28th round and the consensus was that the morbidity rates could be released even though the estimates of NSS 28th round appeared somewhat lower than those of the Pilot Enquiry on Morbidity.

3.2. Table 2 presents incidence and prevalence rates of morbidity by type of ailment for all-India. From the table it is seen that excepting the group 'others' and 'not recorded' the incidence rate of influenza was highest both in the rural & urban areas. It was 2.2 per 1000 persons in both the sectors. In the rural sector influenza was followed by malaria with a rate of 1.1. The incidence rate of malaria in the urban sector was not so high. The prevalence rate of influenza also exceeded the corresponding rates of other ailments in both the sectors. Malaria and dysentery also had a comparatively high rate. The prevalence rates estimated for influenza, malaria and dysentery were 3.1, 1.8 and 1.3 in rural areas and the corresponding rates for the urban sector being 3.2, 1.2 and 1.3.

Table 2 : Incidence and prevalence rate of morbidity by type of temporary ailments for all-India, rural and urban areas

type of ailments (1)	incidence rate		prevalence rate	
	rural (2)	urban (3)	rural (4)	urban (5)
1. cholera	0.03	0.03	0.05	0.04
2. typhoid	0.12	0.19	0.35	0.44
3. dysentery	0.74	0.79	1.34	1.29
4. diarrhoea	0.27	0.22	0.46	0.33
5. diphtheria	0.02	0.01	0.03	0.01
6. whooping cough	0.33	0.25	0.84	0.60
7. tetanus	0.01	0.01	0.02	0.03
8. acute poliomyelitis	0.01	0.02	0.03	0.05
9. small pox	0.38	0.44	0.63	0.79
10. measles	0.17	0.14	0.31	0.30
11. mumps	0.03	0.08	0.05	0.12
12. malaria	1.13	0.71	1.77	1.15
13. influenza	2.16	2.15	3.10	3.24
14. pneumonia	0.13	0.05	0.20	0.11
15. food poisoning	0.02	0.04	0.03	0.04
16. accident	0.39	0.54	0.80	1.01
17. others	4.34	5.16	8.27	8.51
18. not recorded	2.29	2.70	4.18	4.71
19. all ailments	12.57	13.53	22.46	22.77

3.3. Incidence and prevalence rates of morbidity broken down by age is shown in table 3. Both incidence and prevalence rates of sickness as revealed in table 3 was highest in the first year of age and gradually diminished as age increased. It was minimum in the age group 15-24 and then again rose higher with the age. The pattern was observed both in the rural and urban areas of all-India. The incidence rates per 1000 person at age '0' were estimated as 21.09 for rural and 30.29 for urban sectors. The corresponding prevalence rates for rural and urban sectors were 35.72 and 49.34 respectively.

Table 3 : Incidence and prevalence rate of temporary ailments by age for all-India, rural and urban

age in years	incidence rate		prevalence rate	
	rural	urban	rural	urban
(1)	(2)	(3)	(4)	(5)
0	21.09	30.29	35.72	49.34
1-4	20.65	24.36	35.50	39.04
5-14	10.04	11.02	17.15	18.14
15-24	9.53	9.30	16.38	14.97
25-44	12.04	13.64	22.10	23.13
45-59	13.64	13.41	25.51	25.24
60 & above	13.67	12.97	28.78	25.57
not recorded	103.98	-	180.54	-
all ages	12.57	13.53	22.46	22.77

3.4. Average duration of selected temporary ailments which terminated in recovery during the reference period is presented in table 3 by type of ailment. It is seen from the table that at all-India level average duration of all temporary ailments taken together was about 8.2 and 6.7 days in the rural and the urban sectors respectively. There was, however, some variation by type of ailments. In the rural sector, the duration of typhoid was 18.8 days while the duration of measles, mumps, small pox & malaria was about 8-10 days. In the urban sector similarly the duration of typhoid was about 16 days while the duration of small pox, mumps, measles & malaria was about 8-10 days.

Table 4 : Average duration (days) of selected temporary ailments terminating in recovery by type of ailments for all-India, rural and urban areas

type of ailments	duration of temporary ailments	
	rural	urban
(1)	(2)	(3)
1. cholera	5.70	5.32
2. typhoid	18.84	16.43
3. diphtheria	5.15	7.00
4. whooping cough	10.52	6.92
5. small pox	10.33	10.13
6. measles	9.61	8.31
7. mumps	9.02	8.27
8. malaria	8.36	6.54
9. influenza	6.66	5.76
10. pneumonia	7.94	7.46
11. food poisoning	12.95	11.14
12. accident	10.73	7.20
13. all ailments	8.16	6.68

3.5. Information on chronic ailments prevailing among the members of the household are presented in table 5. It is seen that the incidence rate of all chronic ailments taken together was 2098 and 1962 per 1,00,000 persons in the rural and urban sectors. It is also seen that asthma was the most prevalent chronic ailment both in the rural and urban areas. High blood pressure with a rate of 132 in the urban sector compared to 44 in the rural sector appears to be more urban specific. The number of persons suffering from this ailment came to 376 in the rural areas and 355 in the urban areas per 1,00,000 persons. The next most important chronic diseases were rheumatism and tuberculosis in both the rural and urban areas, the number of cases being respectively 251 and 117 in rural whereas 146 and 137

Table 5 : Number of persons suffering from chronic diseases per 1,00,000 persons by type of chronic disease for all-India, rural & urban areas

type of chronic disease (1)	number of persons	
	rural (2)	urban (3)
1. tuberculosis	117	137
2. leprosy	40	25
3. syphilis	8	6
4. cancer	12	14
5. thyroid trouble or goitre	22	16
6. diabetes	39	80
7. mental illness	19	18
8. epilepsy	28	17
9. rheumatic fever	45	26
10. high blood pressure	44	132
11. bronchitis	41	43
12. asthma	376	355
13. peptic ulcer	89	66
14. kidney stone or kidney trouble	37	40
15. arthritis	22	17
16. rheumatism	251	146
17. stroke	13	13
18. piles	65	61
19. others	194	128
20. not recorded	636	629
21. all types	2098	1962

in urban areas. High blood pressure with a rate of 132 in the urban sector compared to 44 in the rural sector appears to be more urban specific.

APPENDIX - I

SAMPLING DESIGN AND ESTIMATION PROCEDURE

A stratified two stage design was adopted for the survey. The first stage units were villages in the rural sector and blocks in the urban sector and the second stage units were households from which the information was collected.

The whole of India was divided into 524 basic strata so that the basic strata did not cut across district boundaries. A district with more than 1.5 million 1971 census rural population formed 2 or more basic strata (by grouping contiguous tehsils) depending on their rural population. In Gujarat, however districts with population less than 1.5 million were divided into two strata in consideration of dry areas and concentration of tribal population at the request of SSB of that state.

Within each State and Union Territory the allotted number of villages and blocks were distributed over the constituent basic strata in proportion to their rural and urban population respectively.

The 1971 census list of villages was used as the sampling frame for selection of villages. Sample villages within each basic stratum were selected with probability proportional to population and with replacement in the form of two independent sub-samples.

All cities and towns within each basic stratum were grouped into a number of urban sub-strata. All towns with 1971 census population less than 20,000 formed sub-stratum 1, towns with 1971 population 20,000 to 49,999 formed sub-stratum 2 and those with 1971 population 50,000 to 99,999 formed sub-stratum 3. Each city with 1971 population 1,00,000 or more constituted a separate urban sub-stratum. Within each urban stratum, the allocated number of blocks was further allocated to the constituent sub-strata circular systematically with probability proportional to urban populations of the respective sub-strata. Within each urban

sub-stratum, the allocated number of blocks were selected with probability proportional to size and with replacement.

Villages and blocks with large population content were divided into two or more compact areal sub-divisions of nearly equal population content. One of these sub-divisions (called hamlet in a village and sub-block in a block) was selected at random and the survey was confined to the selected sub-division only.

All households in a sample village/hamlet group were arranged into 6 classes on the basis of their means of livelihood viz., (i) landless labour, (ii) cultivator and (iii) others and whether (i) visited health centre or (ii) not. All households in a sample block/sub-block were also arranged into 6 classes by occupation divisions (0-1 and 2,3,4,5,6 and 7-8-9 and X). From the arranged frame in a village/block, a sample of households was selected linear systematically using the intervals and random start specified. Sampling intervals were so fixed as to make the design selfweighting at the state level separately for rural and urban sectors.

Unbiased method of estimation was adopted for estimating aggregate values of the characteristics, say, X, Y, Z, etc. under enquiry. Estimates of ratios were derived by dividing unbiased estimates of the numerators by the unbiased estimates of the denominators.

Appendix - II  
Statistical Tables

A list of tables included in this  
is given below :

	page no.
Table (1R) : Incidence rate of temporary ailments by type of ailment separately by sex for different states and all-India : rural households, NSS 28th round.	1
Table (iv) : Incidence rate of temporary ailments by type of ailment separately by sex for different states and all-India : Urban households, NSS 28th round.	4
Table (2R) : Incidence rate of temporary ailments by age separately by sex for different states and all-India : rural households, NSS 28th round.	7
Table (2U) : Incidence rate of temporary ailments by age separately by sex for different states and all-India : urban households, NSS 28th round.	9
Table (3R) : Prevalence rate of temporary ailments by type of ailments separately by sex for different states and all-India : rural households, NSS 28th round	11
Table (3U) : Prevalance rate of temporary ailments by type of ailments separately by sex for different states and all-India : Urban households, NSS 28th round.	14
Table (4R) : Prevalence rate of temporary ailments by age separately by sex for different states and all-India : rural households, NSS round	17
Table (4U) : Prevalence rate of temporary ailments by age separately by sex for different states and all-India : urban households, NSS 28th round.	19
Table (5R) : Average duration (days) of temporary ailments terminating in recovery by type of ailment separately by sex for different states and all-India : rural households, NSS 28th round.	21

page no.

- Table (5U) : Average duration (days) of temporary ailments terminating in recovery by type of ailment separately by sex for different states and all-India : Urban households, NSS 28th round. 24
- Table (6R) : Average duration (days) of temporary ailments terminating in recovery by age separately by sex for different states and all-India : rural households, NSS 28th round. 27
- Table (6U) : Average duration (days) of temporary ailments terminating in recovery by age separately by sex for different states and all-India : urban households, NSS 28th round. 29
- Table (7R) : Number of persons suffering from chronic diseases per 100,000 persons by type of chronic disease separately by sex for different states and all-India : rural households, NSS 28th round. 31
- Table (7U) : Number of persons suffering from chronic disease per 100,000 persons by type of chronic disease separately by sex for different states and all-India : urban households, NSS 28th round. 34
- Table (8R) : Number of persons suffering from chronic diseases per 100,000 persons by age separately by sex for different states and all-India : rural households, NSS 28th round. 37
- Table (8U) : Number of persons suffering from chronic diseases per 100,000 persons by age separately by sex for different states and all-India : urban households, NSS 28th round. 39

Table (1R) : Incidence rate of temporary ailments by type of ailments separately by sex for different states and all-India, rural households : NSS 28th round

states	sex	type of ailments																		all types of ailments	number of sample
		cholera	typhoid	dysentery (all forms)	diarrhoea	diphtheria	whooping cough	tetanus	acute poliomy- elitis	small pox	measles	mumps	malaria	influenza	pneumonia	food poisoning	accident	others			
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)
Andhra Pradesh	M	-	0.13	0.58	0.14	-	0.31	0.05	-	0.70	0.13	-	1.95	4.68	0.18	-	0.35	7.28	5.44	21,92	492
	F	-	0.09	0.40	0.05	-	0.23	-	-	0.63	0.18	0.05	2.35	3.95	0.09	0.05	0.09	5.38	6.30	19.84	436
	T	-	0.11	0.49	0.09	-	0.27	0.02	-	0.66	0.16	0.02	2.14	4.32	0.14	0.02	0.22	6.36	5.87	20.89	928
Assam	M	-	0.30	2.34	0.14	0.36	0.29	-	-	0.29	-	-	0.96	3.04	0.07	-	0.14	0.43	1.07	9.43	130
	F	-	0.08	0.97	0.16	0.16	0.16	-	-	0.16	-	-	0.42	3.71	0.08	-	0.40	0.48	0.56	7.34	91
	T	-	0.19	1.69	0.15	0.27	0.23	-	-	0.23	-	-	0.70	3.35	0.08	-	0.27	0.46	0.84	8.46	221
Bihar	M	0.04	0.04	0.43	0.21	-	0.18	-	-	0.39	0.04	-	0.14	0.96	0.14	-	0.18	2.62	0.11	5.48	154
	F	0.04	0.04	0.15	0.11	-	0.11	0.04	-	0.18	-	-	0.07	0.77	0.04	-	0.04	2.20	0.11	3.90	106
	T	0.04	0.04	0.29	0.16	-	0.14	0.02	-	0.29	0.02	-	0.11	0.86	0.09	-	0.11	2.41	0.11	4.69	260
Gujarat	M	0.08	-	0.16	0.16	-	0.65	-	-	0.16	-	0.08	0.89	0.72	-	-	0.40	0.32	0.56	4.18	52
	F	0.08	0.17	-	0.09	-	0.08	-	-	0.26	-	-	0.94	1.37	0.08	-	0.42	0.61	0.68	4.79	56
	T	0.08	0.08	0.08	0.13	-	0.37	-	-	0.21	-	0.04	0.92	1.04	0.04	-	0.42	0.45	0.62	4.48	108
Haryana	M	0.12	0.12	0.18	0.18	-	0.43	-	-	0.18	0.37	-	1.29	0.61	0.37	0.06	0.31	1.41	4.42	10.05	164
	F	0.14	0.34	0.34	0.07	0.07	0.28	-	-	0.21	0.28	-	0.76	0.28	0.07	-	-	0.76	4.75	8.35	121
	T	0.13	0.23	0.26	0.13	0.03	0.36	-	-	0.19	0.32	-	1.04	0.45	0.23	0.03	0.16	1.10	4.57	9.23	285
Jammu & Kashmir	M	-	0.35	1.29	0.70	0.03	0.38	-	-	0.03	0.07	-	0.53	4.29	0.21	0.03	0.14	2.11	0.73	10.89	318
	F	0.04	0.35	1.48	0.89	-	0.54	-	-	0.04	0.16	0.04	0.40	5.22	0.20	0.04	0.04	1.76	0.78	11.98	308
	T	0.02	0.35	1.38	0.79	0.02	0.46	-	-	0.04	0.11	0.02	0.47	4.73	0.20	0.04	0.09	1.94	0.76	11.42	621

+ M = male, F = female, T = total

Table (1R) Contd. : Incident rate of temporary ailments by type of ailments separately by sex for different states and all-India, rural households : NSS 28th round

states	+ sex	type of ailments																			all type of ailments	number of sample ailments
		cholera	typhoid	dysentery (all forms)	diarrhoea	diphtheria	whooping cough	tetanus	acute poliomy- elitis	small pox	measles	mumps	malaria	influenza	pneumonia	food poisoning	accident	others	not recorded			
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	
Karnataka	M	-	-	1.06	-	-	0.64	-	0.07	0.14	-	-	0.14	0.71	-	0.07	0.07	2.06	3.69	8.65	122	
	F	-	-	0.51	-	-	0.51	-	-	0.29	0.07	0.07	-	0.65	-	-	0.22	2.03	3.85	8.20	113	
	T	-	-	0.79	-	-	0.57	-	0.04	0.22	0.04	0.04	0.07	0.68	-	0.04	0.14	2.04	3.77	8.44	235	
Kerala	M	-	-	2.58	0.84	0.07	0.35	-	-	0.28	0.63	-	-	5.24	-	-	1.82	15.92	11.52	39.25	562	
	F	-	0.07	2.26	1.40	-	0.53	-	-	0.13	1.00	-	-	5.58	0.07	-	0.40	13.95	10.10	35.49	534	
	T	-	0.03	2.42	1.12	0.03	0.44	-	-	0.20	0.82	-	-	5.41	0.03	-	1.09	14.91	10.79	37.29	1096	
Madhya Pradesh	M	-	0.07	0.11	0.22	0.07	0.36	-	0.04	0.43	0.25	0.11	3.09	0.25	0.22	0.07	0.22	5.53	-	11.04	307	
	F	-	0.04	0.26	0.15	0.11	0.26	-	-	0.42	0.04	0.08	2.99	0.30	0.15	0.19	0.04	4.62	-	9.65	255	
	T	-	0.06	0.18	0.18	0.09	0.31	-	0.02	0.42	0.15	0.09	3.04	0.28	0.18	0.13	0.13	5.08	-	10.34	562	
Maharashtra	M	-	0.17	1.08	0.12	-	0.50	0.04	-	0.25	0.33	0.08	1.38	5.43	0.12	-	0.63	4.63	5.20	19.96	479	
	F	0.04	0.15	0.49	0.15	-	0.49	-	-	0.19	0.41	-	1.24	4.26	0.07	-	0.41	2.67	4.13	14.70	391	
	T	0.02	0.16	0.80	0.14	-	0.51	0.02	-	0.22	0.39	0.04	1.35	4.98	0.10	-	0.53	3.72	4.80	17.78	870	
Orissa	M	-	0.08	1.23	-	0.08	0.46	-	-	1.08	0.54	-	1.00	2.69	0.15	-	0.38	5.38	4.62	17.69	230	
	F	-	0.08	0.83	0.45	-	0.15	-	-	0.15	0.15	0.08	1.35	1.80	-	-	0.22	3.62	2.56	11.44	152	
	T	-	0.08	1.03	0.23	0.04	0.30	-	-	0.61	0.34	0.04	1.18	2.24	0.08	-	0.30	4.49	3.57	14.53	382	
Punjab	M	0.66	0.46	1.84	0.86	-	1.12	-	0.13	0.13	0.33	0.07	6.92	1.71	1.12	-	0.92	-	4.41	22.46	341	
	F	0.15	0.51	0.51	0.07	-	0.81	-	-	0.22	0.81	0.07	5.36	1.61	0.59	-	-	-	3.89	15.63	213	
	T	0.42	0.49	1.21	0.49	-	0.97	-	0.07	0.17	0.56	0.07	6.18	1.67	0.87	-	0.49	-	4.17	19.25	554	

+ M = male, F = female, T = total

Table (1a)Contd. : Incidence rate of temporary ailments separately by sex for different states  
and all-India, rural households : NSS 28th round

states	sex	type of ailments										not recorded	all types of ailments	number of respondents							
		cholera	typoid	diarrhoea	whooping cough	cough	teethitus	acute poliomyelitis	small pox	measles	mumps										
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)
Rajasthan	M	-	0.37	0.15	0.15	-	0.37	0.08	-	0.60	-	-	3.21	0.68	0.37	-	0.76	2.69	1.65	11.08	148
	F	-	0.16	0.48	0.08	-	0.08	-	-	0.32	-	-	3.26	0.89	0.32	-	0.24	1.35	2.08	9.26	116
	T	-	0.27	0.31	0.12	-	0.23	0.04	-	0.46	-	-	3.23	0.78	0.35	-	0.51	2.04	1.86	10.20	264
Tamil Nadu	M	-	0.20	1.55	0.20	-	0.54	-	-	0.40	0.14	-	0.14	1.55	-	-	0.74	10.88	3.84	20.18	299
	F	-	0.06	1.70	0.26	-	0.39	-	-	0.79	0.13	-	0.72	-	-	-	0.26	1.66	5.13	21.10	321
	T	-	0.13	1.63	0.23	-	0.46	-	-	0.60	0.14	-	0.07	1.13	-	-	0.50	1.27	4.49	20.65	620
Uttar Pradesh	M	-	0.03	0.35	0.35	0.03	0.45	-	-	0.61	0.19	-	0.77	1.17	0.29	-	0.69	4.34	-	9.27	348
	F	0.06	0.18	0.24	0.15	0.03	0.18	-	0.03	0.39	-	0.03	0.75	0.99	0.09	-	0.27	2.94	-	6.33	211
	T	0.03	0.10	0.30	0.25	0.03	0.32	-	0.01	0.51	0.10	0.01	0.76	1.16	0.20	-	0.49	3.68	-	7.95	559
West Bengal	M	-	0.26	1.36	0.51	-	0.17	0.08	0.04	0.26	0.09	0.04	0.09	4.44	0.04	-	1.07	3.07	0.81	12.37	290
	F	0.09	0.27	1.26	0.63	-	0.04	-	-	0.27	0.13	0.05	0.09	4.05	-	-	0.27	2.48	0.85	10.26	228
	T	0.04	0.26	1.32	0.57	-	0.11	0.04	0.02	0.26	0.11	0.04	0.09	4.25	0.02	-	0.68	2.79	0.83	11.35	518
All-India	M	0.03	0.12	0.84	0.28	0.03	0.40	0.02	0.01	0.43	0.18	0.02	1.16	2.25	0.18	0.01	0.56	4.76	2.25	13.53	4675
	F	0.07	0.11	0.64	0.26	0.02	0.26	0.00	0.00	0.33	0.15	0.03	1.09	2.06	0.09	0.02	0.22	3.90	2.34	11.55	3937
	T	0.03	0.12	0.74	0.27	0.02	0.33	0.01	0.01	0.38	0.17	0.03	1.13	2.16	0.13	0.02	0.39	4.34	2.29	12.57	8612

\* including smaller states and union territories.

\* M = male, F = female, T = total

Table (1U) : Incidence rate of temporary ailments by type of ailiments separately by sex for different states and all-India, urban households : NSS 28th round

states	+ sex	cholesterol	diabetes	dysentery (all forms)	typhoid	tuberculosis	acute polyarthritis	acute poliomyelitis	small pox	measles	influenza	pneumonia	food poisoning	accident	others	not recorded	all types of ailments	sample number	all types of ailments			
Andhra Pradesh	M	0.23	0.54	6.09	0.08	0.46	0.15	1.76	2.38	1.23	5.37	5.52	17.80	232								
	F	0.08	0.79	0.48	0.08	1.27	0.08	0.16	1.03	3.01	1.19	5.69	4.60	18.62	235							
	T	0.16	0.66	0.27	0.08	0.86	0.12	0.09	1.40	2.69	1.21	3.53	5.06	18.20	467							
Assam	M	0.21	0.88	0.42							2.59			0.63	0.42	5.15	24					
	F		1.40							0.82	0.12	0.56	1.66		2.24	0.55	7.23	26				
	T	0.12	1.11	0.24						0.35	0.05	0.25	2.18		1.34	0.35	5.94	50				
Bihar	M	0.19	0.19	0.19						0.29		0.29	1.65	0.10	0.19	3.20	0.48	6.77	70			
	F	0.12	0.23	0.12						0.23		0.12	1.16	0.12	0.12	2.66	0.23	5.23	45			
	T	0.16	0.21	0.05						0.26		0.21	1.42	0.05	0.16	2.95	0.37	6.05	115			
Gujarat	M								0.08	0.08		0.40	0.40	0.32	0.64	0.55	2.47	31				
	F	0.42							0.05	0.10	0.05	0.14	0.86	0.42	0.70	0.70	3.53	25				
	T	0.15							0.05		0.05	0.05	0.56	0.44	0.20	0.67	2.85	56				
Haryana	M	0.21		0.62	0.21	0.41		0.21			1.23	0.21	1.85	3.90	2.67	1.73	57					
	F									0.22		0.24	0.24	0.24	3.30	0.71	4.73	20				
	T	0.11		0.33	0.11					0.11		0.11	0.11	1.10	3.62	1.76	8.46	77				
Jammu & Kashmir	M	0.09	0.27	0.53		0.09		0.09				5.59	0.18	0.09	1.42	0.97	9.32	106				
	F	0.60	0.51	0.52						0.10	6.51	0.10	0.61	0.51	0.76	9.04	90					
	T	0.05	0.42	0.52						0.05	0.05	0.10	0.05	1.05	0.76	9.22	196					

+ M = male, F = female, T = total

Table (1V) Contd : Incidence rate of temporary ailments by type of ailments separately by sex for different states and all-India, urban households, NSS 28th round

		type of ailments																			
		states						all types of ailments													
		Cholera	Typhoid	Dysentery	Diarrhoea	Gastroenteritis	Tetanus	Acute poliomyelitis	Small pox	Influenza	Pneumonia	Food poisoning	Accidental	Others	Total recorded	All types of ailments	Number of sample				
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)
Rajasthan	M	0.15	0.62	0.31		0.15		0.15	0.75	0.61	0.31	0.31	4.73	0.30	8.39	55					
	F	0.33				0.16		0.17	2.51	0.32		0.33	3.13	0.50	6.45	39					
	T	0.23	0.33	0.16		0.16		0.08	0.08	1.11	0.47	0.16	0.32	3.96	0.40	7.46	94				
Uttar Pradesh	M	1.13	0.49			0.50		1.19	0.21		1.92		0.35	10.11	4.97	21.08	298				
	F	1.25	0.22			0.30		0.73	0.14		1.76		0.22	12.26	3.98	20.93	286				
	T	1.18	0.36			0.40		0.96	0.18	0.11	1.84		0.29	11.17	4.48	21.01	584				
West Bengal	M	0.11	0.17	0.06		0.39		0.50	0.11		0.33	0.77	0.06	0.72	3.26		6.54	118			
	F	0.13	0.27			0.20		0.66	0.07		0.20	1.40	0.07	0.20	2.79		5.99	90			
	T	0.03	0.12	0.03		0.30		0.57	0.09		0.27	1.06	0.06	0.48	3.05		6.27	208			
* All-India	M	0.08	0.30	1.80	0.30		0.07		0.23	0.30	0.30	4.37		1.43	6.26	2.71	18.60	247			
	F	0.10	0.10	1.43	0.28		0.10		0.38	0.38	0.10	4.48		0.66	4.88	2.57	15.65	164			
	T	0.09	0.21	1.64	0.30		0.08		0.30	0.33	0.21	4.42		1.10	5.63	2.65	17.30	411			
M	M	0.03	0.17	0.78	0.23	0.01	0.29	0.02	0.39	0.15	0.09	0.73	2.10	0.06	0.70	5.33	2.74	13.87	2306		
	F	0.03	0.21	0.81	0.20		0.21	0.02	0.49	0.13	0.06	0.69	2.22	0.05	0.34	4.97	2.67	13.14	1855		
	T	0.03	0.19	0.79	0.22	0.01	0.25	0.01	0.44	0.14	0.08	0.71	2.15	0.05	0.04	0.54	5.16	2.70	13.53	4161	
		M																			
		F																			
		T																			

\* including smaller states and union territories

+ M = male, F = female, T = total

Table ( 1U ) Contd : Incidence rate of temporary ailments by type of ailments separately by sex for different states  
and-India, urban households : NSS 28th round

states	t y p e o f a i l m e n t s																					
	+ X C S	cholesterol	typoiodia	dysentery	diarrhoea	diptheriae	whooping cough	cough	tetanus	acute poliomyelitis	small pox	measles	influenza	meningitis	food poisoning	accident	others	not recorded	number of II types of illnesses	number of simple types of illnesses		
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	
M	0.12	0.72	0.12	0.72	0.12	0.72	0.12	0.49	0.12	0.36	1.20	0.12	0.24	1.68	2.87	8.76	73					
Karnataka	F			0.48	0.24		0.12		0.36	0.12	1.21			1.09	2.07	5.69	47					
T		0.06	0.60	0.12	0.06	0.42			0.24	0.24	1.21	0.06		0.12	1.39	2.48	7.24	120				
M		0.84	0.66		0.17		0.36	0.17	0.66		6.10	0.18		2.89	11.29	8.96	32.28	190				
Kerala	F		3.43	0.67		0.34		0.18	1.01	0.50		6.70		0.18	11.41	10.80	35.22	205				
T			2.13	0.67		0.26		0.27	0.59	0.59		6.40	0.09		1.54	13.35	9.88	33.77	395			
M	0.11	0.21	0.53	0.21		0.21		0.53	0.11		2.54	0.21		0.53	6.89		12.72	120				
Madhya Pradesh	F	0.23	0.70	0.35	0.35		0.47	0.12	0.23	0.58		1.17	0.58	0.12	0.12	4.78		9.80	84			
T	0.17	0.44	0.44	0.28		0.33	0.06	0.11	0.56	0.06		1.89	0.61	0.17	0.33	5.89		11.34	204			
M	0.26	1.16	0.47		0.31		0.05	0.21	0.05	0.05	0.95	3.74			0.47	7.99	5.99	21.70	413			
Mahara- shtra	F	0.25	0.92	0.37		0.12	0.06	0.19	0.06	0.06	0.74	3.72			0.49	5.37	5.75	18.10	293			
T		0.26	1.05	0.43		0.23	0.03	0.03	0.20	0.06	0.06	0.85	3.73			0.48	6.78	5.87	20.06	706		
M	0.22	2.93	0.45		0.45				1.13			0.45	2.48			0.90	6.08	4.51	19.60	87		
Orissa	F		1.59	2.32	0.24				1.59			0.53	1.86			0.53	2.92	5.32	14.34	54		
T		0.12			0.24				1.34			0.49	2.19			0.73	4.63	4.89	17.19	141		
M	0.19		0.58	0.43	0.43	0.19	0.77		0.19	0.38	3.64	1.73	0.19	0.19	0.19	4.99	10.53	14.95	78			
Punjab	F	0.22	0.87	0.43	0.43	0.20	0.10	0.71		0.22	0.65		6.72	2.60	1.08	0.43	2.60	2.38	18.63	86		
T	0.20	0.41	0.51						0.20	0.20	0.10	5.09	2.14	0.61	0.10	0.31	3.87	1.93	16.68	164		

+ M = male, F = female, T = total

Table (2R): Incidence rate of temporary ailments by age separately by sex for different states and all-India, rural households :  
NSO 28th round.

RURAL

states--		age (years)									all ages	number of sample ailments
		0	1-4	5-14	15-24	25-44	45-59	60 & above	not recorded			
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)		
1. Andhra Pradesh	male	26.10	39.31	14.67	19.82	21.27	24.59	23.88		21.92	492	
	female	22.68	33.72	15.90	17.56	19.64	20.47	15.62		19.84	436	
	total	24.40	36.55	15.26	18.67	20.47	22.46	19.90		20.89	928	
2. Assam	male	4.88	12.01	8.65	5.67	7.85	23.66	14.51		9.43	130	
	female	14.64	9.57	7.64	5.03	7.18	6.95	6.83		7.34	91	
	total	8.76	10.94	8.16	5.34	7.57	12.52	11.69		8.46	221	
3. Bihar	male	10.65	10.50	3.87	3.41	5.63	5.35	6.86		5.48	154	
	female	4.83	8.88	2.10	2.10	4.26	4.85	3.62		3.90	106	
	total	7.83	9.69	3.06	2.78	4.92	5.09	5.21		4.69	260	
4. Gujarat	male	14.76	3.87	3.11	0.87	5.62	9.39	2.83		4.18	52	
	female	3.95	8.37	4.47	2.88	3.71	7.56	4.41		4.79	56	
	total	9.48	6.24	3.74	1.83	4.69	8.46	3.61		4.48	108	
5. Haryana	male	17.18	13.14	6.03	6.52	12.35	14.02	16.82		10.05	164	
	female	14.92	12.25	4.53	6.36	11.53	7.78	11.31		8.35	121	
	total	16.12	12.70	5.33	6.43	11.94	10.97	14.69		9.23	285	
6. Jammu & Kashmir	male	21.19	21.54	8.66	6.29	9.92	12.86	10.35		10.89	313	
	female	15.34	17.96	10.18	7.06	11.58	14.62	18.58		11.98	308	
	total	18.32	19.83	9.39	6.55	10.68	13.71	13.55		11.42	621	
7. Karnataka	male	14.36	13.17	6.40	4.59	11.13	11.93	6.14		8.65	122	
	female	8.52	12.40	6.84	5.53	8.79	8.16	11.57		8.20	113	
	total	11.43	12.77	6.62	5.06	9.93	10.04	8.69		8.44	235	
8. Kerala	male	64.15	82.25	41.36	30.98	28.75	31.78	19.60		39.25	562	
	female	51.29	70.34	36.44	23.79	32.34	32.99	20.19		35.49	534	
	total	57.63	76.46	38.90	28.26	30.68	32.41	19.91		37.29	1096	
9. Madhya Pradesh	male	24.77	12.94	8.18	9.69	11.39	12.89	13.55		11.04	307	
	female	14.11	18.97	7.62	6.04	7.56	8.99	15.32		9.65	255	
	total	19.53	15.95	7.91	7.91	9.57	10.86	14.36		10.34	562	
10. Maharashtra	male	32.32	37.82	15.20	12.02	18.93	25.05	20.29		19.96	479	
	female	29.96	26.74	13.85	9.03	12.93	17.30	11.80		14.70	391	
	total	32.22	33.23	13.01	10.66	16.36	21.52	16.54		17.78	870	
11. Orissa	male	34.58	31.03	15.92	14.01	17.65	12.88	12.16		17.69	230	
	female	9.63	14.97	11.54	6.89	14.28	7.60	13.97		11.44	152	
	total	22.80	23.04	13.70	10.36	15.97	10.01	13.01		14.53	382	

Table (2R) Contd. : Incidence rate of temporary ailments by age separately by sex for different states and all-India, rural households : Ncc 28th round.

RURAL -

states (1)		age (years)									number of sample ailments (11)
		0	1-4	5-14	15-24	25-44	45-59	60 & above	not recor- ded	all ages (10)	
		(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	
12. Punjab	male	48.05	36.43	19.87	18.41	18.26	18.55	26.51		22.46	341
	female	24.12	22.11	12.72	12.46	16.16	15.55	20.08		15.63	213
	total	37.02	29.78	16.58	15.59	17.23	17.08	23.81		19.25	554
13. Rajasthan	male	13.40	13.75	8.00	13.09	10.62	10.51	16.47		11.08	148
	female	20.17	7.44	7.69	10.65	8.18	11.73	10.81		9.26	116
	total	16.68	10.80	7.84	11.92	9.40	11.15	13.76		10.20	264
14. Tamil Nadu	male	28.24	32.58	16.22	12.87	22.27	19.02	24.63		20.18	299
	female	42.67	28.05	12.95	16.08	23.01	25.74	30.27		21.10	321
	total	35.53	30.36	14.59	14.52	22.65	22.47	27.31		20.65	620
15. Uttar Pradesh	male	18.46	14.39	6.33	6.30	8.68	11.89	12.63		9.27	348
	female	4.00	11.90	4.16	4.63	6.17	7.54	8.12		6.33	211
	total	11.69	13.22	5.36	5.53	7.45	9.67	10.60		7.95	559
16. West Bengal	male	33.43	18.94	10.45	11.01	9.45	10.72	17.88		12.37	290
	female	31.25	15.36	9.04	5.82	7.89	13.63	14.19		10.26	228
	total	32.39	17.21	9.76	8.52	8.70	12.22	16.21		11.35	518
17. all-India*	male	24.08	22.36	10.57	10.44	12.88	14.82	14.91	157.82	13.53	4675
	female	17.88	18.81	9.46	8.63	11.20	12.55	12.31	32.04	11.55	3937
	total	21.09	20.65	10.04	9.53	12.04	13.64	13.67	103.98	12.57	8612

\*including smaller states and Union territories.

SNC..

Table (2U). Incidence rate of temporary ailments by age separately by sex for different states and all-India, urban households : NSS 28th round.

URBAN

states (1)		age (years)							60 & not above record- ded	all ages of sample anilments
		0	1-4	5-14	15-24	25-44	45-59	above 59		
		(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
1. Andhra Pradesh	male	19.35	27.80	17.38	14.38	18.60	13.96	14.42	17.80	232
	female	54.88	44.25	11.89	11.52	18.71	13.53	16.60	18.62	235
	total	37.62	36.02	14.67	13.01	18.65	13.74	15.59	18.20	467
2. Assam	male		7.30	1.64	5.84	5.23	13.79		5.15	24
	female		23.83	6.53	9.07	1.17	3.40	5.74	7.23	26
	total		15.22	3.94	7.25	3.65	9.65	2.79	5.94	50
3. Bihar	male	19.24	12.44	5.70	4.67	6.45	7.10	6.11	6.87	70
	female	21.16	5.21	3.15	4.76	6.62	6.95	1.96	5.23	45
	total	20.15	8.99	4.48	4.71	6.52	7.02	3.99	6.05	115
4. Gujarat	male	28.68	0.45	0.96	2.83	4.03	3.79	10.47	2.47	31
	female	6.54	4.19	3.20	3.40	3.45	4.15	2.49	3.53	25
	total	18.21	1.35	1.55	3.09	3.76	3.96	6.38	2.85	56
5. Haryana	male	8.20	17.72	6.99	10.47	16.02	13.02	9.39	11.73	57
	female	20.83	4.29	4.35	1.17	6.30	6.65		4.73	20
	total	15.04	11.30	5.74	6.28	11.56	9.88	5.58	8.46	77
6. Jammu & Kashmir	male	23.48	10.03	7.31	7.27	9.18	11.18	20.78	9.32	106
	female	25.11	14.83	6.35	5.71	10.75	10.14	8.56	9.04	90
	total	24.26	12.30	6.86	6.53	9.90	10.69	15.62	9.22	196
7. Karnataka	male	25.14	23.84	5.48	3.84	8.22	10.78	6.76	8.76	73
	female		12.46	2.78	2.94	7.92	6.37	7.87	5.69	47
	total	13.03	18.14	4.14	3.41	8.08	8.52	7.34	7.24	120
8. Kerala	male	97.20	72.93	33.10	19.75	27.71	21.66	17.94	32.28	190
	female	114.18	76.46	34.62	21.46	28.99	29.37	34.94	35.22	205
	total	104.95	74.59	33.82	20.60	28.37	25.54	27.53	33.77	395
9. Madhya Pradesh	male	20.50	18.60	12.16	7.90	12.08	16.42	14.56	12.72	120
	female	35.00	18.72	6.15	4.42	13.17	10.06	3.90	9.80	84
	total	27.03	18.69	9.27	6.33	12.54	13.36	9.06	11.34	204
10. Maharashtra	male	45.41	38.00	17.86	16.94	21.42	24.65	16.21	21.70	413
	female	22.81	32.98	17.85	13.94	17.57	16.32	10.57	18.10	293
	total	35.37	35.52	17.85	15.65	19.77	20.86	13.25	20.06	706
11. Orissa	male	47.20	28.40	21.38	11.33	17.54	19.07	22.42	19.60	87
	female	43.48	21.69	11.93	10.37	8.44	23.26	20.58	14.34	54
	total	45.45	25.33	17.01	10.92	13.52	21.22	21.46	17.19	141

Table(2U) Contd. : Incidence rate of temporary ailments by age separately by sex for different states and all-India, urban households : NSS 28th round.

URBAN

states (1)		age (years)										all ages of sample ailments (11)
		0	1-4	5-14	15-24	25-44	45-59	60 & above	not recorded	all number (10)		
		(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)		
12. Punjab	male	34.19	33.47	13.77	11.04	14.56	5.91	12.04		14.95	78	
	female	28.57	37.22	16.80	11.21	20.21	12.36	21.98		18.63	86	
	total	31.12	35.22	15.25	11.11	17.15	9.05	16.53		16.68	164	
13. Rajasthan	male	25.16	16.76	3.27	7.85	10.97	4.92	8.30		8.39	55	
	female	5.79	8.22	4.69	5.41	3.84	9.35	17.84		6.45	39	
	total	15.08	12.47	3.95	6.77	8.59	7.17	13.30		7.46	94	
14. Tamil Nadu	male	41.96	41.96	19.12	9.50	24.12	15.40	18.82		21.08	298	
	female	20.06	34.88	14.10	14.42	24.18	24.60	26.93		20.93	286	
	total	31.19	38.48	16.63	11.97	24.14	20.01	23.07		21.01	584	
15. Uttar Pradesh	male	2.57	7.84	4.53	6.48	6.03	8.09	10.31		6.54	118	
	female	16.54	7.89	6.49	4.05	5.38	4.94	5.25		5.99	90	
	total	13.76	3.85	5.43	5.46	5.75	6.63	8.04		6.27	208	
16. West Bengal	male	71.95	52.43	17.92	11.59	11.28	23.18	14.41		18.60	247	
	female	42.59	39.20	16.00	9.73	13.35	8.90	12.93		15.65	164	
	total	58.09	46.07	16.99	10.76	12.09	17.38	13.67		17.30	411	
17. all-India*	male	33.69	23.92	11.40	9.61	13.87	14.36	13.09		13.87	2306	
	female	26.61	24.86	10.61	8.96	13.30	12.36	12.86		13.14	1855	
	total	30.29	24.36	11.02	9.30	13.64	13.41	12.97		13.53	4161	

SNC.

Table (3R) : Prevalence rate of temporary ailments by type of ailments separately by sex for different states and all-India, rural households : NSS 28th round

states	sex +	type of ailments																			not recorded	all types of ailments	number of sample ailments
		cholera	typhoid	dysentery (all forms)	diarrhoea	diphtheria	whooping cough	tetanus	acute poliomy- elitis	small pox	measles	mumps	malaria	influenza	pneumonia	food poisoning	accident	others					
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)		
Andhra Pradesh	M	-	0.36	0.88	0.18	-	0.53	0.05	-	0.96	0.18	-	3.05	6.58	0.22	0.04	0.75	10.28	8.79	32.85	737		
	F	-	0.22	0.62	0.05	-	0.63	-	-	0.80	0.18	0.09	3.21	5.61	0.14	0.05	0.40	8.71	10.24	30.95	681		
	T	-	0.29	0.75	0.12	-	0.58	0.02	-	0.88	0.18	0.05	3.13	6.10	0.18	0.05	0.58	9.51	9.52	31.94	1418		
Assam	M	0.14	0.44	3.56	0.37	0.50	0.72	0.07	-	0.94	-	0.07	1.91	5.94	0.14	-	0.29	1.16	3.30	19.55	270		
	F	-	0.16	1.86	0.40	0.16	0.40	-	-	0.73	-	0.08	0.98	6.53	0.08	-	0.40	0.97	2.25	15.00	186		
	T	0.08	0.31	2.75	0.38	0.34	0.57	0.04	-	0.83	-	0.08	1.47	6.21	0.11	-	0.34	1.07	2.82	17.40	456		
Bihar	M	0.07	0.07	1.13	0.46	-	0.50	-	-	0.71	0.14	-	0.32	1.49	0.18	-	0.43	5.85	0.35	11.70	330		
	F	0.04	0.11	0.81	0.22	-	0.40	0.04	0.04	0.51	0.04	-	0.18	0.99	0.04	-	0.04	4.76	0.40	8.62	235		
	T	0.05	0.09	0.97	0.34	-	0.45	0.02	0.02	0.61	0.09	-	0.25	1.24	0.11	-	0.23	5.31	0.38	10.16	565		
Gujarat	M	0.08	0.16	0.16	0.32	-	1.13	-	-	0.24	-	0.08	2.91	0.80	0.08	-	0.97	1.04	1.37	9.34	116		
	F	0.08	0.26	0.26	0.09	-	0.52	-	-	0.34	-	0.08	3.51	16.62	0.08	-	0.60	1.29	2.32	11.05	129		
	T	0.08	0.21	0.21	0.21	-	0.83	-	-	0.29	-	0.08	3.20	1.20	0.08	-	0.78	1.17	1.83	10.17	245		
Haryana	M	0.12	0.31	0.37	0.31	-	1.23	-	-	0.31	0.43	-	1.78	0.74	0.55	0.06	0.37	2.70	7.67	16.95	276		
	F	0.14	0.34	0.48	0.28	0.07	0.69	-	-	0.83	0.28	-	1.10	0.28	0.14	-	0.07	1.93	8.25	14.88	216		
	T	0.13	0.32	0.42	0.29	0.03	0.97	-	-	0.55	0.36	-	1.46	0.52	0.36	0.03	0.23	2.34	7.95	15.96	492		
Jammu & Kashmir	M	-	0.70	2.16	1.01	0.07	0.87	-	-	0.03	0.07	-	0.99	6.06	0.38	0.03	0.35	3.44	1.53	17.69	508		
	F	0.08	0.63	2.14	1.09	-	1.08	-	-	0.08	0.16	0.04	0.67	7.00	0.23	0.04	0.12	3.09	1.44	17.89	460		
	T	0.04	0.66	2.15	1.05	0.04	0.97	-	-	0.06	0.11	0.02	0.84	6.50	0.31	0.04	0.24	3.27	1.49	17.79	968		

+ M = male F = female T = total

Table (3R) Contd. : Prevalence rate of temporary ailments by type of ailments separately by sex for different states and all-India, rural households : NSS 28th round

states	sex +	type of ailments																			others	not recorded	all types of ailments	number of sample ailments
		cholera	typhoid	dysentery (all forms)	diarrhoea	diphtheria	whooping cough	tetanus	acute poliomyelitis	small pox	measles	mumps	malaria	influenza	pneumonia	food poisoning	accident							
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)			
Karnataka	M	-	0.14	1.35	-	-	1.13	-	0.07	0.14	-	0.14	0.14	1.56	-	0.21	0.35	3.76	3.09	16.08	227			
	F	-	0.07	0.65	-	-	1.16	-	0.07	0.29	0.07	0.15	0.15	1.16	-	-	0.36	3.12	6.46	13.71	189			
	T	-	0.11	1.00	-	-	1.15	-	0.07	0.22	0.04	0.14	0.14	1.36	-	0.11	3.36	3.44	6.78	14.92	416			
Kerala	M	0.14	0.07	3.77	1.05	0.07	1.96	-	-	0.84	0.98	0.07	0.14	7.26	0.14	-	3.56	30.73	23.81	74.59	1068			
	F	0.13	0.20	3.46	1.73	-	1.73	-	-	0.20	1.46	0.07	-	6.78	0.27	-	1.79	27.44	22.72	67.98	1023			
	T	0.14	0.14	3.61	1.40	0.03	1.84	-	-	0.51	1.23	0.07	0.07	7.01	0.20	-	2.66	29.04	23.26	71.21	2091			
Madhya Pradesh	M	-	0.36	0.25	0.32	0.14	0.79	-	0.04	0.72	0.97	0.14	4.95	0.47	0.39	0.07	0.36	1.84	0.04	21.85	609			
	F	-	0.11	0.26	0.23	0.11	0.79	0.04	-	0.91	0.42	0.08	4.96	0.49	0.26	0.19	0.11	0.86	-	19.82	524			
	T	-	0.24	0.26	0.28	0.13	0.79	0.02	0.02	0.81	0.70	0.11	4.96	0.48	0.33	0.13	0.24	11.37	0.02	20.89	1123			
Maharashtra	M	-	0.75	1.58	0.21	-	1.34	0.04	0.08	0.46	0.46	0.17	2.34	7.14	0.21	-	1.13	7.83	7.91	31.65	758			
	F	0.04	0.38	0.83	0.15	-	1.62	0.04	-	0.27	0.49	0.04	2.07	5.88	0.08	-	0.75	4.92	6.34	23.90	635			
	T	0.02	0.57	1.19	0.18	-	1.48	0.04	0.04	0.36	0.47	0.10	2.19	6.48	0.14	-	0.93	6.31	7.09	27.57	1393			
Orissa	M	-	0.38	1.77	0.15	0.08	1.15	-	0.08	1.15	0.62	-	2.46	3.39	0.15	-	0.92	8.85	7.93	29.08	378			
	F	-	0.08	1.58	0.45	-	0.22	-	-	0.38	0.37	0.16	1.73	2.33	-	0.08	0.52	6.18	5.34	19.42	258			
	T	-	0.23	1.67	0.30	0.04	0.68	-	0.04	0.77	0.49	0.08	2.09	2.85	0.08	0.04	0.72	7.50	6.62	24.20	636			
Punjab	M	0.66	0.72	2.24	0.92	-	1.45	-	0.20	0.46	0.53	0.07	8.96	2.50	1.32	-	1.12	2.77	6.52	30.44	462			
	F	0.15	0.66	0.95	0.15	-	2.35	0.07	-	0.51	1.10	0.07	6.53	2.57	0.88	0.07	0.22	1.91	6.02	24.21	330			
	T	0.42	0.69	1.63	0.56	-	1.87	0.03	0.10	0.49	0.80	0.07	7.81	2.53	1.11	0.03	0.69	2.36	6.28	27.47	792			

+ M = male, F = female, T = total

Table (3R) Contd. : Prevalence rate of temporary ailments by type of ailment separately by sex for different states and all-India, rural households : NSS 28th round

state	sex	type of ailments																			not recorded	all types of ailments	number of sample
		stomach	typhoid	dysentery (all forms)	diarrhoea	skin	scabies	scrofula	tuberculosis	small pox	measles	cholera	influenza	meningitis	food poisoning	accident	others						
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)		
Rajasthan	M	0.08	1.05	0.30	0.15	-	0.89	0.08	-	0.75	-	-	4.26	0.82	0.74	-	1.13	5.36	3.67	19.28	258		
	F	0.08	0.87	0.48	0.08	-	0.47	-	0.08	0.56	-	-	4.38	1.28	0.72	-	0.48	2.77	3.13	15.38	193		
	T	0.08	0.96	0.39	0.12	-	0.69	0.04	0.04	0.66	-	-	4.29	1.04	0.73	-	0.81	4.10	3.38	17.33	451		
Tamil Nadu	M	0.13	0.40	2.78	0.27	-	0.88	-	0.07	1.21	0.34	-	0.14	2.16	-	-	1.01	7.86	7.37	34.62	512		
	F	-	0.19	2.77	0.59	-	0.79	-	-	1.18	0.13	-	-	1.84	-	-	0.39	7.44	8.02	33.34	507		
	T	0.07	0.30	2.78	0.43	-	0.83	-	0.03	1.19	0.24	-	0.07	2.00	-	-	0.70	7.65	7.70	33.99	1019		
Uttar Pradesh	M	-	0.29	0.53	0.51	0.03	0.93	-	0.03	0.77	0.56	-	1.04	1.62	0.29	-	1.36	7.27	-	15.23	572		
	F	0.06	0.30	0.51	0.42	0.03	0.60	-	0.06	0.48	0.03	0.03	0.96	1.50	0.12	0.03	0.42	5.40	-	10.95	365		
	T	0.03	0.30	0.52	0.47	0.03	0.78	-	0.04	0.63	0.31	0.01	1.00	1.57	0.21	0.01	0.92	6.39	-	13.22	937		
West Bengal	M	-	1.02	3.62	1.75	-	0.47	0.13	0.04	0.55	0.17	0.09	0.13	7.03	0.17	-	2.44	10.54	2.43	30.58	717		
	F	0.18	0.50	3.32	1.12	-	0.50	-	-	0.54	0.13	0.05	0.14	5.62	-	-	1.26	8.77	2.61	24.74	550		
	T	0.09	0.77	3.47	1.45	-	0.48	0.06	0.02	0.55	0.15	0.07	0.13	6.35	0.08	-	1.86	9.66	2.52	27.71	1267		
* All-India	M	0.05	0.42	1.46	0.50	0.04	0.90	0.02	0.03	0.70	0.37	0.05	1.85	3.26	0.26	0.02	1.09	8.96	4.13	24.11	8219		
	F	0.05	0.28	1.21	0.41	0.02	0.78	0.01	0.02	0.55	0.24	0.05	1.69	2.93	0.14	0.03	0.51	7.55	4.23	20.70	6918		
	T	0.05	0.35	1.34	0.46	0.03	0.84	0.02	0.03	0.63	0.31	0.05	1.77	3.10	0.20	0.03	0.80	8.27	4.18	22.46	5137		

\* including smaller states and union territories.

+ M = male, F = female, T = total

Table (30) : Prevalence rate of temporary ailments by sex for different states and all-India, urban households : NSS 28th round

states	type of ailments												all types not recorded	all types sampled	number of elements						
	+ M	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)
Andhra Pradesh	M	-	0.54	0.92	-	-	0.15	-	0.08	0.77	0.15	0.08	2.45	3.76	-	-	1.61	8.44	9.74	28.69	374
	F	-	0.63	0.87	0.16	-	0.71	-	0.38	1.43	0.16	0.16	2.22	3.72	-	-	1.66	7.68	7.45	26.93	340
	T	-	0.58	0.90	0.08	-	0.43	-	0.08	1.09	0.16	0.12	2.34	3.74	-	-	1.64	8.07	8.61	27.84	714
Assam	M	-	0.42	1.72	0.42	-	-	-	-	-	0.42	-	-	-	-	-	3.65	-	-	0.85	2.53
	F	-	-	2.49	-	-	-	-	-	-	0.82	-	-	-	-	-	1.13	1.93	-	5.24	1.91
	T	-	0.24	2.06	0.24	-	-	-	-	-	0.59	-	-	-	-	-	0.49	2.89	-	0.12	2.76
Bihar	M	-	0.19	0.97	0.19	-	-	-	-	-	0.29	-	-	-	-	-	0.29	1.94	0.10	0.48	6.01
	F	-	0.12	0.58	0.35	-	-	-	-	-	0.12	0.46	0.12	-	-	-	0.23	2.08	-	0.12	4.86
	T	-	0.16	0.79	0.26	-	-	-	-	-	0.21	0.05	0.58	0.05	0.16	0.26	2.00	0.05	-	0.32	5.48
Gujarat	M	-	0.08	0.08	-	-	-	-	-	-	0.08	0.16	0.08	-	-	-	1.27	0.80	0.08	-	0.40
	F	0.14	0.70	-	-	-	-	-	-	-	2.28	-	-	-	-	-	0.29	2.68	0.71	-	0.14
	T	0.05	0.30	0.05	-	-	-	-	-	-	0.26	0.05	0.21	0.05	0.10	1.78	0.76	0.05	-	0.31	1.12
Haryana	M	0.21	0.62	0.82	0.21	-	0.62	-	-	-	0.21	-	-	-	-	-	1.23	0.82	0.41	-	2.26
	F	-	-	0.71	0.24	-	0.71	-	-	-	0.24	-	-	-	-	-	0.24	0.71	0.47	-	0.47
	T	0.11	0.33	0.77	0.22	-	0.66	-	-	-	0.11	0.11	-	-	-	-	0.77	0.77	0.44	-	1.43
Jammu & Kashmir	M	-	0.26	0.53	0.62	-	0.27	-	-	0.09	-	-	-	-	-	-	0.09	6.72	0.18	-	0.35
	F	0.10	0.10	1.11	0.61	-	0.30	-	-	0.10	-	-	-	-	-	-	0.20	7.82	0.10	-	0.40
	T	0.05	0.19	0.80	0.61	-	0.28	-	-	0.05	0.05	-	-	-	-	-	0.14	7.28	0.14	-	0.37

+ M = male, F = female, T = total

Table (3U) Contd. : Prevalence rate of temporary ailments by type of ailment separately by sex for different states and all-India, urban household : NSS 28th round

states	+ X es	type of ailments																			not recorded	all types of ailments	number of sample ailments
		cholera	typhoid	dysentery (all forms)	diarrhoea	diphtheria	whooping cough	tetanus	acute poliomy- elitis	small pox	measles	mumps	malaria	influenza	pneumonia	food poisoning	accident	others					
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)		
Karnataka	M	-	0.12	1.32	-	0.12	1.08	0.12	-	0.61	0.49	-	0.36	2.53	0.12	-	0.36	3.13	4.56	14.92	124		
	F	-	-	0.60	0.24	-	0.85	-	0.12	0.61	-	-	0.12	2.92	-	-	-	2.05	4.25	11.76	97		
	T	-	0.06	0.97	0.12	0.06	0.97	0.06	0.06	0.61	0.24	-	0.24	2.73	0.06	-	0.18	2.59	4.40	13.35	221		
Kerala	M	-	0.36	2.02	1.00	0.17	1.53	-	-	0.36	0.52	0.83	-	7.79	0.18	-	5.30	21.18	21.99	63.23	371		
	F	-	0.18	4.31	1.18	-	1.02	-	-	0.35	1.43	0.50	-	9.30	-	-	0.89	19.01	22.00	60.44	351		
	T	-	0.27	3.61	1.09	0.08	1.28	-	-	0.35	1.11	0.67	-	8.54	0.09	-	3.11	20.10	21.99	61.84	722		
Madhya Pradesh	M	0.11	0.64	1.17	0.21	-	0.85	0.11	0.11	0.95	0.11	0.71	3.71	1.17	0.32	-	1.06	12.93	-	23.56	222		
	F	0.23	1.40	0.47	0.70	-	1.28	0.12	0.35	0.82	-	-	3.62	0.70	0.47	-	0.23	9.33	-	19.72	169		
	T	0.17	1.00	0.83	0.44	-	1.06	0.11	0.22	0.89	0.06	0.06	3.66	0.94	0.39	-	0.67	11.22	-	21.72	391		
Mahara- shtra	M	-	0.68	1.63	0.53	0.05	1.11	-	0.10	0.42	0.32	0.21	1.53	6.22	0.05	-	1.63	12.35	8.66	35.49	675		
	F	-	0.50	1.29	0.37	-	0.31	0.06	0.06	0.37	0.12	0.06	1.05	5.89	-	-	0.99	8.95	8.28	28.30	458		
	T	-	0.60	1.48	0.46	0.03	0.74	0.03	0.08	0.40	0.23	0.14	1.31	6.06	0.03	-	1.33	10.78	8.48	32.18	1133		
Orissa	M	-	0.22	3.38	0.68	-	0.45	-	-	2.03	0.45	-	1.80	2.93	-	-	1.13	9.69	6.76	29.52	131		
	F	-	0.53	1.59	-	-	0.27	-	-	2.12	-	-	1.06	1.86	-	-	0.53	3.98	6.38	18.32	69		
	T	-	0.37	2.56	0.37	-	0.37	-	-	2.07	0.24	-	1.46	2.44	-	-	0.85	7.07	6.58	24.38	200		
Punjab	M	0.38	0.38	0.96	-	0.19	1.34	-	-	0.19	0.77	0.19	4.80	2.11	0.58	0.38	0.58	6.71	3.64	23.20	121		
	F	0.22	1.08	0.65	0.43	-	1.95	-	-	0.87	-	-	7.37	3.90	1.52	-	0.65	4.33	3.25	26.22	121		
	T	0.31	0.71	0.81	0.20	0.10	1.63	-	-	0.51	0.41	0.10	6.00	2.95	1.02	0.20	0.61	5.60	3.46	24.62	242		

+ M = male, F = female, T = total

Table (3U) Contd. : Prevalence rate of temporary ailments by type of ailment separately by sex for different states and all-India, urban households : NSS 28th round

type of ailments																						
states	M	F	male	female	typhoid	diphtheria	whooping cough	tetanus	acute poliomy-	measles	mumps	measles	influenza	food poisoning	accident	other	not recorded	all types of ailments	example of all types of ailments			
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	
M	-	0.15	0.62	0.31	-	0.46	-	0.45	-	1.20	0.61	0.46	-	0.63	6.41	1.66	12.56	85				
Rajasthan	F	-	0.82	0.32	0.33	-	0.32	0.33	-	0.19	0.17	1.67	0.32	0.34	-	0.50	5.28	1.85	12.12	73		
T	-	0.47	0.48	0.32	-	0.39	-	-	0.33	0.08	-	1.43	0.47	0.40	-	0.57	5.86	1.75	12.55	153		
Tamil Nadu	M	0.07	1.03	0.70	-	0.85	0.07	-	1.89	0.49	0.21	-	2.55	-	-	0.71	15.51	6.7	1.62	443		
	F	-	0.22	1.61	0.43	-	0.51	0.07	-	1.83	0.58	0.14	-	2.05	-	-	0.44	17.72	7.20	52.35	443	
	T	-	0.14	1.73	0.57	-	0.68	0.07	-	1.64	0.53	0.18	-	2.30	-	-	0.58	16.50	6.96	31.98	891	
Uttar Pradesh	M	0.05	0.33	0.39	0.17	-	0.50	-	0.05	0.66	0.39	0.44	1.49	0.06	-	1.10	5.52	-	11.23	203		
	F	-	0.40	0.33	0.36	0.09	-	0.20	-	0.03	0.66	0.20	0.33	1.66	0.13	-	0.40	5.025	-	9.76	147	
	T	1.03	-	-	-	-	-	-	-	-	-	-	0.39	1.57	0.09	-	0.78	5.40	-	16.54	350	
West Bengal	M	0.08	0.68	3.99	0.30	0.45	0.06	-	1.28	0.75	0.30	0.15	7.45	0.07	0.53	2.35	11.84	5.35	35.78	475		
	F	0.10	0.57	2.48	0.86	-	0.29	-	0.08	1.34	0.67	0.19	0.10	7.07	0.10	0.18	2.91	10.40	4.96	31.40	329	
	T	0.08	0.63	3.32	0.55	-	0.38	0.07	0.04	1.31	0.71	0.25	0.13	7.26	0.08	0.38	2.32	11.21	5.18	33.85	804	
All-India	M	0.03	0.38	1.41	0.30	0.03	0.65	0.03	0.04	0.77	0.33	0.14	1.13	3.25	0.11	0.06	1.29	8.78	4.66	23.39	3849	
	F	0.04	0.50	1.15	0.37	..	0.53	0.02	0.06	0.22	0.27	1.17	1.47	3.24	0.12	0.02	0.68	8.21	4.76	22.06	3104	
	T	0.04	0.44	1.29	0.33	0.01	0.60	0.03	0.05	0.79	0.30	0.12	1.15	3.24	0.11	0.04	1.01	8.51	4.71	22.77	6953	

\* including smaller states and union territories.

+ M = male, F = female, T = total

Table (4R): Prevalence rate of temporary ailments by age separately by sex for different states and all-India, rural households : 28th round

RURAL,

states	age (years)										number of sample ailments (-11)
	0	1-4	5-14	15-24	25-44	45-59	60 & above	not recor- ed	all ages		
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)		
Andhra pradesh	male	36.66	52.49	21.70	27.22	32.60	40.05	47.23		32.85	737
	female	36.46	46.29	22.45	26.18	31.59	37.14	35.45		30.95	681
	total	36.56	49.40	22.09	26.70	32.17	38.55	41.56		31.94	1418
Assam	male	7.32	22.52	15.77	14.86	16.89	55.67	31.68		19.55	270
	female	25.62	19.08	12.99	10.89	15.67	18.08	20.48		15.00	186
	total	14.61	21.00	14.44	12.83	16.39	30.60	27.56		17.40	456
Bihar	male	16.75	18.75	8.83	7.73	12.32	13.54	15.82		11.70	330
	female	11.27	15.46	5.94	3.98	8.68	12.00	11.87		8.62	235
	total	14.08	17.13	7.48	5.90	10.43	12.74	13.82		10.16	565
Gujarat	male	18.44	11.65	7.25	5.29	10.55	17.99	8.54		9.34	116
	female	11.65	21.63	7.77	6.75	10.40	15.08	13.18		11.05	129
	total	15.12	16.89	7.50	5.99	10.48	16.08	10.83		10.17	245
Haryana	male	28.64	20.90	10.12	12.36	20.92	20.69	30.45		16.95	276
	female	29.83	19.17	8.15	9.53	19.10	22.64	20.11		14.88	216
	total	29.23	20.07	9.21	11.02	19.99	21.61	26.43		15.96	492
Jammu & kashmir	male	29.08	31.45	12.85	8.81	15.55	23.61	29.60		17.69	508
	female	26.40	24.80	13.91	12.00	17.53	23.24	28.99		17.89	460
	total	27.76	28.24	13.36	10.29	16.49	23.42	29.36		17.79	968
Karnataka	male	25.86	18.18	9.96	11.29	18.56	26.84	24.56		16.08	227
	female	17.04	17.98	8.36	8.09	15.73	18.55	28.90		13.71	189
	total	21.44	18.07	9.19	9.70	17.12	22.69	26.62		14.92	416
Kerala	male	134.12	166.35	72.81	55.33	59.50	57.58	40.23		74.59	1068
	female	108.27	147.37	67.83	47.48	59.74	59.64	43.26		67.98	1023
	total	121.02	157.16	70.31	51.23	59.66	58.63	41.81		71.21	2091
Madhya Pradesh	male	43.32	23.66	16.71	17.48	22.31	28.56	30.42		21.85	609
	female	28.24	33.80	16.16	13.10	18.74	17.66	28.10		19.82	524
	total	35.91	28.73	16.45	15.36	20.63	22.90	29.24		20.89	1133
Maharashtra	male	43.64	58.40	24.13	20.49	28.42	41.21	38.22		31.65	759
	female	48.88	41.93	19.69	13.45	23.11	30.46	25.94		23.90	635
	total	46.29	50.07	21.90	16.47	25.62	35.28	32.02		27.57	1393

Contd.

Table (4E) / Prevalence rate of temporary ailments by age separately by sex for different states and all-India, rural households :  
NSS 28th Round

RURAL

states		age (years)									all ages	number of sample ailments
		0	1-4	5-14	15-24	25-44	45-59	60 & above	not recorded			
		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
Orissa	male	43.23	43.18	24.02	25.13	27.42	25.02	42.04		29.08	378	
	female	16.08	27.91	19.94	11.48	20.32	15.84	27.95		19.42	258	
	total	30.40	35.50	21.95	18.13	23.88	20.02	35.48		24.20	636	
Punjab	male	66.37	50.77	24.51	23.02	24.24	29.98	41.10		30.44	462	
	female	37.52	33.15	18.42	18.28	24.73	31.10	31.04		24.21	330	
	total	53.08	42.60	21.68	20.76	24.50	30.55	36.85		27.47	792	
Rajasthan	male	18.66	22.68	14.46	17.75	19.52	25.49	28.64		19.28	258	
	female	20.17	16.15	11.70	24.18	16.31	17.23	19.24		15.35	193	
	total	19.39	19.62	13.17	17.00	17.91	21.18	24.15		17.33	451	
Tamil Nadu	male	41.21	53.17	24.33	19.29	40.00	37.07	55.37		34.62	512	
	female	55.25	42.90	19.86	24.27	37.98	39.93	56.02		33.34	507	
	total	48.32	48.11	22.12	21.88	38.96	38.54	55.14		33.99	1019	
Uttar Pradesh	male	26.38	21.89	9.58	10.57	16.04	19.48	22.43		15.23	572	
	female	14.99	20.17	6.15	7.54	10.94	12.18	16.62		10.95	365	
	total	21.06	21.08	8.04	9.14	13.53	15.77	19.83		13.22	937	
West Bengal	male	74.86	47.34	25.20	23.18	26.39	30.73	44.72		30.58	717	
	female	66.05	38.76	19.82	15.42	23.24	27.26	35.46		24.74	550	
	total	70.65	43.19	22.57	19.44	24.87	28.94	40.29		27.71	1267	
All- India*	male	38.71	37.83	18.24	18.00	23.44	28.30	31.32	157.82	24.11	8219	
	female	32.55	33.07	15.98	14.78	20.76	22.97	25.96	210.95	20.70	6918	
	total	35.72	35.50	17.15	16.38	22.10	25.51	28.78	180.54	22.46	15137	

including Smaller states and union Territories.

Table (4U) : Prevalence rate of temporary ailments by age separately by sex  
for different states and all-India, urban households : NSS  
28th round

URBAN

states	0	age ( years )								all ages	number of sample ailments
		1-4	5-14	15-24	25-44	45-59	60 & above	not recorded			
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	
Andhra Pradesh	male	48.40	43.47	25.79	19.69	31.32	22.99	36.86	28.69	374	
	female	70.14	61.39	17.54	16.45	25.62	26.31	26.28	26.93	340	
	total	59.56	52.42	21.72	18.13	28.62	24.72	31.19	27.84	714	
Assam	male	12.77	16.82	4.11	9.69	7.46	18.28	32.60	10.01	47	
	female	29.00	13.13	15.30	7.04	6.80	28.71		13.79	50	
	total	6.10	22.65	8.34	12.14	7.29	13.71	30.70	11.66	97	
Bihar	male	33.66	22.95	9.26	8.41	11.01	16.23	12.23	12.21	126	
	female	21.16	13.56	7.88	6.80	13.70	15.06	3.92	10.55	91	
	total	27.71	18.48	8.60	7.76	12.20	15.67	7.99	11.42	217	
Gujarat	male	34.55	2.71	2.10	5.09	8.55	6.34	23.50	5.03	63	
	female	6.54	5.59	6.41	6.14	11.40	13.77	10.02	8.47	60	
	total	21.30	3.40	3.24	5.56	9.88	9.91	16.59	6.26	123	
Jammu & Kashmir	male	23.48	14.14	10.06	9.98	12.58	15.88	26.99	12.66	144	
	female	25.10	17.64	8.67	7.80	16.12	19.36	17.20	12.98	129	
	total	24.26	15.82	9.40	8.98	14.19	17.53	22.85	12.80	273	
Karnataka	male	50.41	30.65	10.53	6.07	14.06	24.33	13.54	14.92	124	
	female	10.77	16.95	7.89	6.46	17.36	10.13	17.63	11.76	97	
	total	31.32	23.78	9.22	6.26	15.69	17.01	15.72	13.35	221	
Kerala	male	171.36	142.92	62.00	38.69	49.08	54.96	53.63	63.23	371	
	female	133.80	138.39	59.62	40.38	45.77	54.29	63.01	60.44	351	
	total	154.40	140.76	60.84	39.53	47.38	54.61	58.91	61.84	722	
Madhya pradesh	male	32.79	35.14	22.04	15.78	24.59	26.69	22.87	23.56	222	
	female	65.00	33.27	13.11	6.96	23.27	27.91	19.52	19.72	169	
	total	47.29	34.24	17.74	11.78	23.95	27.26	21.15	21.72	391	
Maharashtra	male	72.85	62.02	29.33	25.99	33.12	45.67	34.72	35.49	675	
	female	34.19	49.14	25.27	21.23	29.64	27.63	22.16	28.30	458	
	total	55.72	55.67	23.37	23.94	31.64	37.45	28.13	32.18	1133	

Table (4U) Contd. : Prevalence rate of temporary ailments by age separately by sex for different states and all-India, urban households : NSS 28th round

URBAN

states	age (years)									all ages	number of sample ailments
	0	1-4	5-14	15-24	25-44	45-59	60 & above	not recor- ded			
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	
Orissa	male	66.04	38.54	33.36	14.74	26.73	43.60	22.42	29.52	131	
	female	43.48	28.92	14.91	11.85	11.60	31.01	28.81	18.32	69	
	total	55.55	34.14	24.83	13.49	20.04	37.13	25.75	24.38	200	
Punjab	male	51.29	40.91	17.82	15.28	27.61	23.63	15.05	23.20	121	
	female	35.71	43.78	21.00	17.30	30.32	26.79	29.31	26.22	121	
	total	42.89	42.25	19.37	16.20	29.68	25.16	21.49	24.62	242	
Rajasthan	male	31.63	18.17	6.54	11.33	16.46	12.88	19.60	12.96	85	
	female	23.16	18.19	8.80	7.31	9.21	15.63	27.83	12.12	73	
	total	27.22	18.18	7.62	9.55	14.76	14.28	23.92	12.55	158	
Tamil Nadu	male	67.14	58.13	28.13	16.13	34.26	26.00	34.87	31.62	448	
	female	43.18	56.35	20.32	21.84	36.87	37.24	43.89	32.35	443	
	total	55.35	57.28	24.28	18.97	35.19	31.63	39.61	31.98	891	
Uttar Pradesh	male	33.40	11.76	8.87	8.10	10.44	15.64	17.19	11.23	203	
	female	18.91	11.52	9.14	6.62	10.78	9.88	10.50	9.76	147	
	total	26.38	11.63	8.97	7.49	10.60	12.97	14.18	10.54	350	
West Bengal	male	127.14	97.84	33.16	20.92	25.73	41.48	30.28	35.78	475	
	female	80.50	66.40	30.91	20.37	27.11	24.01	34.51	31.40	329	
	total	107.37	82.73	32.07	20.67	26.26	34.38	32.40	33.85	804	
All- India	male	56.96	38.87	18.88	15.26	23.39	27.02	26.05	23.39	3849	
	female	41.10	39.25	17.28	14.56	22.80	23.28	25.06	22.06	3104	
	total	49.34	39.04	18.14	14.97	23.13	25.24	25.57	22.77	6953	

\* including smaller states and union Territories.

Table(5.R): Average duration (days) of temporary ailments terminating in recovery by type of ailment separately by sex for different states and all-India, rural households : NSS 28th round

states	sex	type of ailments																			not recorded	all types of ailments	number of sample ailments
		cholera	typhoid	dysentery (all forms)	diarrhoea	diphtheria	whooping cough	tetanus	acute poliomyelitis	small pox	measles	mumps	malaria	influenza	pneumonia	food poisoning	accident	others					
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)		
Andhra Pradesh	M	19.23	4.12				8.38			6.64	13.94		6.89	7.16	4.00		6.33	7.50	6.25	7.25	266		
	F		5.40							7.17	7.00		9.83	6.84				6.43	4.97	6.60	250		
	T	19.23	4.65				8.38			6.89	12.23		8.27	7.00	4.00		6.33	7.05	5.46	6.94	516		
Assam	M	9.50	8.00	6.44	5.00	7.00	16.00			13.33		10.00	5.89	6.78	9.50		8.67	5.83	9.00	7.28	106		
	F		8.00	13.75	6.00	10.00	8.00			14.60			6.33	7.20	8.00			5.00	4.65	8.33	78		
	T	9.50	8.00	9.08	5.40	7.75	4.00			14.13		10.00	6.07	6.98	9.00		8.67	5.83	6.08	7.72	184		
Bihar	M	3.00	4.00	9.11	3.88		14.57			10.00	7.00		7.20	6.50	10.00			6.75	11.26	10.00	9.36	137	
	F	5.00	20.00	7.00	4.67		32.40			8.25	7.00		16.00	7.07				8.08	9.00	9.86	77		
	T	4.00	14.67	8.27	4.09		22.00			9.22	7.00		10.50	6.69	10.00			6.75	10.09	9.33	9.54	214	
Gujarat	M	5.00			3.00			8.33						6.79	4.00			8.64	2.00	7.16		42	
	F	4.00						8.00		10.50				7.71	5.10			2.50	6.24	2.50		54	
	T	4.50			3.00			8.25		10.50				7.34	4.92			6.20	5.38	5.60		96	
Haryana	M	4.00	20.00	10.00	6.33			6.67			10.00	4.00		8.60	4.33	4.33	3.00	2.00	6.42	12.98		86	
	F	1.00		6.50	22.33					18.63	6.50		6.83	8.00				5.00	8.96			55	
	T	2.50	20.00	8.25	14.33			6.67			16.90	5.43		7.94	5.25	4.33	3.00	2.00	6.06	11.39		141	
Jammu & Kashmir	M		10.85	9.58	5.95		11.66			8.00	6.00		9.60	5.87	18.64	4.00	10.00	8.22	11.00			218	
	F	4.00	11.67	5.29	4.94		11.96				6.32		9.01	6.35	10.00			7.99	11.53			189	
	T	4.00	11.08	6.35	5.47		11.81			8.00	6.24		9.44	6.11	16.93	4.00	10.00	8.11	11.30			407	

+ M = male, F = female, T = total

Table (5R) Contd : Average duration (days) of temporary ailments terminating in recovery by type of ailment separately by sex for different states and all-India, rural households : NSS 28th round

states	+ x e s	type of ailments																			not recorded	all types of ailments	number of sample ailments
		cholera	typhoid	dysentery (all forms)	diarrhoea	diphtheria	whooping cough	tetanus	acute polioy- elitis	small pox	measles	mumps	malaria	influenza	pneumonia	food poisoning	accident	others					
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)		
Karnataka	M		7.00	11.12			4.50						6.00	5.44		15.00	20.00	6.50	4.47	6.74	65		
	F			3.50			4.33		30.00			8.00	10.00	5.09			60.05	4.65	5.82	5.66	59		
	T		7.00	9.60			4.40		30.00			8.00	8.00	5.25		15.00	9.50	5.61	5.11	6.13	124		
Kerala	M	6.50		4.80	5.60		7.50			15.00	10.43		12.00	5.48				9.43	6.74	5.57	8.24	350	
	F	6.50		5.93	3.59		10.50			20.00	11.07			6.85	12.00		15.00	6.24	6.33	6.66	321		
	T	6.50		5.37	4.33		9.00			15.71	10.86		12.00	6.17	12.00		11.83	6.52	6.96	6.75	671		
Madhya Pradesh	M		13.50	6.50	14.17		14.75			10.00	18.80		8.02	11.57	7.40			5.00	8.72		9.28	181	
	F		14.00		7.00		10.00	8.00		8.38	7.00		8.41	9.50			5.00	8.59		8.56	109		
	T	13.67	6.50	11.78		13.17	8.00		9.00	15.43		8.18	10.82	7.40			5.00	8.67		9.01	290		
Mahara- shtra	M		15.60	4.83	6.50		9.80		45.00	14.24	9.25		7.17	7.86	10.77			6.00	5.85	7.05	7.58	265	
	F		21.56	7.38	3.00		8.40			21.00	8.00		8.88	5.97				7.33	8.05	7.74	7.73	220	
	T		17.30	5.85	5.80		9.10		45.00	16.49	8.63		8.00	6.97	10.77			6.75	6.80	7.39	7.65	485	
Orissa	M		15.50	2.11	12.00	4.00	6.33			10.50	7.40		7.52	6.47	9.00			8.00	7.33	9.81	7.94	173	
	F			11.00	4.75		7.00			6.00	8.00	8.00	8.80	6.25			12.00	6.50	11.73	9.08	117		
	T		15.50	8.48	6.20	4.00	6.50			8.70	7.50	8.00	7.94	6.39	9.00			9.00	7.65	10.65	8.40	290	
Punjab	M	3.88	11.33	4.07	3.78		13.00		14.00	11.20	8.00		6.26	5.65	8.17			3.00	6.00	6.96	6.42	191	
	F	5.00	63.00	7.60	4.00		5.00	20.00		8.00	8.00	1.00	6.78	3.92	8.67			8.00	10.58	5.57	7.61	135	
	T	4.00	32.00	5.00	3.80		10.71	20.00	14.00	9.78	8.00	1.00	6.46	4.90	8.42			5.50	8.50	6.20	6.91	326	

+ M = male, F = female, T = total

Table(5R) Contd. : Average duration (days) of temporary ailments terminating in recovery by type of ailment separately by sex for different states and all-India, rural households : NSS 28th round

states	sex	type of ailments																			others	not recorded	all types of ailments	number of sample ailments	
		cholera	typhoid	dysentery (all forms)	diarrhoea	diphtheria	whooping cough	tetanus	acute poliomy- elitis	small pox	measles	mumps	malaria	influenza	pneumonia	food poisoning	accident	(19)	(20)	(21)	(22)				
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)				
Rajasthan	M	2.00	28.58				3.00			13.32			13.81	8.00	4.76		6.00	7.36	9.49	11.13	78				
	F												10.34	31.74	7.00		7.00	10.13	4.26	10.60	52				
	T	2.00	28.47				3.00			13.32			12.20	22.31	5.21		6.50	8.43	7.66	10.92	130				
Camil Nadu	M	11.00	8.66	4.16	3.00		3.50			11.29	7.00		4.00	5.90			6.50	6.18	6.11	6.18	191				
	F		19.00	5.48	7.41					10.59	9.00			7.35			2.00	5.11	7.24	6.36	198				
	T	11.00	13.82	4.89	6.44		3.50			10.94	8.00		4.00	6.58			3.71	5.62	6.67	6.30	389				
Uttar Pradesh	M		23.33	5.00	6.40		14.33			8.00	5.00		8.21	5.65	5.83		23.56	6.59		8.21	146				
	F		33.50	6.00	5.00		6.50			4.67			8.26	7.70	3.50		9.86	9.53		8.90	88				
	T		27.40	5.20	6.00		11.20			7.09	5.00		8.24	6.68	5.25		17.56	7.49		8.47	234				
West Bengal	M		16.53	10.63	5.93		8.33	10.00		13.24	7.67	14.00	2.00	6.24	11.99		14.00	9.90	25.02	9.42	275				
	F		8.50	27.40	6.31	4.64		9.20		14.25	5.00		4.50	6.12					8.15	6.63	7.54	178			
	T		8.50	20.76	8.95	5.52	8.72	10.00		13.75	7.00	14.00	3.67	6.19	11.99		14.00	9.12	15.33	8.68	453				
* All-India	M	5.49	15.92	6.65	5.96	4.38	9.55	10.00	19.98	10.50	10.06	12.48	8.11	6.61	7.98	12.95	12.07	8.15	6.89	7.88	2817				
	F	6.06	24.81	7.19	5.25	10.00	12.26	12.26	30.00	10.13	9.05	6.73	8.68	6.72	7.76		8.20	11.23	5.66	8.52	2240				
	T	5.70	18.84	6.89	5.66	5.15	10.52	10.87	22.97	10.33	9.61	9.02	8.36	6.66	7.94	12.95	10.73	9.45	6.27	8.16	5057				
	M																								
	F																								
	T																								

Table (50) : Average duration (days) of temporary ailments terminating in recovery by type of ailment separately by sex for different states and all-India, urban households :  
NSS 28th round.

states	+ sex	type of ailments																			not recorded	all types of ailments	number of sample ailments	
		cholera	typhoid	dysentery (all forms)	diarrhoea	diphtheria	whooping cough	tetanus	acute poliomy- elitis	small pox	measles	mumps	malaria	influenza	pneumonia	food poisoning	accident	others						
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)			
Andhra Pradesh	M		24.00	2.50					7.00	9.83	7.00		5.91	5.70				1.50	6.96	5.20	6.35	165		
	F		14.40	3.75	1.50				7.50	6.83	5.00		5.24	5.26				11.71	15.26	5.43	8.59	133		
	T		18.00	3.33	1.50				7.00	8.14	6.00		5.63	5.46				9.44	10.46	5.27	7.35	298		
Assam	M			3.67											6.73				6.00	15.07	8.01	21		
	F			7.35											7.00				10.17		7.94	8		
	T			5.53											6.99				8.12	15.07	7.99	29		
Bihar	M			3.00	10.00			9.00			12.00		5.00	4.11				7.50	4.82	18.00	6.44	54		
	F			15.00	3.00						7.00		13.00	4.60				4.00	6.91	10.00	6.65	40		
	T			9.00	5.33			9.00			12.00	7.00	8.20	4.37				6.33	5.71	16.67	6.53	94		
Gujarat	M		10.00	2.33							10.00	15.00		5.66	4.99				4.32	9.00	3.52	5.85	28	
	F		7.00										4.00	8.27	9.00				3.00	7.00	12.94	8.51	24	
	T		8.49									10.00	15.00	4.00	7.10	6.32				4.00	7.80	7.27	7.07	52
Haryana	M		33.00	2.33															10.00	4.86	5.88	8.43	28	
	F		33.00	2.33															5.88	8.67	5.33	15		
	T		33.00	2.33															10.00	5.40	6.64	7.35	43	
Jammu & Kashmir	M			7.46	6.03								7.00		5.00	4.90	12.00				5.80	5.00	5.32	63
	F			7.34	4.00								7.00		5.00	4.44				9.08	7.77	5.60	54	
	T			7.37	5.35								7.00		5.00	4.68	12.00				6.89	6.16	5.45	117

Table (5U) Contd. : Average duration (days) of temporary ailments terminating in recovery by type of ailment separately by sex for different states and all-India, urban households :  
NSS 28th round

states	+ sex	type of ailments																			not recorded	all types of ailments	number of sample ailments
		cholera	typhoid	dysentery (all forms)	diarrhoea	diphtheria	whooping cough	tetanus	acute poliomy- elitis	small pox	measles	mumps	malaria	influenza	pneumonia	food poisoning	accident	others					
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)		
Karnataka	M			3.76	3.00	7.00	4.47			8.00	8.00		3.00	6.83			4.97	8.08	5.47	6.26	47		
	F			3.49	3.00	7.00	4.50			7.50			5.00	4.74			4.97	5.36	3.43	4.57	33		
	T			3.67	3.00	7.00	4.48			7.80	8.00		4.01	5.63				6.10	4.57	5.56	80		
Kerala	M			6.23	7.00						7.00	9.60		4.69			9.80	6.93	7.83	7.07	129		
	F			4.13	7.80						9.40	11.00		5.91			4.00	5.29	8.81	6.45	141		
	T			4.00	7.50						9.09	10.00		5.42			9.33	6.25	8.35	6.75	270		
Madhya Pradesh	M		13.00	4.50	7.00		10.00	10.00	10.00	4.00			7.00	5.40	6.00					7.21	70		
	F		2.00	21.00	9.33		13.50		4.00	7.50			4.92	4.50	11.00		10.00	7.10		7.45	49		
	T		2.00	17.00	4.50	8.40	12.33	10.00	7.00	5.75			6.04	5.14	8.50		10.00	7.30		7.31	119		
Mahara- shtra	M		14.23	6.43	3.34		8.10		30.00	7.00	11.53		7.64	6.98	8.00		6.57	5.40	4.09	5.92	317		
	F		16.33	6.38	1.50				1.00	25.00	7.00		6.19	6.15			12.00	4.86	5.14	5.68	201		
	T		14.13	6.41	2.61		8.10	1.00	27.50	7.00	11.53		7.29	6.66	8.00		8.20	5.19	4.52	5.82	518		
Orissa	M			3.33	7.50		3.00			10.00	10.00		8.00	5.28					5.56	6.80	6.29	62	
	F			15.00	3.20					9.00			4.00	4.33					7.75	5.50	5.92	26	
	T			15.00	3.29	7.50		3.00		9.75	10.00		7.00	5.00					6.00	5.67	6.18	88	
Punjab	M		7.50	30.00	3.33			2.00					7.54	4.67	3.00				4.67	6.50	6.30	40	
	F		8.00	8.50		6.00		4.00			11.50			5.05	5.18	8.25				3.25	5.25	5.56	50
	T		7.67	15.67	2.50	6.00		2.50			11.50	8.00		6.86	5.07	7.20				4.23	5.88	5.89	90

+ M = male, F = female, T = total

Table (50) Contd. : Average duration (days) of temporary ailments terminating in recovery by type  
of ailments separately by sex for different states and all-India, urban households:  
NSS 28th round

states	sex	type of ailments																			not recorded	all types of ailments	number of sample ailments
		cholera	typhoid	syphilitic forms (all forms)	diarrhoea	diphtheria	whooping cough	tetanus	acute poliomyelitis	small pox	measles	mumps	malaria	influenza	pneumonia	food poisoning	accident	others					
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)		
Rajasthan	M			1.00	3.00		6.00			8.00			11.86	3.00	5.00				6.54		6.71	27	
	F				*		4.00				4.00			3.27	6.00	3.00				5.35	6.43	4.93	22
	T			1.00	3.00		5.00			8.00	4.00		7.50	4.01	4.00				6.06	5.43	5.90	49	
Tamil Nadu	M			6.97	2.25		5.00	10.00		11.39	7.16	7.00		5.01					1.67	5.59	4.69	5.64	192
	F			15.66	5.20	9.34		8.23	2.00		12.85	9.60	8.50		4.84				2.66	5.33	5.69	6.15	189
	T			15.66	6.04	5.27		7.14	6.00		12.15	8.37	1.19		4.93				2.17	5.46	5.15	5.89	381
Uttar Pradesh	M			30.00	3.50				15.00		8.00			11.50	7.18				12.67	6.95		8.33	52
	F			16.50	7.75						7.63			6.00	6.53	7.00			7.00	6.50		7.11	57
	T			21.00	6.33				15.00		7.82			9.14	6.79	7.00			10.40	6.79		7.69	109
West Bengal	M			14.33	5.63	5.33		6.01			15.00	8.55		5.50	5.30				12.43	5.82	11.06	5.64	213
	F			3.00	18.20	5.60	5.00		8.50		11.83	6.60	7.00	11.00	6.50				6.00	7.73	11.64	6.92	135
	T			3.00	16.09	5.62	5.14		7.01		13.54	7.85	7.00	7.33	5.84				11.14	6.43	11.25	6.18	348
* all-India	M			7.50	17.59	5.32	4.79	7.00	6.39	11.44	15.92	10.21	8.57	9.02	6.91	5.78	6.89	12.43	6.76	6.90	5.05	6.58	1548
	F			4.09	15.33	5.57	5.64		7.84	1.56	14.59	10.03	7.98	7.67	6.09	5.75	7.85	6.00	8.06	7.04	6.23	6.80	1212
	T			5.32	16.43	5.44	5.23	7.00	6.92	7.34	15.34	10.13	8.31	8.27	6.54	5.76	7.46	11.14	7.20	6.96	5.55	6.68	2760

sc/

+ M = male, F = female, T = total \* including smaller states and union territories.

ole (GR): Average duration (days) of temporary ailments terminating in recovery by age separately by sex for different states and all-India, rural households : NSS 28th round.

States		age (years)									all ages of sample ailme- nts	number
		0	1-4	5-14	15-24	25-44	15-59	60 & above	not recor- ded			
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)		
1. Andhra- Pradesh	male	7.32	5.57	7.53	7.85	7.60	7.81	7.97		7.25	266	
	female	6.47	5.38	6.94	5.77	7.45	7.08	7.98		6.60	250	
	total	6.40	5.49	7.26	6.49	7.53	7.43	7.98		6.94	516	
2. Assam	male	5.00	8.21	7.10	6.71	7.17	5.91	8.00		7.28	106	
	female	6.53	11.67	8.72	8.31	7.36	5.89	6.00		8.33	78	
	total	5.25	9.50	7.82	7.40	7.61	5.90	7.33		7.72	184	
3. Bihar	male	8.67	7.54	7.57	7.50	13.80	7.33	10.18		9.56	157	
	female	16.20	11.21	11.71	5.50	7.51	9.20	8.50		9.86	77	
	total	13.58	8.76	8.87	6.64	11.76	8.27	9.73		9.54	214	
4. Gujarat	male	5.00	4.99	5.46	7.53	8.86	6.98			6.53	42	
	female		6.91	5.86	7.75	6.86	7.50	8.00		6.53	54	
	total	5.00	5.61	5.67	7.62	7.49	7.24	8.00		6.53	96	
5. Haryana	male	5.53	4.71	7.38	5.80	7.67	6.42	34.55		9.91	86	
	female	20.50	9.79	11.00	4.33	5.83	9.33	4.50		10.15	55	
	total	17.71	7.00	9.00	5.46	6.75	7.00	29.92		10.00	141	
6. Jammu & Kashmir	male	10.79	7.47	6.41	8.25	7.24	8.47	9.55		7.62	218	
	female	6.66	5.32	7.27	3.28	6.89	7.68	10.16		7.25	189	
	total	8.88	6.69	6.77	8.27	7.08	8.08	9.70		7.45	407	
7. Karnata- ka	male	5.00	5.50	8.56	4.45	7.30	5.00	14.00		6.74	65	
	female	2.00	3.82	3.64	6.50	5.88	7.00	11.00		5.66	59	
	total	4.40	4.62	6.56	5.18	6.60	5.79	11.86		6.13	124	
8. Kerala	male	3.77	6.53	6.13	6.77	3.24	5.33	6.86		8.24	350	
	female	3.57	7.36	6.44	6.20	6.13	6.97	6.92		6.66	321	
	total	8.70	6.92	6.27	6.51	10.00	6.88	6.42		6.75	671	
9. Madhya- Pradesh	male	4.83	9.06	7.14	6.53	9.97	13.48	10.89		9.28	181	
	female	11.55	7.94	6.85	7.46	7.71	9.92	13.25		8.56	109	
	total	3.67	9.10	7.01	6.91	9.20	12.26	11.81		9.01	290	

Table (6R) (Contd.) : Average duration (days) of temporary ailments terminating in recovery by age separately by sex for different states and all-India, rural households : NSS 20th round.

RURAL

States		age years									all ages of sample	number of ailme- nts
		0	1-4	5-14	15-24	25-44	45-59	60 & above	not record ded	all ages		
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)		
10. Maharashtra	male	4.50	6.89	7.91	7.35	9.08	6.48	9.02	7.58	265		
	female	7.68	6.65	8.47	6.93	8.74	6.18	9.21	7.73	220		
	total	5.74	6.78	8.17	7.16	8.93	6.35	9.11	7.65	485		
11. Orissa	male	4.25	3.48	5.98	6.71	11.05	7.87	11.95	7.94	173		
	female	4.50	12.93	9.41	11.64	7.87	7.18	5.70	9.08	117		
	total	4.35	8.28	7.54	8.26	9.62	7.58	9.87	8.40	290		
12. Punjab	male	6.11	6.33	5.92	7.00	7.41	6.23	4.42	6.42	191		
	female	5.00	10.65	5.42	5.91	6.62	13.50	9.43	7.61	135		
	total	5.71	7.86	5.73	6.52	6.98	10.76	5.77	6.91	326		
13. Rajasthan	male	6.32	9.67	10.42	7.79	9.25	11.91	29.01	11.13	78		
	female	8.36	13.69	8.14	11.60	7.99	8.51	10.60	10.60	52		
	total	1.32	9.32	11.79	7.93	10.46	10.16	23.96	10.92	130		
14. Tamil Nadu	male	4.21	5.01	6.37	4.92	6.90	7.79	5.18	6.18	191		
	female	4.45	6.81	6.53	4.49	6.42	9.82	5.37	6.36	198		
	total	4.32	5.83	6.45	4.61	6.67	8.66	5.28	6.30	389		
15. Uttar Pradesh	male	5.67	6.33	8.73	9.05	8.58	6.81	11.60	8.21	146		
	female	5.50	6.96	12.42	6.85	10.11	7.00	14.43	8.90	88		
	total	5.60	6.65	9.79	8.23	9.09	6.88	12.76	8.47	234		
16. West Bengal	male	7.55	10.45	9.88	9.14	8.10	9.55	9.62	9.42	275		
	female	6.8	9.12	8.00	7.03	7.36	8.54	5.50	7.54	178		
	total	6.71	10.00	9.19	8.33	7.77	9.20	7.97	8.68	453		
17. all- India	male	6.00	7.01	7.52	7.41	8.69	8.02	10.39	7.88	2817		
	female	7.83	7.61	7.89	13.53	7.41	7.86	8.00	11.34	8.52	2240	
	total	6.80	7.27	7.68	10.22	8.11	7.95	9.37	6.96	5057		

\* including smaller states and Union Territories.

sd/

contd.

Table (6U) : Average duration (days) of temporary ailments terminating in recovery by age separately by sex for different states and all-India, urban households : NSS 28th round.

URBAN

States		age (years)									all number ages of sample ailme- nts
		0	1-4	5-14	15-24	25-44	45-59	60 & above	not recor- ded		
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	
10. Maharashtra	male	4.01	5.56	5.52	6.43	6.14	6.25	7.15		5.92	317
	female	10.40	5.20	5.7	6.74	5.21	5.07	7.00		5.68	201
	total	5.78	5.41	5.55	6.54	5.80	5.88	7.12		5.82	518
11. Orissa	male	8.67	5.13	6.83	4.50	6.26	7.29	3.00		6.29	62
	female	9.00	5.75	8.57	1.33	6.00	5.17	3.00		5.92	26
	total	8.80	5.33	7.32	3.44	6.24	6.31	3.00		6.18	88
12. Punjab	male	1.50	6.60	5.43	6.83	7.00	7.25	5.00		6.30	40
	female	5.67	5.71	7.00	4.10	5.08	4.75	8.00		5.56	50
	total	4.00	6.24	6.42	5.13	5.95	6.00	7.00		5.89	90
13. Rajasthan	male	10.00	4.84	11.74	3.61	9.19		3.99		6.71	27
	female	6.00	4.95	4.19	3.03		6.38	6.45		4.93	22
	total	8.67	4.89	7.04	3.41	9.19	6.38	5.63		5.90	49
14. Tamil Nadu	male	4.90	4.57	5.86	7.70	5.65	5.92	4.62		5.64	192
	female	8.92	5.37	7.08	5.79	5.08	7.30	7.28		6.15	189
	total	6.43	4.93	6.47	6.60	5.40	6.84	5.65		5.89	381
15. Uttar Pradesh	male	5.40	6.78	8.67	8.30	11.00	6.20	10.50		8.33	52
	female	7.50	8.83	6.85	5.18	8.64	6.50	7.00		7.11	57
	total	6.33	7.60	7.41	6.67	9.87	6.53	9.33		7.69	109
16. West Bengal	male	5.80	8.25	6.77	6.11	6.73	9.57	15.87		7.70	213
	female	7.12	6.60	10.37	6.95	6.55	10.25	9.76		8.29	135
	total	6.38	7.65	8.39	6.47	6.67	9.74	12.09		7.93	348
17. all- India*	male	5.19	6.28	5.88	6.52	6.80	8.20	7.85		6.58	1548
	female	6.73	6.07	7.02	6.27	6.09	6.99	12.32		6.80	1212
	total	5.85	6.18	6.40	6.41	6.51	7.67	9.99		6.68	2760

\* including smaller states and Union Territories.

sd/

Table (7R) : Number of persons suffering from chronic diseases per 100,000 persons by type of chronic disease, separately by sex for different states and all-India, rural households : NSS 28th round

states		type of chronic diseases																				all types	number of sample ailments
		tuberculosis	leprosy	syphilis	cancer	thyroid trouble or joint trouble	diabetes	mental illness	epilepsy	rheumatic fever	high blood pressure	bronchitis	asthma	peptic ulcer	kidney stone or kidney trouble	arthritis	rheumatism	stroke	piles	others	not recorded		
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)	
1. Andhra Pradesh	m	126	91	13	13	9	17	9	22	34	26	27	277	73	22		189	13	17		774	1752	404
	f	59	23	9	26	9	9	13	13	35	83	13	231	40			177	31	13		867	1651	371
	t	93	57	11	19	9	13	11	17	35	54	20	255	57	11		183	22	15		820	1702	775
2. Assam	m	51	36	14	14	43	65		14	36	79	79	281	194	116		158	14	51	554	144	1943	270
	f	24	9			56	16	16	8	40	16	16	88	129	40	8	153		32	459	72	1182	147
	t	38	23	8	8	49	42	8	11	38	49	49	190	163	80	4	155	8	42	509	110	1584	417
3. Bihar	m	172	63	18	4	14	25	21	39	18	25	25	549	151	25	11	141	11	120	1076	28	2536	720
	f	98	22		4	11	4	18	15	26	15	18	489	91	26	4	219	7	40	1028	11	2146	588
	t	136	43	9	4	13	14	20	27	21	20	21	519	122	25	7	179	9	81	1053	20	2343	1308
4. Gujarat	m	105	8		8	24	24		16	16	24	24	249	8	16	32	8	16	16	81	113	780	97
	f	26	17	9		8	9		8	17		162	34			17			9	42	60	418	49
	t	66	12	4	4	12	17	4	8	12	21	13	207	21	8	25	4	8	8	62	89	605	146
5. Haryana	m	54	24		6	12	18	18	12	90	18	60	409	30	30	26	138		120	968		2043	340
	f	107	7		20	7		13	33	60	20	20	354	27	7	107	87	7	87	1015		1978	296
	t	79	16		13	9	9	16	22	76	19	41	383	28	19	70	114	3	104	991		2012	636
6. Jammu & Kashmir	m	45	12	2	5	17	5	17	13	38	93	14	159	69	66		84	3	43		259	934	542
	f	63	4	2	6	23	17	4	17	64	99	23	91	86	31		90	6	16		321	963	500
	t	54	8	2	5	20	11	11	10	50	96	18	127	77	49		87	5	30		288	948	1042

m = male, f = female, t = total

Table 9R) Contd. : Number of persons suffering from chronic diseases per 100,000 by type of chronic disease separately by sex for different states and all-India, rural households : NSS 28th round

states	type of chronic diseases																				number of sample ailments		
	tuberculosis	leprosy	syphilis	cancer	thyroid trouble or jointre	diabetes	mental illness	epilepsy	rheumatic fever	high blood pressure	bronchitis	asthma	peptic ulcer	kidney stone or kidney trouble	arthritis	rheumatism	stroke	piles	others	not recorded	all types		
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)	
Karnataka	m	50	7	14		14	83	21	14	7	55	21	346	48	7		28	7	26		582	1330	192
	f	43	14		14	14	21	14	14	14	43		257	29			22	14	14		392	919	129
	t	46	11	7	7	14	53	18	14	11	49	18	302	39	4		25	11	21		487	1137	321
Kerala	m	584	21	7	55	41	309	21	110	103	151	288	1057	206	117		2052	21	584		2504	8231	1200
	f	505	26	7	66	85	269		131	236	144	72	1057	66	59	20	2710	20	164		2861	8498	1294
	t	544	23	7	60	64	289	10	121	171	148	178	1057	134	87	10	2389	20	369		2687	8368	2494
Madhya Pradesh	m	59	53	7	11	25	12	6	39	46	12	78	328	35	46	18	198	14	32	406		1425	413
	f	75	22		4	15	34	19	30	37	26	52	75	26	19	22	157	7	11	220		851	228
	t	80	38	4	7	20	24	15	34	42	20	65	205	31	33	20	178	11	22	316		1165	641
Maharashtra	m	127	119	8	12	16	37	20	16	21	12	12	352	114	33	94	41		65		736	1835	449
	f	67	59		8	16	4	12	12	8	28	12	279	51	16	130	28	4	24		639	1397	354
	t	96	88	4	10	16	20	16	14	14	20	12	315	82	24	112	34	2	44		686	1609	803
Orissa	m	53	61	15		23	46	69		76	38	53	274	160	107	23	343	38	61	1463	46	2949	387
	f	60		15	23	15	30	150	15	128	38	45	128	128	53	38	451	45	22	1172	60	2616	348
	t	57	30	15	11	19	38	110	8	102	38	49	200	144	79	30	397	42	42	1317	53	2781	735
Punjab	m	102		45	13	13	13	51	70	26	45	102	774	26	96	6	154	26	204	582		2348	367
	f	130		22		7	50	50	58	65	123	86	468	72	72	14	195	14	86	685		2197	305
	t	115		34	7	10	30	51	65	44	81	95	630	48	85	10	173	20	149	630		2277	672

m = male, f = female, t = total

Table (SR) Contd. : Number of persons suffering from chronic disease per 100,000 persons by type of chronic disease separately by sex for different states and all-India, rural households : NSS 28th round

states		type of chronic diseases																			number of sample ailments		
		tuberculosis	leprosy	syphilis	cancer	thyroid trouble or goitre	diabetes	mental illness	epilepsy	rheumatic fever	high blood pressure	bronchitis	asthma	peptic ulcer	kidney stone or kidney trouble	arthritis	rheumatism	stroke	piles	others	not recorded		
(1)		(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)
13. Rajasthan	m	116	7	7	7	7	29	14	29	36		51	436	37	14	22	88	7	50		468	1425	196
	f	47			8		23	16	16	47	24	8	272	8		16	116	8	15		493	1117	143
	t	83	4	4	7	4	26	15	22	41	11	30	357	23	7	19	102	8	33		480	1276	339
14. Tamil Nadu	m	235	128	13	13		47	14	40		53		242	174	81	13	167	33	53		1147	2453	367
	f	92	59	7	20	7	7	27	33	46	33	13	204	111	7	7	85	13	20		1217	2008	306
	t	163	93	10	17	3	26	20	36	23	43	7	223	142	43	10	125	23	37		1182	2226	673
15. Uttar Pradesh	m	129	31	10	10	16	26	10	29	31	8	34	485	50	21	3	137	10	81		464	1585	177
	f	74	23	3	21	27	35	9	27	65	12	26	345	27	24	6	159	9	15		486	1393	165
	t	103	28	7	15	21	30	10	28	47	8	31	419	39	22	4	147	10	50		474	1493	342
16. West Bengal	m	181	71	21	8	67	86	4	21	29	143	134	634	361	101	34	260	8	160	1137	458	3920	934
	f	58	36	4	4	45	18	15	36	36	138	27	299	98	90	27	368		62	1205	465	2969	665
	t	121	54	13	6	56	54	9	28	32	141	82	472	234	95	30	284	4	113	1170	461	3459	1599
17. All-India	m	144	54	13	11	22	48	17	30	36	41	56	440	115	48	19	228	14	95	204	625	2260	7783
	f	89	25	4	14	21	30	21	27	54	47	25	309	60	26	26	275	12	33	184	647	1947	6480
	t	117	40	8	12	22	39	19	28	45	44	41	376	89	37	22	251	13	65	194	636	2098	14263

\* including smaller states and union territories

m = male, f = female, t = total

Table (VU) : Number of persons suffering from chronic diseases per 100,000 persons by type of chronic disease separately by sex for different states and all-India, urban households : NSS 28th round

states		type of chronic diseases																				all types	number of sample ailments	
		tuberculosis	leprosy	syphilis	cancer	thyroid trouble or goitre	diabetes	mental illness	epilepsy	rheumatic fever	high blood pressure	bronchitis	asthma	peptic ulcer	kidney stone or kidney trouble	arthritis	rheumatism	stroke	piles	others	not recorded			
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)		
Andhra Pradesh	m	98	45	8		8	75		8	8	60	38	150	30	30		38	23	15		466	1100	146	
	f	31	8	8		47	8	8		109	16	186	8	8		47	47	23		637	1191	153		
	t	65	23	8	4	4	61	4	8	4	84	27	168	19	19		42	34	19		550	1143	299	
Assam	m	63			21	63	104			21	229	42	229	231	42		125		63	812	168	2213	106	
	f	27			27	27	81		27	81	109	27	163	27	81		190		1060	136	2063	76		
	t	47			24	47	94		12	47	177	35	201	142	59		153		35	921	154	2148	182	
Bihar	m	230	57		10	67	29	48		29	29	527	67	48		105	29	67	910	10	2262	236		
	f	183			23	34	11	34	11		149	34	561	23	57		57	11	46	778	134	2046	179	
	t	209	31		10	21	42	31	31		83	31	542	47	52		83	21	57	850	21	2162	415	
Gujarat	m	162	12			12	50			12	37		211	12	13	12	12			164	88	797	64	
	f	56				28				14	70	14	98		14	14	14			56	238	616	44	
	t	112	6			7	39			13	53	7	158	7	13	13	13			113	158	712	108	
Haryana	m	121	20				60	40	20	80		623	141	101	20	80		161	1246			2713	135	
	f	138			46	46	-	23	46		92	46	367	69		23	46	23	92	1171			2228	97
	t	129	11		21	21	32	11	43	11	86	21	504	107	54	21	64	11	129	1211			2487	232
Jammu & Kashmir	m	48	13			4	4	4	4	26	126	9	152	126	39		70	9	44		114	792	182	
	f	30	20			5	15	10	5	20	160		90	105	30	10	70		10		262	842	168	
	t	39	17			5	9	7	5	23	141	5	124	116	35	5	70	5	28		182	816	350	

m = male, f = female, t = total

Contd.

Table (7U) Contd. : Number of persons suffering from chronic diseases per 100,000 persons by type of chronic disease separately by sex for different states and all-India, urban households : NSS 28th round

states	type of chronic diseases																				number of sample ailments		
	tuberculosis	leprosy	syphilis	cancer	thyroid trouble or jointre	diabetes	mental illness	epilepsy	rheumatic fever	high blood pressure	bronchitis	asthma	peptic ulcer	kidney stone or kidney trouble	arthritis	rheumatism	stroke	piles	others	not recorded	all types		
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)	
Karnataka	m	128	23			129	23	12		58		453	47		12	35	23	59		288	1290	111	
	f	82	12		12		83	24	12		141		329			48	12			316	1071	91	
	t	105	17		6		106	23	12		100		392	24		6	41	18	29		302	1181	202
Kerala	m	768	16	33		480	33	33	182	434	278	1824	248	181		1584		511		2260	8865	534	
	f	303	34	119	66	321		33	268	372	133	1725	135	116	18	2291	51	269		2565	8884	527	
	t	570	25	75	33	401	16	33	225	403	200	1775	192	149	9	1935	25	391		2411	8874	10611	
Madhya Pradesh	m	135	31			21	41	21	10	10	83		425	62	52	10	31		41	456		1429	138
	f	113		23	45	11	11	23	23	113	11	204	34	23	23	57	11	23	522		1270	112	
	t	125	16		11	32	27	16	16	16	97	5	319	49	38	16	43	5	32	487		1350	250
Maharashtra	m	190	31		26	41	103	10	15	5	180	31	330	72	21	16	21	10	46		463	1570	305
	f	140	18		12	6	91	24	18	6	218	12	339	24	24	24	42		24		588	1610	265
	t	167	25		20	3	97	17	17	6	197	23	334	50	22	20	31	6	36		520	1591	570
Orissa	m	10	113	22		158	68	23			90	248	23	45	23	135	22	68	946	113	2180	97	
	f	53	53	53			264		26	212	26	132	53	106		344	53	26	952	106	2459	93	
	t	73	85	24	12		85	158	12	12	146	12	195	36	73	12	232	36	49	950	110	2312	190
Punjab	m	75	56	19		19	37	19	37		130	130	391	19	56		56	37	540		1621	87	
	f	107	65	21	43	62	22	43	21		215	86	407	43		22	257	107	472		1993	93	
	t	90	60	20	20	40	30	30	30		169	110	398	30	30	10	149	70	508		1794	180	

Table (9U) Contd. Number of persons suffering from chronic diseases per 100,000 persons by type of chronic disease separately by sex for different states and all-India, urban households ; NSS 28th round

states	type of chronic diseases																				all types	number of sample ailments	
	tuberculosis	leprosy	syphilis	cancer	thyroid trouble or goitre	diabetes	mental illness	epilepsy	rheumatic fever	high blood pressure	bronchitis	asthma	peptic ulcer	kidney stone or kidney trouble	arthritis	rheumatism	stroke	piles	others	not recorded			
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)	
Rajasthan	m	89	30	15	14		29	15		30		93		45				15		459	834	55	
	f	64				36	48	48	16	16	48	33	114	33			65		32		384	937	57
	t	77	15	8	7	17	39	31	8	8	39	16	103	16	23		31		23		423	884	112
Tamil Nadu	m	195	55	21	7	35	138	14	7	6	76	14	270	131	21		21	14	27		997	2049	296
	f	86	22	21	14		29	14	14	7	138	22	223	57	7	15	79		14		1165	1927	268
	t	142	39	21	11	18	85	14	10	7	106	18	247	95	14	7	49	7	21		1079	1990	564
Uttar Pradesh	m	184	32	5	5		16	5	11	11	16	38	276	32	54		54	5	22	416		1182	77
	f	143	13	20	26	7	26	19	20	32	78	26	176	33	72	7	84	13	7	449		1251	69
	t	165	23	12	15	3	21	12	15	21	44	32	230	32	62	3	68	9	15	431		1213	146
West Bengal	m	111	22		7	37	193	7	22	44	222	163	651	229	74	82	193	37	282	1052	496	3924	529
	f	47	19		19	28	38	9	47	104	348	132	301	122	75	132	319	9	103	1449	601	3902	414
	t	83	21		12	33	125	8	33	70	278	149	497	182	74	104	248	25	203	1227	542	3914	943
All-India	m	169	34	5	8	15	105	13	16	21	108	48	397	86	45	13	113	14	81	130	582	2003	3373
	f	102	14	7	20	18	52	22	18	33	159	38	308	44	35	22	182	12	38	127	680	1931	2912
	t	137	25	6	14	16	80	18	17	26	132	43	355	66	40	17	146	13	61	128	629	1962	6285

\* including smaller states and union territories.

m = male, f = female, t = total

Table (18R): Number of persons suffering from chronic diseases per 100,000 persons by age separately by sex for different states and all-India, rural households : NSS 28th round.

RURAL

states		age (years)							number of sam- ple cases
		0-14	15-25	25-44	45-59	60 & above	not re- corded	all ages	
		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
1. Andhra Pra- desh	male	241	388	1731	5346	8114	—	1752	404
	female	284	340	2618	4045	4633	—	1651	371
	total	261	611	2162	4666	6770	—	1702	775
2. Assam	male	265	650	2520	6770	9184	—	1943	270
	female	172	753	2245	3771	3868	—	1182	147
	total	216	702	2396	5379	7241	—	1584	417
3. Bihar	male	455	1084	3490	7204	9223	—	2536	720
	female	367	1088	3067	5784	5727	—	2146	588
	total	411	1084	3273	6466	7460	—	2343	1308
4. Gujarat	male	202	264	953	2035	4117	—	780	97
	female	—	97	593	1343	2182	—	418	49
	total	105	184	777	1687	3163	—	605	146
5. Haryana	male	549	941	2359	6602	7488	—	2043	340
	female	392	684	4040	5424	5119	—	1978	296
	total	476	914	3214	6034	6577	—	2012	636
6. Jammu & Kashmir	male	134	321	173	2572	4093	—	934	542
	female	96	371	1867	2404	3043	—	963	500
	total	116	344	1443	2491	3684	—	948	1042
7. Karnataka	male	401	285	1168	4069	6375	—	1330	192
	female	417	272	1481	1593	2831	—	919	129
	total	409	288	1325	2840	4716	—	1137	321
8. Kerala	male	2054	2093	10325	22301	35649	—	8231	1200
	female	1805	3515	11457	24547	26859	—	8498	1294
	total	1931	2839	10927	23439	31099	—	8368	2494
9. Madhya Pradesh	male	313	553	1879	3891	5757	—	1425	413
	female	78	461	1329	2448	2088	—	851	228
	total	200	512	1616	3141	3891	—	1165	641
10. Maharashtra	male	438	698	2271	4612	7833	—	1835	449
	female	276	355	2552	3869	3343	—	1397	354
	total	361	506	2413	4221	5652	—	1609	803

Table (8R): contd. Number of persons suffering from chronic diseases per 100,000 persons by age separately by sex for different states and all-India, rural households NSS 28th round.

RURAL

states		age (years)							number of sam- ple cases
		0-14	15-25	25-44	45-59	60 & above	not re- corded	all ages	
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	
11. Orissa	male	651	1236	3719	7496	11341	-	2949	387
	female	626	1280	3456	7561	7143	-	2616	348
	total	638	1259	3589	7530	9403	-	2781	735
12. Punjab	male	247	759	2459	6689	9717	-	2348	367
	female	269	900	3503	5772	6708	-	2197	305
	total	257	827	2970	6238	8458	-	2277	672
13. Rajasthan	male	246	540	1534	4912	5803	-	1425	196
	female	187	570	1940	2714	2728	-	1117	143
	total	218	554	1731	3770	4345	-	1276	339
14. Tamil Nadu	male	509	1088	2821	6486	8538	-	2453	367
	female	431	1090	3387	4323	3804	-	2008	306
	total	470	1089	3110	5382	6306	-	2226	673
15. Uttar Pradesh	male	263	485	1835	4498	6082	-	1585	177
	female	248	856	1963	3533	3338	-	1393	165
	total	256	661	1897	4009	4848	-	1493	342
16. West Bengal	male	1028	1260	4692	11763	18549	-	3920	934
	female	532	1173	5526	8224	848	-	2969	665
	total	786	1221	5088	9959	13743	-	3459	1599
17. All-India*	male	474	793	2703	6373	9248	13015	2260	7783
	female	361	917	2966	5109	5340	16625	1929	6400
	total	421	855	2833	5726	7404	14582	2098	14263

\* including smaller states and Union Territories.

Table (8U) : Number of persons suffering from chronic diseases per 100,000 persons by age separately by sex for different states and all-India, urban households NSS 28th round.

URBAN

states		age (years)							number of sam- ple cases
		0-14	15-24	25-44	45-59	60 & above	all ages		
		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
1. Andhra Pradesh	male	268	299	1521	3177	4747	1100	146	
	female	193	565	2259	2574	3225	1191	153	
	total	231	426	1868	2864	3837	1143	299	
2. Assam	male	232	670	2845	8379	9231	2213	106	
	female	194	992	4222	419	5128	2063	76	
	total	214	810	3372	7604	7464	2148	182	
3. Bihar	male	267	1249	2679	4797	15726	2262	236	
	female	270	605	3451	5385	7589	2046	179	
	total	270	985	3022	5069	11583	2162	415	
4. Gujarat	male	-	281	1050	1636	6552	797	64	
	female	145	137	1245	1232	1728	616	44	
	total	69	216	1141	1442	4080	712	108	
5. Haryana	male	461	746	3649	7806	12037	2713	135	
	female	277	905	3768	5870	8929	2228	97	
	total	373	1818	3704	6852	10796	2487	232	
6. Jammu & Kashmir	male	116	226	856	2605	5465	792	182	
	female	94	390	1363	2576	3942	842	168	
	total	106	302	1084	2591	4822	816	350	
7. Karnataka	male	444	375	1490	3726	5975	1290	111	
	female	272	280	1393	3012	4388	1071	91	
	total	359	328	1442	3356	5137	1181	202	
8. Kerala	male	3684	2256	10721	22137	36613	8865	534	
	female	2643	2253	11976	24232	25537	8884	527	
	total	3186	2256	11360	23175	30393	8874	1061	
9. Madhya Pradesh	male	179	311	1301	5185	8686	1429	138	
	female	135	366	2514	3514	3426	1270	112	
	total	157	336	1868	4387	5980	1350	250	
10. Maharashtra	male	289	655	1744	5465	6241	1570	305	
	female	360	741	1949	4435	6544	1610	265	
	total	321	692	1831	5000	6399	1591	570	

Table (S.U): contd. Number of persons suffering from chronic diseases per 100,000 persons by age separately by sex for different states and all-India, urban households : NCS 28th round.

URBAN

states	(1)	age (years)						number of sam- ple cases (8)
		0-14	15-24	25-44	45-59	60 & above ages	all	
		(2)	(3)	(4)	(5)	(6)	(7)	
11. Orissa	male	619	1012	2732	6720	8297	2180	97
	female	260	732	4493	6170	6719	2459	93
	total	452	891	3510	6439	7469	2312	190
12. Punjab	male	473	416	1724	4097	8036	1621	87
	female	334	607	2342	7042	7326	1993	93
	total	405	502	1998	5513	7718	1794	180
13. Rajasthan	male	191	280	1343	2256	3243	834	55
	female	313	524	2263	1424	994	937	57
	total	251	388	1778	1639	2061	884	112
14. Tamil Nadu	male	941	489	2080	5412	8637	2049	296
	female	349	1040	3086	4059	6004	1927	268
	total	647	764	2551	4737	7252	1990	564
15. Uttar Pradesh	male	222	637	1363	3361	4592	1182	77
	female	158	683	2156	2619	4357	1251	69
	total	192	657	1718	3017	4486	1213	146
16. West Bengal	male	1596	1217	3565	8910	18640	3924	529
	female	787	1367	5651	10446	12016	3902	414
	total	1207	1284	4379	9531	15316	3914	943
17. All India*	male	591	656	2232	5662	9450	2003	3373
	female	379	763	2982	4954	6388	1931	2912
	total	488	704	2571	5321	7893	1969	6285

\* including smaller states and Union Territories.